



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: C12N 15/57, C07K 14/47, C07K 16/18, C07K 19/00, C12N 1/21, C12N 5/10, C12N 9/64, C12N 15/12, C12N 15/62, C12N 15/85, C12Q 1/37, G01N 33/68	A2	(11) International Publication Number: WO 00/17369 (43) International Publication Date: 30 March 2000 (30.03.2000)
--	----	--

(21) International Application Number: PCT/US99/20881

(22) International Filing Date: 23 September 1999 (23.09.1999)

(30) Priority Data:
60/101,594 24 September 1998 (24.09.1998) US

(60) Parent Application or Grant

PHARMACIA & UPJOHN COMPANY []; () GURNEY,
Mark, E. []; () BIENKOWSKI, Michael, Jerome [];
() HEINRIKSON, Robert, Leroy []; () PARODI, Luis, A. [];
() YAN, Riqiang []; () GURNEY, Mark, E. [];
() BIENKOWSKI, Michael, Jerome []; () HEINRIKSON,
Robert, Leroy []; () PARODI, Luis, A. []; () YAN, Riqiang
[]; () WOOTTON, Thomas, A. []

Published

(54) Title: ALZHEIMER'S DISEASE SECRETASE

(54) Titre: SECRETASE DE LA MALADIE D'ALZHEIMER

(57) Abstract

The present invention provides the enzyme and enzymatic procedures for cleaving the 'beta' secretase cleavage site of the APP protein and associated nucleic acids, peptides, vectors, cells and cell isolates and assays.

(57) Abrégé

La présente invention porte sur l'enzyme et les procédures enzymatiques de clivage du site de clivage de la 'beta' sécrétase de la protéine APP et des acides nucléiques, des peptides, des vecteurs, des cellules et des isolats cellulaires associés, et sur des dosages.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷: C12N 15/57, 15/62, 15/85, 5/10, 9/64, C07K 19/00, 14/47, C12N 15/12, C07K 16/18, C12Q 1/37, G01N 33/68, C12N 1/21	A2	(11) International Publication Number: WO 00/17369 (43) International Publication Date: 30 March 2000 (30.03.00)
(21) International Application Number: PCT/US99/20881 (22) International Filing Date: 23 September 1999 (23.09.99) (30) Priority Data: 60/101,594 24 September 1998 (24.09.98) US (71) Applicant (for all designated States except US): PHARMACIA & UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): GURNEY, Mark, E. [US/US]; 910 Rosewood Avenue, S.E., Grand Rapids, MI 49506 (US). BIENKOWSKI, Michael, Jerome [US/US]; 3431 Hollow Wood, Portage, MI 49024 (US). HEINRIK- SON, Robert, Leroy [US/US]; 81 South Lake Doster Drive, Plainwell, MI 49080 (US). PARODI, Luis, A. [US/SE]; Grevgafan 24, S-115 43 Stockholm (SE). YAN, Riqiang [US/US]; 5026 Queen Victoria Street, Kalamazoo, MI 49009 (US).	(74) Agent: WOOTTON, Thomas, A.; Pharmacia & Upjohn Com- pany, Intellectual Property Legal Services, 301 Henrietta Street, Kalamazoo, MI 49001 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(54) Title: ALZHEIMER'S DISEASE SECRETASE (57) Abstract The present invention provides the enzyme and enzymatic procedures for cleaving the β secretase cleavage site of the APP protein and associated nucleic acids, peptides, vectors, cells and cell isolates and assays.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
RE	Reunion	LR	Liberia	SG	Singapore		

Description

5

10

15

20

25

30

35

40

45

50

55

Alzheimer's Disease Secretase

FIELD OF THE INVENTION

The present invention related to the field of Alzheimer's Disease, APP, amyloid beta peptide, and human aspartyl proteases as well as a method for the identification of agents that modulate the activity of these polypeptides.

BACKGROUND OF THE INVENTION

Alzheimer's disease (AD) causes progressive dementia with consequent formation of amyloid plaques, neurofibrillary tangles, gliosis and neuronal loss. The disease occurs in both genetic and sporadic forms whose clinical course and pathological features are quite similar. Three genes have been discovered to date which when mutated cause an autosomal dominant form of Alzheimer's disease. These encode the amyloid protein precursor (APP) and two related proteins, presenilin-1 (PS1) and presenilin-2 (PS2), which as their names suggest are both structurally and functionally related. Mutations in any of the three enhance proteolytic processing of APP via an intracellular pathway that produces amyloid beta peptide or the A β peptide (or sometimes here as Abeta), a 40-42 amino acid long peptide that is the primary component of amyloid plaque in AD. Dysregulation of intracellular pathways for proteolytic processing may be central to the pathophysiology of AD. In the case of plaque formation, mutations in APP, PS1 or PS2 consistently alter the proteolytic processing of APP so as to enhance formation of A β 1-42, a form of the A β peptide which seems to be particularly amyloidogenic, and thus very important in AD. Different forms of APP range in size from 695-770 amino acids, localize to the cell surface, and have a single C-terminal transmembrane domain. The Abeta peptide is derived from a region of APP adjacent to and containing a portion of the transmembrane domain. Normally, processing of APP at the α -secretase site cleaves the midregion of the A β sequence adjacent to the membrane and releases the soluble, extracellular domain of APP from the cell surface. This α -secretase APP processing, creates soluble APP- α , and it is normal and not thought to contribute to AD.

Pathological processing of APP at the β - and γ -secretase sites produces a very different result than processing at the α site. Sequential processing at the β - and γ -secretase sites releases the A β peptide, a peptide possibly very important in AD pathogenesis. Processing at the β - and γ -secretase sites can occur in both the endoplasmic reticulum (in neurons) and in the endosomal/lysosomal pathway after reinternalization of cell surface

5 APP (in all cells). Despite intense efforts, for 10 years or more, to identify the enzymes responsible for processing APP at the β and γ sites, to produce the A β peptide, those proteases remained unknown until this disclosure. Here, for the first time, we report the identification and characterization of the β secretase enzyme. We disclose some known and
10 some novel human aspartic proteases that can act as β -secretase proteases and, for the first time, we explain the role these proteases have in AD. We describe regions in the proteases critical for their unique function and for the first time characterize their substrate. This is the first description of expressed isolated purified active protein of this type, assays that use
15 the protein, in addition to the identification and creation of useful cell lines and inhibitors.

10 SUMMARY OF THE INVENTION

Here we disclose a number of variants of the asp2 gene and peptide.

20 Any isolated or purified nucleic acid polynucleotide that codes for a protease capable of cleaving the beta (β) secretase cleavage site of APP that contains two or more sets of special nucleic acids, where the special nucleic acids are separated by nucleic acids
25 that code for about 100 to 300 amino acid positions, where the amino acids in those positions may be any amino acids, where the first set of special nucleic acids consists of the nucleic acids that code for the peptide DTG, where the first nucleic acid of the first special set of nucleic acids is, the first special nucleic acid, and where the second set of nucleic
30 acids code for either the peptide DSG or DTG, where the last nucleic acid of the second set of nucleic acids is the last special nucleic acid, with the proviso that the nucleic acids disclosed in SEQ ID NO. 1 and SEQ. ID NO. 5 are not included. The nucleic acid
35 polynucleotide of claim 1 where the two sets of nucleic acids are separated by nucleic acids that code for about 125 to 222 amino acid positions, which may be any amino acids. The nucleic acid polynucleotide of claim 2 that code for about 150 to 172 amino acid positions,
25 which may be any amino acids. The nucleic acid polynucleotide of claim that code for about 172 amino acid positions, which may be any amino acids. The nucleic acid polynucleotide of claim 4 where the nucleotides are described in SEQ. ID. NO. 3 The
40 nucleic acid polynucleotide of claim 2 where the two sets of nucleic acids are separated by nucleic acids that code for about 150 to 196 amino acid positions. The nucleic acid polynucleotide of claim 6 where the two sets of nucleotides are separated by nucleic acids
30 that code for about 196 amino acids (positions). The nucleic acid polynucleotide of claim 7 where the two sets of nucleic acids are separated by the same nucleic acid sequences that
50 separate the same set of special nucleic acids in SEQ. ID. NO. 5. The nucleic acid

5 polynucleotide of claim 4 where the two sets of nucleic acids are separated by nucleic acids
that code for about 150 to 190, amino acid (positions). The nucleic acid polynucleotide of
claim 9 where the two sets of nucleotides are separated by nucleic acids that code for about
190 amino acids (positions). The nucleic acid polynucleotide of claim 10 where the two
10 5 sets of nucleotides are separated by the same nucleic acid sequences that separate the same
set of special nucleotides in SEQ. ID. NO. 1. Claims 1-11 where the first nucleic acid of
the first special set of amino acids, that is, the first special nucleic acid, is operably linked
to any codon where the nucleic acids of that codon codes for any peptide comprising from 1
15 to 10,000 amino acid (positions). The nucleic acid polynucleotide of claims 1-12 where the
first special nucleic acid is operably linked to nucleic acid polymers that code for any
peptide selected from the group consisting of: any any reporter proteins or proteins which
facilitate purification. The nucleic acid polynucleotide of claims 1-13 where the first special
20 nucleic acid is operably linked to nucleic acid polymers that code for any peptide selected
from the group consisting of: immunoglobulin-heavy chain, maltose binding protein,
glutathion S transfection, Green Fluorescent protein, and ubiquitin. Claims 1-14 where the
25 last nucleic acid of the second set of special amino acids, that is, the last special nucleic
acid, is operably linked to nucleic acid polymers that code for any peptide comprising any
amino acids from 1 to 10,000 amino acids. Claims 1-15 where the last special nucleic acid
is operably linked to any codon linked to nucleic acid polymers that code for any peptide
30 selected from the group consisting of: any reporter proteins or proteins which facilitate
purification. The nucleic acid polynucleotide of claims 1-16 where the first special nucleic
acid is operably linked to nucleic acid polymers that code for any peptide selected from the
group consisting of: immunoglobulin-heavy chain, maltose binding protein, glutathion S
35 transfection, Green Fluorescent protein, and ubiquitin.

25 Any isolated or purified nucleic acid polynucleotide that codes for a protease
capable of cleaving the beta secretase cleavage site of APP that contains two or more sets of
40 special nucleic acids, where the special nucleic acids are separated by nucleic acids that
code for about 100 to 300 amino acid positions, where the amino acids in those positions
may be any amino acids, where the first set of special nucleic acids consists of the nucleic
45 acids that code for DTG, where the first nucleic acid of the first special set of nucleic acids
is, the first special nucleic acid, and where the second set of nucleic acids code for either
DSG or DTG, where the last nucleic acid of the second set of special nucleic acids is the
50 last special nucleic acid, where the first special nucleic acid is operably linked to nucleic

5 acids that code for any number of amino acids from zero to 81 amino acids and where each
of those codons may code for any amino acid. The nucleic acid polynucleotide of claim 18
10 , where the first special nucleic acid is operably linked to nucleic acids that code for any
number of from 64 to 77 amino acids where each codon may code for any amino acid. The
5 nucleic acid polynucleotide of claim 19 , where the first special nucleic acid is operably
linked to nucleic acids that code for 71 amino acids. The nucleic acid polynucleotide of
claim 20, where the first special nucleic acid is operably linked to 71 amino acids and
15 where the first of those 71 amino acids is the amino acid T. The nucleic acid
polynucleotide of claim 21, where the polynucleotide comprises a sequence that is at least
10 95% identical to SEQ. ID. (Example 11). The nucleic acid polynucleotide of claim 22,
where the complete polynucleotide comprises SEQ. ID. (Example 11). The nucleic acid
polynucleotide of claim 18 , where the first special nucleic acid is operably linked to nucleic
20 acids that code for any number of from 40 to 54 amino acids where each codon may code
for any amino acid. The nucleic acid polynucleotide of claim 24, where the first special
nucleic acid is operably linked to nucleic acids that code for 47 amino acids. The nucleic
25 acid polynucleotide of claim 20, where the first special nucleic acid is operably linked to 47
codons where the first those 47 amino acids is the amino acid E. The nucleic acid
polynucleotide of claim 21, where the polynucleotide comprises a sequence that is at least
30 95% identical to SEQ. ID. (Example 10). The nucleic acid polynucleotide of claim 22,
where the complete polynucleotide comprises SEQ. ID. (Example 10).

Any isolated or purified nucleic acid polynucleotide that codes for a protease
35 capable of cleaving the beta (β) secretase cleavage site of APP that contains two or more
sets of special nucleic acids, where the special nucleic acids are separated by nucleic acids
that code for about 100 to 300 amino acid positions, where the amino acids in those
25 positions may be any amino acids, where the first set of special nucleic acids consists of the
nucleic acids that code for the peptide DTG, where the first nucleic acid of the first special
40 set of amino acids is, the first special nucleic acid, and where the second set of special
nucleic acids code for either the peptide DSG or DTG, where the last nucleic acid of the
second set of special nucleic acids, the last special nucleic acid, is operably linked to
45 nucleic acids that code for any number of codons from 50 to 170 codons. The nucleic acid
polynucleotide of claim 29 where the last special nucleic acid is operably linked to nucleic
30 acids comprising from 100 to 170 codons. The nucleic acid polynucleotide of claim 30
where the last special nucleic acid is operably linked to nucleic acids comprising from 142
50

5 to 163 codons. The nucleic acid polynucleotide of claim 31 where the last special nucleic acid is operably linked to nucleic acids comprising about 142 codons. The nucleic acid polynucleotide of claim 32 where the polynucleotide comprises a sequence that is at least 95% identical to SEQ. ID. (Example 9 or 10). The nucleic acid polynucleotide of claim 10 5 33, where the complete polynucleotide comprises SEQ. ID. (Example 9 or 10). The nucleic acid polynucleotide of claim 31 where the last special nucleic acid is operably linked to nucleic acids comprising about 163 codons. The nucleic acid polynucleotide of claim 35 where the polynucleotide comprises a sequence that is at least 95% identical to SEQ. ID. (Example 9 or 10). The nucleic acid polynucleotide of claim 36, where the 15 complete polynucleotide comprises SEQ. ID. (Example 9 or 10). The nucleic acid polynucleotide of claim 31 where the last special nucleic acid is operably linked to nucleic acids comprising about 170 codons. Claims 1-38 where the second set of special nucleic acids code for the peptide DSG, and optionally the first set of nucleic acid polynucleotide is operably linked to a peptide purification tag. Claims 1-39 where the nucleic acid 20 polynucleotide is operably linked to a peptide purification tag which is six histidine. Claims 1-40 where the first set of special nucleic acids are on one polynucleotide and the second set of special nucleic acids are on a second polynucleotide, where both first and second polynucleotides have at least 50 codons. Claims 1-40 where the first set of special nucleic acids are on one polynucleotide and the second set of special nucleic acids are on a 30 second polynucleotide, where both first and second polynucleotides have at least 50 codons where both said polynucleotides are in the same solution. A vector which contains a polynucleotide described in claims 1-42. A cell or cell line which contains a polynucleotide described in claims 1-42.

Any isolated or purified peptide or protein comprising an amino acid polymer that is 25 a protease capable of cleaving the beta (β) secretase cleavage site of APP that contains two or more sets of special amino acids, where the special amino acids are separated by about 100 to 300 amino acid positions, where each amino acid position can be any amino acid, where the first set of special amino acids consists of the peptide DTG, where the first amino acid of the first special set of amino acids is, the first special amino acid, where the second 45 set of amino acids is selected from the peptide comprising either DSG or DTG, where the last amino acid of the second set of special amino acids is the last special amino acid, with the proviso that the proteases disclosed in SEQ ID NO. 2 and SEQ. ID NO. 6 are not 50 included. The amino acid polypeptide of claim 45 where the two sets of amino acids are

5 separated by about 125 to 222 amino acid positions where in each position it may be any amino acid. The amino acid polypeptide of claim 46 where the two sets of amino acids are separated by about 150 to 172 amino acids. The amino acid polypeptide of claim 47 where the two sets of amino acids are separated by about 172 amino acids. The amino acid
10 5 polypeptide of claim 48 where the protease is described in SEQ. ID. NO. 4 The amino acid polypeptide of claim 46 where the two sets of amino acids are separated by about 150 to 196 amino acids. The amino acid polypeptide of claim 50 where the two sets of amino acids are separated by about 196 amino acids. The amino acid polypeptide of claim 51
15 where the two sets of amino acids are separated by the same amino acid sequences that separate the same set of special amino acids in SEQ. ID. NO. 6. The amino acid
20 polypeptide of claim 46 where the two sets of amino acids are separated by about 150 to 190, amino acids. The amino acid polypeptide of claim 53 where the two sets of nucleotides are separated by about 190 amino acids. The amino acid polypeptide of claim 54 where the two sets of nucleotides are separated by the same amino acid sequences that
25 15 separate the same set of special amino acids in SEQ. ID. NO. 2. Claims 45-55 where the first amino acid of the first special set of amino acids, that is, the first special amino acid, is operably linked to any peptide comprising from 1 to 10,000 amino acids. The amino acid polypeptide of claims 45-56 where the first special amino acid is operably linked to any peptide selected from the group consisting of: any any reporter proteins or proteins which
30 20 facilitate purification. The amino acid polypeptide of claims 45-57 where the first special amino acid is operably linked to any peptide selected from the group consisting of: immunoglobulin-heavy chain, maltose binding protein, glutathion S transfection, Green
35 Fluorescent protein, and ubiquitin. Claims 45-58, where the last amino acid of the second set of special amino acids, that is, the last special amino acid, is operably linked to any
25 peptide comprising any amino acids from 1 to 10,000 amino acids. Claims 45-59 where the
40 last special amino acid is operably linked any peptide selected from the group consisting of any reporter proteins or proteins which facilitate purification. The amino acid polypeptide of claims 45-60 where the first special amino acid is operably linked to any peptide selected from the group consisting of: immunoglobulin-heavy chain, maltose binding protein,
45 30 glutathion S transfection, Green Fluorescent protein, and ubiquitin.

Any isolated or purified peptide or protein comprising an amino acid polypeptide that codes for a protease capable of cleaving the beta secretase cleavage site of APP that
50 contains two or more sets of special amino acids, where the special amino acids are

5 separated by about 100 to 300 amino acid positions, where each amino acid in each position
can be any amino acid, where the first set of special amino acids consists of the amino acids
DTG, where the first amino acid of the first special set of amino acids is, the first special
amino acid, D, and where the second set of amino acids is either DSG or DTG, where the
10 last amino acid of the second set of special amino acids is the last special amino acid, G,
where the first special amino acid is operably linked to amino acids that code for any
number of amino acids from zero to 81 amino acid positions where in each position it may
be any amino acid. The amino acid polypeptide of claim 62, where the first special amino
15 acid is operably linked to a peptide from about 64 to 77 amino acids positions where each
amino acid position may be any amino acid. The amino acid polypeptide of claim 63,
where the first special amino acid is operably linked to a peptide of 71 amino acids. The
amino acid polypeptide of claim 64, where the first special amino acid is operably linked to
20 71 amino acids and the first of those 71 amino acids is the amino acid T. The amino acid
polypeptide of claim 65, where the polypeptide comprises a sequence that is at least 95%
15 identical to SEQ. ID. (Example 11). The amino acid polypeptide of claim 66, where the
complete polypeptide comprises SEQ. ID. (Example 11). The amino acid polypeptide of
claim 62, where the first special amino acid is operably linked to any number of from 40 to
54 amino acids (positions) where each amino acid position may be any amino acid. The
30 amino acid polypeptide of claim 68, where the first special amino acid is operably linked to
amino acids that code for a peptide of 47 amino acids. The amino acid polypeptide of claim
69, where the first special amino acid is operably linked to a 47 amino acid peptide where
the first those 47 amino acids is the amino acid E. The amino acid polypeptide of claim 70,
35 where the polypeptide comprises a sequence that is at least 95% identical to SEQ. ID.
(Example 10). The amino acid polypeptide where the polypeptide comprises Example 10).

25 Any isolated or purified amino acid polypeptide that is a protease capable of
40 cleaving the beta (β) secretase cleavage site of APP that contains two or more sets of
special amino acids, where the special amino acids are separated by about 100 to 300 amino
acid positions, where each amino acid in each position can be any amino acid, where the
45 first set of special amino acids consists of the amino acids that code for DTG, where the
30 first amino acid of the first special set of amino acids is, the first special amino acid, D, and
where the second set of amino acids are either DSG or DTG, where the last amino acid of
the second set of special amino acids is the last special amino acid, G, which is operably
50 linked to any number of amino acids from 50 to 170 amino acids, which may be any amino

acids. The amino acid polypeptide of claim 73 where the last special amino acid is operably linked to a peptide of about 100 to 170 amino acids. The amino acid polypeptide of claim 74 where the last special amino acid is operably linked to a peptide of about 142 to 163 amino acids. The amino acid polypeptide of claim 75 where the last special amino acid is operably linked to a peptide of about 142 amino acids. The amino acid polypeptide of claim 76 where the polypeptide comprises a sequence that is at least 95% identical to SEQ. ID. (Example 9 or 10). The amino acid polypeptide of claim 75 where the last special amino acid is operably linked to a peptide of about 163 amino acids. The amino acid polypeptide of claim 79 where the polypeptide comprises a sequence that is at least 95% identical to SEQ. ID. (Example 9 or 10). The amino acid polypeptide of claim 79, where the complete polypeptide comprises SEQ. ID. (Example 9 or 10). The amino acid polypeptide of claim 74 where the last special amino acid is operably linked to a peptide of about 170 amino acids. Claim 46-81 where the second set of special amino acids is comprised of the peptide with the amino acid sequence DSG. Claims 45-82 where the amino acid polypeptide is operably linked to a peptide purification tag. Claims 45-83 where the amino acid polypeptide is operably linked to a peptide purification tag which is six histidine. Claims 45-84 where the first set of special amino acids are on one polypeptide and the second set of special amino acids are on a second polypeptide, where both first and second polypeptide have at least 50 amino acids, which may be any amino acids. Claims 45-84 where the first set of special amino acids are on one polypeptide and the second set of special amino acids are on a second polypeptide, where both first and second polypeptides have at least 50 amino acids where both said polypeptides are in the same vessel. A vector which contains a polypeptide described in claims 45-86. A cell or cell line which contains a polynucleotide described in claims 45-87. The process of making any of the polynucleotides, vectors, or cells of claims 1-44. The process of making any of the polypeptides, vectors or cells of claims 45-88. Any of the polynucleotides, polypeptides, vectors, cells or cell lines described in claims 1-88 made from the processes described in claims 89 and 90.

Any isolated or purified peptide or protein comprising an amino acid polypeptide that codes for a protease capable of cleaving the beta secretase cleavage site of APP that contains two or more sets of special amino acids, where the special amino acids are separated by about 100 to 300 amino acid positions, where each amino acid in each position can be any amino acid, where the first set of special amino acids consists of the amino acids

5 DTG, where the first amino acid of the first special set of amino acids is, the first special amino acid, D, and where the second set of amino acids is either DSG or DTG, where the last amino acid of the second set of special amino acids is the last special amino acid, G, where the first special amino acid is operably linked to amino acids that code for any
10 5 number of amino acids from zero to 81 amino acid positions where in each position it may be any amino acid.

15 The amino acid polypeptide of claim 62, where the first special amino acid is operably linked to a peptide from about 30 to 77 amino acids positions where each amino acid position may be any amino acid. The amino acid polypeptide of claim 63, where the
20 10 first special amino acid is operably linked to a peptide of 35, 47, 71, or 77 amino acids.

25 The amino acid polypeptide of claim 63, where the first special amino acid is operably linked to the same corresponding peptides from SEQ. ID. NO. 3 that are 35, 47, 71, or 77 peptides in length, beginning counting with the amino acids on the first special sequence, DTG, towards the N-terminal of SEQ. ID. NO. 3.

30 15 The amino acid polypeptide of claim 65, where the polypeptide comprises a sequence that is at least 95% identical to the same corresponding amino acids in SEQ. ID. NO. 4, that is, identical to that portion of the sequences in SEQ. ID. NO. 4, including all the sequences from both the first and or the second special nucleic acids, toward the N-terminal, through and including 71, 47, 35 amino acids before the first special amino acids.
20 (Examples 10 and 11).

35 The amino acid polypeptide of claim 65, where the complete polypeptide comprises the peptide of 71 amino acids, where the first of the amino acid is T and the second is Q. The nucleic acid polynucleotide of claim 21, where the polynucleotide comprises a sequence that is at least 95% identical to the same corresponding amino acids in SEQ. ID.
25 30 NO. 3, that is, identical to the sequences in SEQ. ID. NO. 3 including the sequences from both the first and or the second special nucleic acids, toward the N-Terminal, through and including 71 amino acids, see Example 10, beginning from the DTG site and including the nucleotides from that code for 71 amino acids).

45 30 The nucleic acid polynucleotide of claim 22, where the complete polynucleotide comprises identical to the same corresponding amino acids in SEQ. ID. NO. 3, that is, identical to the sequences in SEQ. ID. NO. 3 including the sequences from both the first and or the second special nucleic acids, toward the N-Terminal, through and including 71
50

amino acids, see Example 10, beginning from the DTG site and including the nucleotides from that code for 71 amino acids).

The nucleic acid polynucleotide of claim 18, where the first special nucleic acid is operably linked to nucleic acids that code for any number of from about 30 to 54 amino acids where each codon may code for any amino acid.

The nucleic acid polynucleotide of claim 20, where the first special nucleic acid is operably linked to 47 codons where the first those 35 or 47 amino acids is the amino acid E or G.

The nucleic acid polynucleotide of claim 21, where the polynucleotide comprises a sequence that is at least 95% identical to the same corresponding amino acids in SEQ. ID. NO. 3, that is, identical to that portion of the sequences in SEQ. ID. NO. 3 including the sequences from both the first and or the second special nucleic acids, toward the N-Terminal, through and including 35 or 47 amino acids, see Example 11 for the 47 example, beginning from the DTG site and including the nucleotides from that code for the previous 35 or 47 amino acids before the DTG site). The nucleic acid polynucleotide of claim 22, where the polynucleotide comprises identical to the same corresponding amino acids in SEQ. ID. NO. 3, that is, identical to the sequences in SEQ. ID. NO. 3 including the sequences from both the first and or the second special nucleic acids, toward the N-Terminal, through and including 35 or 47 amino acids, see Example 11 for the 47 example, beginning from the DTG site and including the nucleotides from that code for the previous 35 or 47 amino acids before the DTG site).

An isolated nucleic acid molecule comprising a polynucleotide, said polynucleotide encoding a Hu-Asp polypeptide and having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:

(a) a nucleotide sequence encoding a Hu-Asp polypeptide selected from the group consisting of Hu-Asp1, Hu-Asp2(a), and Hu-Asp2(b), wherein said Hu-Asp1, Hu-Asp2(a) and Hu-Asp2(b) polypeptides have the complete amino acid sequence of SEQ ID No. 2, SEQ ID No. 4, and SEQ ID No. 6, respectively; and

(b) a nucleotide sequence complementary to the nucleotide sequence of (a).

The nucleic acid molecule of claim 92, wherein said Hu-Asp polypeptide is Hu-Asp1, and said polynucleotide molecule of 1(a) comprises the nucleotide sequence of SEQ ID No. 1. The nucleic acid molecule of claim 92, wherein said Hu-Asp polypeptide is Hu-

5 Asp2(a), and said polynucleotide molecule of 1(a) comprises the nucleotide sequence of
SEQ ID No. 4. The nucleic acid molecule of claim 92, wherein said Hu-Asp polypeptide is
Hu-Asp2(b), and said polynucleotide molecule of 1(a) comprises the nucleotide sequence of
10 SEQ ID No. 5. An isolated nucleic acid molecule comprising polynucleotide which
hybridizes under stringent conditions to a polynucleotide having the nucleotide sequence in
(a) or (b) of claim 92. A vector comprising the nucleic acid molecule of claim 96. The
vector of claim 97, wherein said nucleic acid molecule is operably linked to a promoter for
15 the expression of a Hu-Asp polypeptide. The vector of claim 98, wherein said Hu-Asp
polypeptide is Hu-Asp1. The vector of claim 98, wherein said Hu-Asp polypeptide is Hu-
20 Asp2(a). The vector of claim 98, wherein said Hu-Asp polypeptide is Hu-Asp2(b). A host
cell comprising the vector of claim 98. A method of obtaining a Hu-Asp polypeptide
comprising culturing the host cell of claim 102 and isolating said Hu-Asp polypeptide. An
isolated Hu-Asp1 polypeptide comprising an amino acid sequence at least 95% identical to
25 a sequence comprising the amino acid sequence of SEQ ID No. 2. An isolated Hu-Asp2(a)
polypeptide comprising an amino acid sequence at least 95% identical to a sequence
comprising the amino acid sequence of SEQ ID No. 4. An isolated Hu-Asp2(a) polypeptide
comprising an amino acid sequence at least 95% identical to a sequence comprising the
amino acid sequence of SEQ ID No. 8. An isolated antibody that binds specifically to the
30 Hu-Asp polypeptide of any of claims 104-107.

20 Here we disclose numerous methods to assay the enzyme.

A method to identify a cell that can be used to screen for inhibitors of β
secretase activity comprising:

35 (a) identifying a cell that expresses a protease capable of cleaving APP at the β
secretase site, comprising:

- 25 i) collect the cells or the supernatant from the cells to be identified
- 40 ii) measure the production of a critical peptide, where the critical
peptide is selected from the group consisting of either the APP C-
terminal peptide or soluble APP,
- 45 iii) select the cells which produce the critical peptide.

30 The method of claim 108 where the cells are collected and the critical peptide is the
APP C-terminal peptide created as a result of the β secretase cleavage. The method of claim
108 where the supernatant is collected and the critical peptide is soluble APP where the
soluble APP has a C-terminal created by β secretase cleavage. The method of claim 108

5 where the cells contain any of the nucleic acids or polypeptides of claims 1-86 and where
the cells are shown to cleave the β secretase site of any peptide having the following
peptide structure, P2, P1, P1', P2', where P2 is K or N, where P1 is M or L, where P1' is
D, where P2' is A. The method of claim 111 where P2 is K and P1 is M.. The method of
10 claim 112 where P2 is N and P1 is L.

Any bacterial cell comprising any nucleic acids or peptides in claims 1-86
and 92-107. A bacterial cell of claim 114 where the bacteria is *E coli*. Any eukaryotic cell
comprising any nucleic acids or polypeptides in claims 1-86 and 92-107.

15 Any insect cell comprising any of the nucleic acids or polypeptides in claims
1-86 and 92-107. A insect cell of claim 117 where the insect is sf9, or High 5. A insect
cell of claim 100 where the insect cell is High 5. A mammalian cell comprising any of the
20 nucleic acids or polypeptides in claims 1-86 and 92-107. A mammalian cell of claim 120
where the mammalian cell is selected from the group consisting of, human, rodent,
lagomorph, and primate. A mammalian cell of claim 121 where the mammalian cell is
15 selected from the group consisting of human cell. A mammalian cell of claim 122 where
the human cell is selected from the group comprising HEK293, and IMR-32. A
mammalian cell of claim 121 where the cell is a primate cell. A primate cell of claim 124
where the primate cell is a COS-7 cell. A mammalian cell of claim 121 where cell is
30 selected from a rodent cells. A rodent cell of claim 126 selected from, CHO-K1, Neuro-
2A, 3T3 cells. A yeast cell of claim 115. An avian cell of claim 115.

Any isoform of APP where the last two carboxy terminus amino acids of that
isoform are both lysine residues. In written descrip. Define isoform is any APP
35 polypeptide, including APP variants (including mutations), and APP fragments that exists
in humans such as those described in US 5,766,846, col 7, lines 45-67, incorporated into this
25 document by reference. The isoform of APP from claim 114, comprising the isoform
known as APP695 modified so that its last two having two lysine residues as its last two
carboxy terminus amino acids. The isoform of claim 130 comprising SEQ. ID. 16. The
40 isoform variant of claim 130 comprising SEQ. ID. NO. 18, and 20. Any eukaryotic cell
line, comprising nucleic acids or polypeptides of claim 130-132. Any cell line of claim 133
45 that is a mammalian cell line (HEK293, Neuro2a, best - plus others. A method for
30 identifying inhibitors of an enzyme that cleaves the beta secretase cleavable site of APP
comprising:

5 a) culturing cells in a culture medium under conditions in which the enzyme causes processing of APP and release of amyloid beta-peptide into the medium and causes the accumulation of CTF99 fragments of APP in cell lysates,

10 b) exposing the cultured cells to a test compound; and specifically determining whether the test compound inhibits the function of the enzyme by measuring the amount of amyloid beta-peptide released into the medium and or the amount of CTF99 fragments of APP in cell lysates;

15 c) identifying test compounds diminishing the amount of soluble amyloid beta peptide present in the culture medium and diminution of CTF99 fragments of APP in cell lysates as Asp2 inhibitors.

20 The method of claim 135 wherein the cultured cells are a human, rodent or insect cell line. The method of claim 136 wherein the human or rodent cell line exhibits β secretase activity in which processing of APP occurs with release of amyloid beta-peptide into the culture medium and accumulation of CTF99 in cell lysates. A method as in claim 25 137 wherein the human or rodent cell line treated with the antisense oligomers directed against the enzyme that exhibits β secretase activity, reduces release of soluble amyloid beta-peptide into the culture medium and accumulation of CTF99 in cell lysates. A method for the identification of an agent that decreases the activity of a Hu-Asp polypeptide selected from the group consisting of Hu-Asp1, Hu-Asp2(a), and Hu-Asp2(b), the method 30 comprising:

a) determining the activity of said Hu-Asp polypeptide in the presence of a test agent and in the absence of a test agent; and

35 b) comparing the activity of said Hu-Asp polypeptide determined in the presence of said test agent to the activity of said Hu-Asp polypeptide determined in the absence of said test agent;

40 whereby a lower level of activity in the presence of said test agent than in the absence of said test agent indicates that said test agent has decreased the activity of said Hu-Asp polypeptide. The nucleic acids, peptides, proteins, vectors, cells and cell lines, and assays described herein.

45 30 The present invention provides isolated nucleic acid molecules comprising a polynucleotide that codes for a polypeptide selected from the group consisting of human aspartyl proteases. In particular, human aspartyl protease 1 (Hu-Asp1) and two alternative splice variants of human aspartyl protease 2 (Hu-Asp2), designated herein as Hu-Asp2(a) and 50

5 Hu-Asp2(b). As used herein, all references to "Hu-Asp" should be understood to refer to all of Hu-Asp1, Hu-Asp2(a), and Hu-Asp2(b). In addition, as used herein, all references to "Hu-Asp2" should be understood to refer to both Hu-Asp2(a) and Hu-Asp2(b). Hu-Asp1 is expressed most abundantly in pancreas and prostate tissues, while Hu-Asp2(a) and Hu-Asp2(b) are expressed most abundantly in pancreas and brain tissues. The invention also provides isolated Hu-Asp1, Hu-Asp2(a), and Hu-Asp2(b) polypeptides, as well as fragments thereof which exhibit aspartyl protease activity.

10 In a preferred embodiment, the nucleic acid molecules comprise a polynucleotide having a nucleotide sequence selected from the group consisting of residues 1-1554 of SEQ ID NO:1, encoding Hu-Asp1, residues 1-1503 of SEQ ID NO:3, encoding Hu-Asp2(a), and residues 1-1428 of SEQ ID NO:5, encoding Hu-Asp2(b). In another aspect, the invention provides an isolated nucleic acid molecule comprising a polynucleotide which hybridizes under stringent conditions to a polynucleotide encoding Hu-Asp1, Hu-Asp2(a), Hu-Asp2(b), or fragments thereof. European patent application EP 0 848 062 discloses a polypeptide referred to as "Asp 1," that bears substantial homology to Hu-Asp1, while international application WO 98/22597 discloses a polypeptide referred to as "Asp 2," that bears substantial homology to Hu-Asp2(a).

15 The present invention also provides vectors comprising the isolated nucleic acid molecules of the invention, host cells into which such vectors have been introduced, and recombinant methods of obtaining a Hu-Asp1, Hu-Asp2(a), or Hu-Asp2(b) polypeptide comprising culturing the above-described host cell and isolating the relevant polypeptide.

20 In another aspect, the invention provides isolated Hu-Asp1, Hu-Asp2(a), and Hu-Asp2(b) polypeptides, as well as fragments thereof. In a preferred embodiment, the Hu-Asp1, Hu-Asp2(a), and Hu-Asp2(b) polypeptides have the amino acid sequence given in SEQ ID NO:2, SEQ ID NO:4, or SEQ ID NO:6, respectively. The present invention also describes active forms of Hu-Asp2, methods for preparing such active forms, methods for preparing soluble forms, methods for measuring Hu-Asp2 activity, and substrates for Hu-Asp2 cleavage. The invention also describes antisense oligomers targeting the Hu-Asp1, Hu-Asp2(a) and Hu-Asp2(b) mRNA transcripts and the use of such antisense reagents to decrease such mRNA and consequently the production of the corresponding polypeptide. Isolated antibodies, both polyclonal and monoclonal, that binds specifically to any of the Hu-Asp1, Hu-Asp2(a), and Hu-Asp2(b) polypeptides of the invention are also provided.

The invention also provides a method for the identification of an agent that modulates the activity of any of Hu-Asp-1, Hu-Asp2(a), and Hu-Asp2(b). The invention describes methods to test such agents in cell-free assays to which Hu-Asp2 polypeptide is added, as well as methods to test such agents in human or other mammalian cells in which Hu-Asp2 is present.

BRIEF DESCRIPTION OF THE SEQUENCE LISTINGS

Sequence ID No. 1—Human Asp-1, nucleotide sequence
 Sequence ID No. 2—Human Asp-1, predicted amino acid sequence
 Sequence ID No. 3—Human Asp-2(a), nucleotide sequence
 Sequence ID No. 4—Human Asp-2(a), predicted amino acid sequence
 Sequence ID No. 5—Human Asp-2(b), nucleotide sequence
 Sequence ID No. 6—Human Asp-2(b), predicted amino acid sequence
 Sequence ID No. 7—Murine Asp-2(a), nucleotide sequence
 Sequence ID No. 8—Murine Asp-2(a), predicted amino acid sequence
 Sequence ID No. 9—Human APP695, nucleotide sequence
 Sequence ID No. 10—Human APP695, predicted amino acid sequence
 Sequence ID No. 11—Human APP695-Sw, nucleotide sequence
 Sequence ID No. 12—Human APP695-Sw, predicted amino acid sequence
 Sequence ID No. 13—Human APP695-VF, nucleotide sequence
 Sequence ID No. 14—Human APP695-VF, predicted amino acid sequence
 Sequence ID No. 15—Human APP695-KK, nucleotide sequence
 Sequence ID No. 16—Human APP695-KK, predicted amino acid sequence
 Sequence ID No. 17—Human APP695-Sw-KK, nucleotide sequence
 Sequence ID No. 18—Human APP695-Sw-KK, predicted amino acid sequence
 Sequence ID No. 19—Human APP695-VF-KK, nucleotide sequence
 Sequence ID No. 20—Human APP695-VF-KK, predicted amino acid sequence
 Sequence ID No. 21—T7-Human-pro-Asp-2(a) Δ TM, nucleotide sequence
 Sequence ID No. 22—T7-Human-pro-Asp-2(a) Δ TM, amino acid sequence
 Sequence ID No. 23—T7-Caspase-Human-pro-Asp-2(a) Δ TM, nucleotide sequence
 Sequence ID No. 24—T7-Caspase-Human-pro-Asp-2(a) Δ TM, amino acid sequence
 Sequence ID No. 25—Human-pro-Asp-2(a) Δ TM (low GC), nucleotide sequence
 Sequence ID No. 26—Human-pro-Asp-2(a) Δ TM, (low GC), amino acid sequence
 Sequence ID No. 27—T7-Caspase-Caspase 8 cleavage-Human-pro-Asp-2(a) Δ TM, nucleotide sequence
 Sequence ID No. 28—T7-Caspase-Caspase 8 cleavage-Human-pro-Asp-2(a) Δ TM, amino acid sequence
 Sequence ID No. 29—Human Asp-2(a) Δ TM, nucleotide sequence
 Sequence ID No. 30—Human Asp-2(a) Δ TM, amino acid sequence
 Sequence ID No. 31—Human Asp-2(a) Δ TM(His)₆, nucleotide sequence
 Sequence ID No. 32—Human Asp-2(a) Δ TM(His)₆, amino acid sequence
 Sequence ID No.s 33-46 are described below in the Detailed Description of the Invention.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1: Figure 1 shows the nucleotide (SEQ ID NO:1) and predicted amino acid sequence (SEQ ID NO:2) of human Asp1.

Figure 2: Figure 2 shows the nucleotide (SEQ ID NO:3) and predicted amino acid sequence (SEQ ID NO:4) of human Asp2(a).

Figure 3: Figure 3 shows the nucleotide (SEQ ID NO:5) and predicted amino acid sequence (SEQ ID NO:6) of human Asp2(b). The predicted transmembrane domain of Hu-Asp2(b) is enclosed in brackets.

Figure 4: Figure 4 shows the nucleotide (SEQ ID No. 7) and predicted amino acid sequence (SEQ ID No. 8) of murine Asp2(a)

Figure 5: Figure 5 shows the BestFit alignment of the predicted amino acid sequences of Hu-Asp2(a) and murine Asp2(a)

Figure 6: Figure 6 shows the nucleotide (SEQ ID No. 21) and predicted amino acid sequence (SEQ ID No. 22) of T7-Human-pro-Asp-2(a) Δ TM

Figure 7: Figure 7 shows the nucleotide (SEQ ID No. 23) and predicted amino acid sequence (SEQ ID No. 24) of T7-caspase-Human-pro-Asp-2(a) Δ TM

Figure 8: Figure 8 shows the nucleotide (SEQ ID No. 25) and predicted amino acid sequence (SEQ ID No. 26) of Human-pro-Asp-2(a) Δ TM (low GC)

Figure 9: Western blot showing reduction of CTF99 production by HEK125.3 cells transfected with antisense oligomers targeting the Hu-Asp2 Mrna

Figure 10: Western blot showing increase in CTF99 production in mouse Neuro-2a cells cotransfected with APP-KK with and without Hu-Asp2 only in those cells cotransfected with Hu-Asp2. A further increase in CTF99 production is seen in cells cotransfected with APP-Sw-KK with and without Hu-Asp2 only in those cells cotransfected with Hu-Asp2

Figure 11: Figure 11 shows the predicted amino acid sequence (SEQ ID No. 30) of Human-Asp2(a) Δ TM

Figure 12: Figure 11 shows the predicted amino acid sequence (SEQ ID No. 30) of Human-Asp2(a) Δ TM(His)₆

DETAILED DESCRIPTION OF THE INVENTION

A few definitions used in this invention follow, most definitions to be used are those that would be used by one ordinarily skilled in the art.

When the β amyloid peptide any peptide resulting from beta secretase cleavage of APP. This includes, peptides of 39, 40, 41, 42 and 43 amino acids, extending from the β -

5 secretase cleavage site to 39, 40, 41, 42 and 43 amino acids. β amyloid peptide also means sequences 1-6, SEQ. ID. NO. 1-6 of US 5,750,349, issued 12 May 1998 (incorporated into this document by reference). A β -secretase cleavage fragment disclosed here is called CTF-99, which extends from β -secretase cleavage site to the carboxy terminus of APP.

10 5 When an isoform of APP is discussed then what is meant is any APP polypeptide, including APP variants (including mutations), and APP fragments that exists in humans such as those described in US 5,766,846, col 7, lines 45-67, incorporated into this document by reference and see below.

15 The term " β -amyloid precursor protein" (APP) as used herein is defined as a polypeptide that is encoded by a gene of the same name localized in humans on the long arm of chromosome 21 and that includes " β AP - here " β -amyloid protein" see above, within its carboxyl third. APP is a glycosylated, single-membrane spanning protein expressed in a wide variety of cells in many mammalian tissues. Examples of specific isotypes of APP which are currently known to exist in humans are the 695-amino acid polypeptide described by Kang et. al. (1987) Nature 325:733-736 which is designated as the "normal" APP; the 751-amino acid polypeptide described by Ponte et al. (1988) Nature 331:525-527 (1988) and Tanzi et al. (1988) Nature 331:528-530; and the 770-amino acid polypeptide described by Kitaguchi et. al. (1988) Nature 331:530-532. Examples of specific variants of APP include point mutation which can differ in both position and phenotype (for review of known variant mutation see Hardy (1992) Nature Genet. 1:233-234). All references cited here incorporated by reference. The term "APP fragments" as used herein refers to fragments of APP other than those which consist solely of β AP or β AP fragments. That is, APP fragments will include amino acid sequences of APP in addition to those which form intact 3AP or a fragment of β AP.

25 20 When the term "any amino acid" is used, the amino acids referred to are to be selected from the following, three letter and single letter abbreviations - which may also be used, are provided as follows:

40 Alanine, Ala, A; Arginine, Arg, R; Asparagine, Asn, N; Aspartic acid, Asp, D; Cystein, Cys, C; Glutamine, Gln, Q; l;E-Glutamic Acid, Glu, E; Glycine, Gly, G; 45 Histidine, His, H; Isoleucine, Ile, I; Leucine, Leu, L; Lysine, Lys, K; Methionine, Met, M; Phenylalanine, Phe, F; Proline, Pro, P; Serine, Ser, S; Threonine, Thr, T; Tryptophan, Trp, W; Tyrosine, Tyr, Y; Valine, Val, V; Aspartic acid or Asparagine, Asx, B; Glutamic acid or Glutamine, Glx, Z; Any amino acid, Xaa, X..

5 The present invention describes a method to scan gene databases for the simple active site motif characteristic of aspartyl proteases. Eukaryotic aspartyl proteases such as pepsin and renin possess a two-domain structure which folds to bring two aspartyl residues into proximity within the active site. These are embedded in the short tripeptide motif
10 5 DTG, or more rarely, DSG. Most aspartyl proteases occur as proenzyme whose N-terminus must be cleaved for activation. The DTG or DSG active site motif appears at about residue 65-70 in the proenzyme (prorenin, pepsinogen), but at about residue 25-30 in the active enzyme after cleavage of the N-terminal prodomain. The limited length of the active site motif makes it difficult to search collections of short, expressed sequence tags (EST) for
15 10 novel aspartyl proteases. EST sequences typically average 250 nucleotides or less, and so would encode 80-90 amino acid residues or less. That would be too short a sequence to span the two active site motifs. The preferred method is to scan databases of hypothetical or assembled protein coding sequences. The present invention describes a computer method to identify candidate aspartyl proteases in protein sequence databases. The method
20 15 was used to identify seven candidate aspartyl protease sequences in the *Caenorhabditis elegans* genome. These sequences were then used to identify by homology search Hu-Asp1 and two alternative splice variants of Hu-Asp2, designated herein as Hu-Asp2(a) and Hu-Asp2(b).

30 In a major aspect of the invention disclosed here we provide new information about APP processing. Pathogenic processing of the amyloid precursor protein (APP) via the $A\beta$ pathway requires the sequential action of two proteases referred to as β -secretase and γ -secretase. Cleavage of APP by the β -secretase and γ -secretase generates the N-terminus and C-terminus of the $A\beta$ peptide, respectively. Because over production of the $A\beta$ peptide, particularly the $A\beta_{1-42}$, has been implicated in the initiation of Alzheimer's disease,
35 25 inhibitors of either the β -secretase and/or the γ -secretase have potential in the treatment of Alzheimer's disease. Despite the importance of the β -secretase and γ -secretase in the pathogenic processing of APP, molecular definition of these enzymes has not been accomplished to date. That is, it was not known what enzymes were required for cleavage at either the β -secretase or the γ -secretase cleavage site. The sites themselves were
40 45 known because APP was known and the $A\beta_{1-42}$ peptide was known, see US 5,766,846 and US 5,837,672, (incorporated by reference, with the exception to reference to "soluble" peptides). But what enzyme was involved in producing the $A\beta_{1-42}$ peptide was unknown.

5 The present invention involves the molecular definition of several novel human
aspartyl proteases and one of these, referred to as Hu-Asp-2(a) and Hu-Asp2(b), has been
characterized in detail. Previous forms of asp1 and asp 2 have been disclosed, see EP
0848062 A2 and EP 0855444A2, inventors David Powel et. al., assigned to Smith Kline
10 Beecham Corp. (incorporated by reference). Herein are disclosed old and new forms of
Hu-Asp 2. For the first time they are expressed in active form, their substrates are
disclosed, and their specificity is disclosed. Prior to this disclosure cell or cell extracts were
required to cleave the β -secretase site, now purified protein can be used in assays, also
15 described here. Based on the results of (1) antisense knock out experiments, (2) transient
transfection knock in experiments, and (3) biochemical experiments using purified
recombinant Hu-Asp-2, we demonstrate that Hu-Asp-2 is the β -secretase involved in the
processing of APP. Although the nucleotide and predicted amino acid sequence of Hu-
Asp-2(a) has been reported, see above, see EP 0848062 A2 and EP 0855444A2, no
20 functional characterization of the enzyme was disclosed. Here the authors characterize the
Hu-Asp-2 enzyme and are able to explain why it is a critical and essential enzyme required
25 in the formation of $A\beta_{1-42}$ peptide and possible a critical step in the development of AD.

In another embodiment the present invention also describes a novel splice variant of
Hu-Asp2, referred to as Hu-Asp2(b), that has never before been disclosed.

30 In another embodiment, the invention provides isolated nucleic acid molecules
comprising a polynucleotide encoding a polypeptide selected from the group consisting of
human aspartyl protease 1 (Hu-Asp1) and two alternative splice variants of human aspartyl
protease 2 (Hu-Asp2), designated herein as Hu-Asp2(a) and Hu-Asp2(b). As used herein, all
35 references to "Hu-Asp2" should be understood to refer to both Hu-Asp2(a) and Hu-Asp2(b).
Hu-Asp1 is expressed most abundantly in pancreas and prostate tissues, while Hu-Asp2(a)
and Hu-Asp2(b) are expressed most abundantly in pancreas and brain tissues. The invention
25 also provides isolated Hu-Asp1, Hu-Asp2(a), and Hu-Asp2(b) polypeptides, as well as
fragments thereof which exhibit aspartyl protease activity.

The predicted amino acid sequences of Hu-Asp1, Hu-Asp2(a) and Hu-Asp2(b) share
45 significant homology with previously identified mammalian aspartyl proteases such as
pepsinogen A, pepsinogen B, cathepsin D, cathepsin E, and renin. P.B.Szecs, *Scand. J. Clin.*
30 *Lab. Invest.* 52:(Suppl. 210 5-22 (1992)). These enzymes are characterized by the presence of
a duplicated DTG/DSG sequence motif. The Hu-Asp1 and HuAsp2 polypeptides disclosed

5 herein also exhibit extremely high homology with the ProSite consensus motif for aspartyl proteases extracted from the SwissProt database.

10 The nucleotide sequence given as residues 1-1554 of SEQ ID NO:1 corresponds to the nucleotide sequence encoding Hu-Asp1, the nucleotide sequence given as residues 1-1503 of SEQ ID NO:3 corresponds to the nucleotide sequence encoding Hu-Asp2(a), and the nucleotide sequence given as residues 1-1428 of SEQ ID NO:5 corresponds to the nucleotide sequence encoding Hu-Asp2(b). The isolation and sequencing of DNA encoding Hu-Asp1, Hu-Asp2(a), and Hu-Asp2(b) is described below in Examples 1 and 2.

15 As is described in Examples 1 and 2, automated sequencing methods were used to obtain the nucleotide sequence of Hu-Asp1, Hu-Asp2(a), and Hu-Asp2(b). The Hu-Asp nucleotide sequences of the present invention were obtained for both DNA strands, and are believed to be 100% accurate. However, as is known in the art, nucleotide sequence obtained by such automated methods may contain some errors. Nucleotide sequences determined by automation are typically at least about 90%, more typically at least about 95% to at least about 20 99.9% identical to the actual nucleotide sequence of a given nucleic acid molecule. The actual sequence may be more precisely determined using manual sequencing methods, which are well known in the art. An error in sequence which results in an insertion or deletion of one or more nucleotides may result in a frame shift in translation such that the predicted amino acid sequence will differ from that which would be predicted from the actual nucleotide sequence of the nucleic acid molecule, starting at the point of the mutation. The Hu-Asp DNA of the present invention includes cDNA, chemically synthesized DNA, DNA isolated by PCR, genomic DNA, and combinations thereof. Genomic Hu-Asp DNA may be obtained by screening a genomic library with the Hu-Asp2 cDNA described herein, using methods that are well known in the art, or with oligonucleotides chosen from the Hu-Asp2 25 sequence that will prime the polymerase chain reaction (PCR). RNA transcribed from Hu-Asp DNA is also encompassed by the present invention.

30 Due to the degeneracy of the genetic code, two DNA sequences may differ and yet encode identical amino acid sequences. The present invention thus provides isolated nucleic acid molecules having a polynucleotide sequence encoding any of the Hu-Asp polypeptides of the invention, wherein said polynucleotide sequence encodes a Hu-Asp polypeptide having the complete amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, or fragments thereof.

Also provided herein are purified Hu-Asp polypeptides, both recombinant and non-recombinant. Most importantly, methods to produce Hu-Asp2 polypeptides in active form are provided. These include production of Hu-Asp2 polypeptides and variants thereof in bacterial cells, insect cells, and mammalian cells, also in forms that allow secretion of the Hu-Asp2 polypeptide from bacterial, insect or mammalian cells into the culture medium, also methods to produce variants of Hu-Asp2 polypeptide incorporating amino acid tags that facilitate subsequent purification. In a preferred embodiment of the invention the Hu-Asp2 polypeptide is converted to a proteolytically active form either in transformed cells or after purification and cleavage by a second protease in a cell-free system, such active forms of the Hu-Asp2 polypeptide beginning with the N-terminal sequence TQHGIR or ETDEEP. Variants and derivatives, including fragments, of Hu-Asp proteins having the native amino acid sequences given in SEQ ID Nos: 2, 4, and 6 that retain any of the biological activities of Hu-Asp are also within the scope of the present invention. Of course, one of ordinary skill in the art will readily be able to determine whether a variant, derivative, or fragment of a Hu-Asp protein displays Hu-Asp activity by subjecting the variant, derivative, or fragment to a standard aspartyl protease assay. Fragments of Hu-Asp within the scope of this invention include those that contain the active site domain containing the amino acid sequence DTG, fragments that contain the active site domain amino acid sequence DSG, fragments containing both the DTG and DSG active site sequences, fragments in which the spacing of the DTG and DSG active site sequences has been lengthened, fragments in which the spacing has been shortened. Also within the scope of the invention are fragments of Hu-Asp in which the transmembrane domain has been removed to allow production of Hu-Asp2 in a soluble form. In another embodiment of the invention, the two halves of Hu-Asp2, each containing a single active site DTG or DSG sequence can be produced independently as recombinant polypeptides, then combined in solution where they reconstitute an active protease.

Hu-Asp variants may be obtained by mutation of native Hu-Asp-encoding nucleotide sequences, for example. A Hu-Asp variant, as referred to herein, is a polypeptide substantially homologous to a native Hu-Asp polypeptide but which has an amino acid sequence different from that of native Hu-Asp because of one or more deletions, insertions, or substitutions in the amino acid sequence. The variant amino acid or nucleotide sequence is preferably at least about 80% identical, more preferably at least about 90% identical, and most preferably at least about 95% identical, to a native Hu-Asp sequence. Thus, a variant nucleotide sequence which contains, for example, 5 point mutations for every one hundred

5 nucleotides, as compared to a native Hu-Asp gene, will be 95% identical to the native protein.
The percentage of sequence identity, also termed homology, between a native and a variant
Hu-Asp sequence may also be determined, for example, by comparing the two sequences
using any of the computer programs commonly employed for this purpose, such as the Gap
10 5 program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer
Group, University Research Park, Madison Wisconsin), which uses the algorithm of Smith
and Waterman (*Adv. Appl. Math.* 2: 482-489 (1981)).

15 Alterations of the native amino acid sequence may be accomplished by any of a
number of known techniques. For example, mutations may be introduced at particular
10 locations by procedures well known to the skilled artisan, such as oligonucleotide-directed
mutagenesis, which is described by Walder *et al.* (*Gene* 42:133 (1986)); Bauer *et al.* (*Gene*
20 37:73 (1985)); Craik (*BioTechniques*, January 1985, pp. 12-19); Smith *et al.* (*Genetic
Engineering: Principles and Methods*, Plenum Press (1981)); and U.S. Patent Nos. 4,518,584
and 4,737,462.

15 Hu-Asp variants within the scope of the invention may comprise conservatively
substituted sequences, meaning that one or more amino acid residues of a Hu-Asp polypeptide
are replaced by different residues that do not alter the secondary and/or tertiary structure of the
Hu-Asp polypeptide. Such substitutions may include the replacement of an amino acid by a
30 residue having similar physicochemical properties, such as substituting one aliphatic residue
(Ile, Val, Leu or Ala) for another, or substitution between basic residues Lys and Arg, acidic
20 residues Glu and Asp, amide residues Gln and Asn, hydroxyl residues Ser and Tyr, or
aromatic residues Phe and Tyr. Further information regarding making phenotypically silent
amino acid exchanges may be found in Bowie *et al.*, *Science* 247:1306-1310 (1990). Other
35 Hu-Asp variants which might retain substantially the biological activities of Hu-Asp are those
25 where amino acid substitutions have been made in areas outside functional regions of the
protein.
40

In another aspect, the invention provides an isolated nucleic acid molecule comprising
a polynucleotide which hybridizes under stringent conditions to a portion of the nucleic acid
45 molecules described above, *e.g.*, to at least about 15 nucleotides, preferably to at least about
30 20 nucleotides, more preferably to at least about 30 nucleotides, and still more preferably to at
least about from 30 to at least about 100 nucleotides, of one of the previously described
nucleic acid molecules. Such portions of nucleic acid molecules having the described lengths
50 refer to, *e.g.*, at least about 15 contiguous nucleotides of the reference nucleic acid molecule.

5 By stringent hybridization conditions is intended overnight incubation at about 42°C for about 2.5 hours in 6 X SSC/0.1% SDS, followed by washing of the filters in 1.0 X SSC at 65°C, 0.1% SDS.

10 5 Fragments of the Hu-Asp-encoding nucleic acid molecules described herein, as well as polynucleotides capable of hybridizing to such nucleic acid molecules may be used as a probe or as primers in a polymerase chain reaction (PCR). Such probes may be used, e.g., to detect the presence of Hu-Asp nucleic acids in *in vitro* assays, as well as in Southern and northern blots. Cell types expressing Hu-Asp may also be identified by the use of such probes. Such
15 procedures are well known, and the skilled artisan will be able to choose a probe of a length suitable to the particular application. For PCR, 5' and 3' primers corresponding to the termini of a desired Hu-Asp nucleic acid molecule are employed to isolate and amplify that sequence using conventional techniques.

20 Other useful fragments of the Hu-Asp nucleic acid molecules are antisense or sense oligonucleotides comprising a single-stranded nucleic acid sequence capable of binding to a target Hu-Asp mRNA (using a sense strand), or Hu-Asp DNA (using an antisense strand)
25 sequence. In a preferred embodiment of the invention these Hu-Asp antisense oligonucleotides reduce Hu-Asp mRNA and consequent production of Hu-Asp polypeptides.

30 In another aspect, the invention includes Hu-Asp polypeptides with or without associated native pattern glycosylation. Both Hu-Asp1 and Hu-Asp2 have canonical acceptor sites for Asn-linked sugars, with Hu-Asp1 having two of such sites, and Hu-Asp2 having four. Hu-Asp expressed in yeast or mammalian expression systems (discussed below) may be
35 similar to or significantly different from a native Hu-Asp polypeptide in molecular weight and glycosylation pattern. Expression of Hu-Asp in bacterial expression systems will provide non-glycosylated Hu-Asp.

40 25 The polypeptides of the present invention are preferably provided in an isolated form, and preferably are substantially purified. Hu-Asp polypeptides may be recovered and purified from tissues, cultured cells, or recombinant cell cultures by well-known methods, including ammonium sulfate or ethanol precipitation, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity
45 chromatography, hydroxylapatite chromatography, lectin chromatography, and high performance liquid chromatography (HPLC). In a preferred embodiment, an amino acid tag is added to the Hu-Asp polypeptide using genetic engineering techniques that are well known to
50 practitioners of the art which include addition of six histidine amino acid residues to allow

5 purification by binding to nickel immobilized on a suitable support, epitopes for polyclonal or monoclonal antibodies including but not limited to the T7 epitope, the myc epitope, and the V5a epitope, and fusion of Hu-Asp2 to suitable protein partners including but not limited to glutathione-S-transferase or maltose binding protein. In a preferred embodiment these
10 5 additional amino acid sequences are added to the C-terminus of Hu-Asp but may be added to the N-terminus or at intervening positions within the Hu-Asp2 polypeptide.

The present invention also relates to vectors comprising the polynucleotide molecules of the invention, as well as host cell transformed with such vectors. Any of the polynucleotide
15 molecules of the invention may be joined to a vector, which generally includes a selectable marker and an origin of replication, for propagation in a host. Because the invention also
10 provides Hu-Asp polypeptides expressed from the polynucleotide molecules described above, vectors for the expression of Hu-Asp are preferred. The vectors include DNA encoding any of the Hu-Asp polypeptides described above or below, operably linked to suitable transcriptional or translational regulatory sequences, such as those derived from a mammalian, microbial,
20 viral, or insect gene. Examples of regulatory sequences include transcriptional promoters, operators, or enhancers, mRNA ribosomal binding sites, and appropriate sequences which control transcription and translation. Nucleotide sequences are operably linked when the regulatory sequence functionally relates to the DNA encoding Hu-Asp. Thus, a promoter
25 nucleotide sequence is operably linked to a Hu-Asp DNA sequence if the promoter nucleotide sequence directs the transcription of the Hu-Asp sequence.

Selection of suitable vectors to be used for the cloning of polynucleotide molecules encoding Hu-Asp, or for the expression of Hu-Asp polypeptides, will of course depend upon
35 the host cell in which the vector will be transformed, and, where applicable, the host cell from which the Hu-Asp polypeptide is to be expressed. Suitable host cells for expression of Hu-Asp polypeptides include prokaryotes, yeast, and higher eukaryotic cells, each of which is
25 discussed below.

The Hu-Asp polypeptides to be expressed in such host cells may also be fusion proteins which include regions from heterologous proteins. Such regions may be included to allow, *e.g.*, secretion, improved stability, or facilitated purification of the polypeptide. For
45 example, a sequence encoding an appropriate signal peptide can be incorporated into expression vectors. A DNA sequence for a signal peptide (secretory leader) may be fused in frame to the Hu-Asp sequence so that Hu-Asp is translated as a fusion protein comprising
30 the signal peptide. A signal peptide that is functional in the intended host cell promotes
50

5 extracellular secretion of the Hu-Asp polypeptide. Preferably, the signal sequence will be
cleaved from the Hu-Asp polypeptide upon secretion of Hu-Asp from the cell. Non-limiting
examples of signal sequences that can be used in practicing the invention include the yeast
1-factor and the honeybee melatin leader in sf9 insect cells.

10 5 In a preferred embodiment, the Hu-Asp polypeptide will be a fusion protein which
includes a heterologous region used to facilitate purification of the polypeptide. Many of the
available peptides used for such a function allow selective binding of the fusion protein to a
binding partner. For example, the Hu-Asp polypeptide may be modified to comprise a
15 peptide to form a fusion protein which specifically binds to a binding partner, or peptide tag.
Non-limiting examples of such peptide tags include the 6-His tag, thioredoxin tag,
hemagglutinin tag, GST tag, and OmpA signal sequence tag. As will be understood by one of
20 skill in the art, the binding partner which recognizes and binds to the peptide may be any
molecule or compound including metal ions (e.g., metal affinity columns), antibodies, or
fragments thereof, and any protein or peptide which binds the peptide, such as the FLAG tag.

15 25 Suitable host cells for expression of Hu-Asp polypeptides includes prokaryotes, yeast,
and higher eukaryotic cells. Suitable prokaryotic hosts to be used for the expression of Hu-
Asp include bacteria of the genera *Escherichia*, *Bacillus*, and *Salmonella*, as well as members
of the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*. For expression in, e.g., *E.*
30 *coli*, a Hu-Asp polypeptide may include an N-terminal methionine residue to facilitate
expression of the recombinant polypeptide in a prokaryotic host. The N-terminal Met may
optionally then be cleaved from the expressed Hu-Asp polypeptide. Other N-terminal amino
acid residues can be added to the Hu-Asp polypeptide to facilitate expression in *Escherichia*
35 *coli* including but not limited to the T7 leader sequence, the T7-caspase 8 leader sequence, as
well as others leaders including tags for purification such as the 6-His tag (Example 9). Hu-
Asp polypeptides expressed in *E. coli* may be shortened by removal of the cytoplasmic tail,
40 the transmembrane domain, or the membrane proximal region. Hu-Asp polypeptides
expressed in *E. coli* may be obtained in either a soluble form or as an insoluble form which
may or may not be present as an inclusion body. The insoluble polypeptide may be rendered
soluble by guanidine HCl, urea or other protein denaturants, then refolded into a soluble form
45 before or after purification by dilution or dialysis into a suitable aqueous buffer. If the
inactive proform of the Hu-Asp was produced using recombinant methods, it may be rendered
active by cleaving off the prosegment with a second suitable protease such as human
immunodeficiency virus protease.
50

5 Expression vectors for use in prokaryotic hosts generally comprises one or more phenotypic selectable marker genes. Such genes generally encode, *e.g.*, a protein that confers antibiotic resistance or that supplies an auxotrophic requirement. A wide variety of such vectors are readily available from commercial sources. Examples include pSPORT vectors, 10 5 pGEM vectors (Promega), pPROEX vectors (LTI, Bethesda, MD), Bluescript vectors (Stratagene), pET vectors (Novagen) and pQE vectors (Qiagen).

Hu-Asp may also be expressed in yeast host cells from genera including *Saccharomyces*, *Pichia*, and *Kluveromyces*. Preferred yeast hosts are *S. cerevisiae* and *P. pastoris*. 15 10 Yeast vectors will often contain an origin of replication sequence from a 2T yeast plasmid, an autonomously replicating sequence (ARS), a promoter region, sequences for polyadenylation, sequences for transcription termination, and a selectable marker gene. Vectors replicable in both yeast and *E. coli* (termed shuttle vectors) may also be used. In 20 addition to the above-mentioned features of yeast vectors, a shuttle vector will also include sequences for replication and selection in *E. coli*. Direct secretion of Hu-Asp polypeptides 15 expressed in yeast hosts may be accomplished by the inclusion of nucleotide sequence encoding the yeast I-factor leader sequence at the 5' end of the Hu-Asp-encoding nucleotide sequence. 25

Insect host cell culture systems may also be used for the expression of Hu-Asp polypeptides. In a preferred embodiment, the Hu-Asp polypeptides of the invention are 30 20 expressed using an insect cell expression system (*see* Example 10). Additionally, a baculovirus expression system can be used for expression in insect cells as reviewed by Luckow and Summers, *BioTechnology* 6:47 (1988).

In another preferred embodiment, the Hu-Asp polypeptide is expressed in mammalian host cells. Non-limiting examples of suitable mammalian cell lines include the COS-7 line of 35 25 monkey kidney cells (Gluzman *et al.*, *Cell* 23:175 (1981)), human embryonic kidney cell line 293, and Chinese hamster ovary (CHO) cells. Preferably, Chinese hamster ovary (CHO) cells are used for expression of Hu-Asp proteins (Example 11). 40

The choice of a suitable expression vector for expression of the Hu-Asp polypeptides of the invention will of course depend upon the specific mammalian host cell to be used, and 45 30 is within the skill of the ordinary artisan. Examples of suitable expression vectors include pcDNA3 (Invitrogen) and pSVL (Pharmacia Biotech). A preferred vector for expression of Hu-Asp polypeptides is pcDNA3.1-Hygro (Invitrogen). Expression vectors for use in 50 mammalian host cells may include transcriptional and translational control sequences derived

5 from viral genomes. Commonly used promoter sequences and enhancer sequences which
may be used in the present invention include, but are not limited to, those derived from human
cytomegalovirus (CMV), Adenovirus 2, Polyoma virus, and Simian virus 40 (SV40).
Methods for the construction of mammalian expression vectors are disclosed, for example, in
10 5 Okayama and Berg (*Mol. Cell. Biol.* 3:280 (1983)); Cosman *et al.* (*Mol. Immunol.* 23:935
(1986)); Cosman *et al.* (*Nature* 312:768 (1984)); EP-A-0367566; and WO 91/18982.

The polypeptides of the present invention may also be used to raise polyclonal and
monoclonal antibodies, which are useful in diagnostic assays for detecting Hu-Asp
15 polypeptide expression. Such antibodies may be prepared by conventional techniques. See,
for example, *Antibodies: A Laboratory Manual*, Harlow and Land (eds.), Cold Spring Harbor
Laboratory Press, Cold Spring Harbor, N.Y., (1988); *Monoclonal Antibodies, Hybridomas: A
New Dimension in Biological Analyses*, Kennet *et al.* (eds.), Plenum Press, New York (1980).
20 Synthetic peptides comprising portions of Hu-Asp containing 5 to 20 amino acids may also be
used for the production of polyclonal or monoclonal antibodies after linkage to a suitable
carrier protein including but not limited to keyhole limpet hemacyanin (KLH), chicken
ovalbumin, or bovine serum albumin using various cross-linking reagents including
carbodiimides, glutaraldehyde, or if the peptide contains a cysteine, N-methylmaleimide. A
25 preferred peptide for immunization when conjugated to KLH contains the C-terminus of
Hu_Asp1 or Hu-Asp2 comprising QRRPRDPEVVNDESSLVRHRWK or
30 LRQQHDDFADDISLLK. respectively.
20

The Hu-Asp nucleic acid molecules of the present invention are also valuable for
chromosome identification, as they can hybridize with a specific location on a human
35 chromosome. Hu-Asp1 has been localized to chromosome 21, while Hu-Asp2 has been
localized to chromosome 11q23.3-24.1. There is a current need for identifying particular sites
25 on the chromosome, as few chromosome marking reagents based on actual sequence data
(repeat polymorphisms) are presently available for marking chromosomal location. Once a
40 sequence has been mapped to a precise chromosomal location, the physical position of the
sequence on the chromosome can be correlated with genetic map data. The relationship
between genes and diseases that have been mapped to the same chromosomal region can then
45 be identified through linkage analysis, wherein the coinheritance of physically adjacent genes
is determined. Whether a gene appearing to be related to a particular disease is in fact the
cause of the disease can then be determined by comparing the nucleic acid sequence between
50 affected and unaffected individuals.

5 In another embodiment, the invention relates to a method of assaying Hu-Asp function, specifically Hu-Asp2 function which involves incubating in solution the Hu-Asp polypeptide with a suitable substrate including but not limited to a synthetic peptide containing the β -secretase cleavage site of APP, preferably one containing the mutation found in a Swedish
10 kindred with inherited AD in which KM is changed to NL, such peptide comprising the sequence SEVNLDAEFR in an acidic buffering solution, preferably an acidic buffering solution of pH5.5 (see Example 12) using cleavage of the peptide monitored by high
15 performance liquid chromatography as a measure of Hu-Asp proteolytic activity. Preferred assays for proteolytic activity utilize internally quenched peptide assay substrates. Such
20 suitable substrates include peptides which have attached a paired fluorophore and quencher including but not limited to coumarin and dinitrophenol, respectively, such that cleavage of the peptide by the Hu-Asp results in increased fluorescence due to physical separation of the fluorophore and quencher. Preferred colorimetric assays of Hu-Asp proteolytic activity utilize
25 other suitable substrates that include the P2 and P1 amino acids comprising the recognition site for cleavage linked to o-nitrophenol through an amide linkage, such that cleavage by the Hu-Asp results in an increase in optical density after altering the assay buffer to alkaline pH.

In another embodiment, the invention relates to a method for the identification of an agent that increases the activity of a Hu-Asp polypeptide selected from the group consisting of
30 Hu-Asp1, Hu-Asp2(a), and Hu-Asp2(b), the method comprising

- (a) determining the activity of said Hu-Asp polypeptide in the presence of a test agent and in the absence of a test agent; and
- (b) comparing the activity of said Hu-Asp polypeptide determined in the
35 presence of said test agent to the activity of said Hu-Asp polypeptide determined in the absence of said test agent;

25 whereby a higher level of activity in the presence of said test agent than in the absence of said test agent indicates that said test agent has increased the activity of said Hu-Asp polypeptide. Such tests can be performed with Hu-Asp polypeptide in a cell free system and with cultured cells that express Hu-Asp as well as variants or isoforms thereof.
40

45 In another embodiment, the invention relates to a method for the identification of an agent that decreases the activity of a Hu-Asp polypeptide selected from the group consisting of Hu-Asp1, Hu-Asp2(a), and Hu-Asp2(b), the method comprising
50

5 (a) determining the activity of said Hu-Asp polypeptide in the presence of a test agent and in the absence of a test agent; and

(b) comparing the activity of said Hu-Asp polypeptide determined in the presence of said test agent to the activity of said Hu-Asp polypeptide
10 5 determined in the absence of said test agent;

whereby a lower level of activity in the presence of said test agent than in the absence of said test agent indicates that said test agent has decreased the activity of said Hu-Asp polypeptide. Such tests can be performed with Hu-Asp polypeptide in a cell free system and with cultured
15 cells that express Hu-Asp as well as variants or isoforms thereof.

10 In another embodiment, the invention relates to a novel cell line (HEK125.3 cells) for measuring processing of amyloid β peptide ($A\beta$) from the amyloid protein precursor (APP). The cells are stable transformants of human embryonic kidney 293 cells (HEK293) with a bicistronic vector derived from pIRES-EGFP (Clontech) containing a modified
20 human APP cDNA, an internal ribosome entry site and an enhanced green fluorescent protein (EGFP) cDNA in the second cistron. The APP cDNA was modified by adding two
15 lysine codons to the carboxyl terminus of the APP coding sequence. This increases processing of $A\beta$ peptide from human APP by 2-4 fold. This level of $A\beta$ peptide processing is 60 fold higher than is seen in nontransformed HEK293 cells. HEK125.3 cells
30 will be useful for assays of compounds that inhibit $A\beta$ peptide processing. This invention also includes addition of two lysine residues to the C-terminus of other APP isoforms
20 including the 751 and 770 amino acid isoforms, to isoforms of APP having mutations found in human AD including the Swedish KM \rightarrow NL and V717 \rightarrow F mutations, to C-terminal
35 fragments of APP, such as those beginning with the β -secretase cleavage site, to C-terminal fragments of APP containing the β -secretase cleavage site which have been operably linked
25 to an N-terminal signal peptide for membrane insertion and secretion, and to C-terminal fragments of APP which have been operably linked to an N-terminal signal peptide for
40 membrane insertion and secretion and a reporter sequence including but not limited to green fluorescent protein or alkaline phosphatase, such that β -secretase cleavage releases
45 the reporter protein from the surface of cells expressing the polypeptide.


30 Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

EXAMPLES

Example 1: Development of a Search Algorithm Useful for the Identification of Aspartyl Proteases, and Identification of C. elegans Aspartyl Protease Genes in Wormpep 12:


Materials and Methods:

Classical aspartyl proteases such as pepsin and renin possess a two-domain structure which folds to bring two aspartyl residues into proximity within the active site. These are embedded in the short tripeptide motif DTG, or more rarely, DSG. The DTG or DSG active site motif appears at about residue 25-30 in the enzyme, but at about 65-70 in the proenzyme (prorenin, pepsinogen). This motif appears again about 150-200 residues downstream. The proenzyme is activated by cleavage of the N-terminal prodomain. This pattern exemplifies the double domain structure of the modern day aspartyl enzymes which apparently arose by gene duplication and divergence. Thus;



where X denotes the beginning of the enzyme, following the N-terminal prodomain, and Y denotes the center of the molecule where the gene repeat begins again.

In the case of the retroviral enzymes such as the HIV protease, they represent only a half of the two-domain structures of well-known enzymes like pepsin, cathepsin D, renin, etc. They have no prosegment, but are carved out of a polyprotein precursor containing the gag and pol proteins of the virus. They can be represented by:



This "monomer" only has about 100 aa, so is extremely parsimonious as compared to the other aspartyl protease "dimers" which have of the order of 330 or so aa, not counting the N-terminal prodomain.

The limited length of the eukaryotic aspartyl protease active site motif makes it difficult to search EST collections for novel sequences. EST sequences typically average 250 nucleotides, and so in this case would be unlikely to span both aspartyl protease active site motifs. Instead, we turned to the *C. elegans* genome. The *C. elegans* genome is estimated to contain around 13,000 genes. Of these, roughly 12,000 have been sequenced and the corresponding hypothetical open reading frame (ORF) has been placed in the database Wormpep12. We used this database as the basis for a whole genome scan of a higher eukaryote for novel aspartyl proteases, using an algorithm that we developed

specifically for this purpose. The following AWK script for locating proteins containing two DTG or DSG motifs was used for the search, which was repeated four times to recover all pairwise combinations of the aspartyl motif.

```

BEGIN{RS=">"}          /* defines ">" as record separator for FASTA format */
{
  pos = index($0,"DTG")  /* finds "DTG" in record */
  if (pos>0) {
    rest = substr($0,pos+3) /* get rest of record after first DTG */
    pos2 = index(rest,"DTG") /* find second DTG */
    if (pos2>0) printf ("%s%s\n", ">", $0) /* report hits */
  }
}

```

The AWK script shown above was used to search Wormpep12, which was downloaded from ftp.sanger.ac.uk/pub/databases/wormpep, for sequence entries containing at least two DTG or DSG motifs. Using AWK limited each record to 3000 characters or less. Thus, 35 or so larger records were eliminated manually from Wormpep12 as in any case these were unlikely to encode aspartyl proteases.

Results and Discussion:

The Wormpep 12 database contains 12,178 entries, although some of these (<10%) represent alternatively spliced transcripts from the same gene. Estimates of the number of genes encoded in the *C. elegans* genome is on the order of 13,000 genes, so Wormpep12 may be estimated to cover greater than 90% of the *C. elegans* genome.

Eukaryotic aspartyl proteases contain a two-domain structure, probably arising from ancestral gene duplication. Each domain contains the active site motif D(S/T)G located from 20-25 amino acid residues into each domain. The retroviral (e.g., HIV protease) or retrotransposon proteases are homodimers of subunits which are homologous to a single eukaryotic aspartyl protease domain. An AWK script was used to search the Wormpep12 database for proteins in which the D(S/T)G motif occurred at least twice. This identified >60 proteins with two DTG or DSG motifs. Visual inspection was used to select proteins in which the position of the aspartyl domains was suggestive of a two-domain structure meeting the criteria described above.

In addition, the PROSITE eukaryotic and viral aspartyl protease active site pattern PS00141 was used to search Wormpep12 for candidate aspartyl proteases. (Bairoch A., Bucher P., Hofmann K., The PROSITE database: its status in 1997, *Nucleic Acids Res.* 24:217-221(1997)). This generated an overlapping set of Wormpep12 sequences. Of these,

5 seven sequences contained two DTG or DSG motifs and the PROSITE aspartyl protease active site pattern. Of these seven, three were found in the same cosmid clone (F21F8.3, F21F8.4, and F21F8.7) suggesting that they represent a family of proteins that arose by ancestral gene duplication. Two other ORFs with extensive homology to F21F8.3, F21F8.4 and F21F8.7 are present in the same gene cluster (F21F8.2 and F21F8.6), however, these contain only a single DTG motif. Exhaustive BLAST searches with these seven sequences against Wormpep12 failed to reveal additional candidate aspartyl proteases in the *C. elegans* genome containing two repeats of the DTG or DSG motif.

15 BLASTX search with each *C. elegans* sequence against SWISS-PROT, GenPep and TREMBL revealed that R12H7.2 was the closest worm homologue to the known mammalian aspartyl proteases, and that T18H9.2 was somewhat more distantly related, while CEASP1, F21F8.3, F21F8.4, and F21F8.7 formed a subcluster which had the least sequence homology to the mammalian sequences.

Discussion:

15 APP, the presenilins, and p35, the activator of cdk5, all undergo intracellular proteolytic processing at sites which conform to the substrate specificity of the HIV protease. Dysregulation of a cellular aspartyl protease with the same substrate specificity, might therefore provide a unifying mechanism for causation of the plaque and tangle pathologies in AD. Therefore, we sought to identify novel human aspartyl proteases. A whole genome scan in *C. elegans* identified seven open reading frames that adhere to the aspartyl protease profile that we had identified. These seven aspartyl proteases probably comprise the complete complement of such proteases in a simple, multicellular eukaryote. These include four closely related aspartyl proteases unique to *C. elegans* which probably arose by duplication of an ancestral gene. The other three candidate aspartyl proteases (T18H9.2, R12H7.2 and C11D2.2) were found to have homology to mammalian gene sequences.

Example 2: Identification of Novel Human Aspartyl Proteases Using Database Mining by Genome Bridging

Materials and Methods:

- 5 Computer-assisted analysis of EST databases, cDNA, and predicted polypeptide sequences:

10 Exhaustive homology searches of EST databases with the CEASP1, F21F8.3, F21F8.4, and F21F8.7 sequences failed to reveal any novel mammalian homologues. TBLASTN searches with R12H7.2 showed homology to cathepsin D, cathepsin E, 15 pepsinogen A, pepsinogen C and renin, particularly around the DTG motif within the active site, but also failed to identify any additional novel mammalian aspartyl proteases. This indicates that the *C. elegans* genome probably contains only a single lysosomal aspartyl protease which in mammals is represented by a gene family that arose through duplication and consequent modification of an ancestral gene.

- 15 TBLASTN searches with T18H9.2, the remaining *C. elegans* sequence, identified several ESTs which assembled into a contig encoding a novel human aspartyl protease (Hu-ASP1). As is described above in Example 1, BLASTX search with the Hu-ASP1 contig 25 against SWISS-PROT revealed that the active site motifs in the sequence aligned with the active sites of other aspartyl proteases. Exhaustive, repetitive rounds of BLASTN searches against LifeSeq, LifeSeqFL, and the public EST collections identified 102 EST from 30 multiple cDNA libraries that assembled into a single contig. The 51 sequences in this contig found in public EST collections also have been assembled into a single contig (THC213329) by The Institute for Genome Research (TIGR). The TIGR annotation indicates that they failed to find any hits in the database for the contig. Note that the TIGR 35 contig is the reverse complement of the LifeSeq contig that we assembled. BLASTN search of Hu-ASP1 against the rat and mouse EST sequences in ZooSeq revealed one homologous EST in each database (Incyte clone 700311523 and IMAGE clone 313341, GenBank 40 accession number W10530, respectively).

- TBLASTN searches with the assembled DNA sequence for Hu-ASP1 against both 30 LifeSeqFL and the public EST databases identified a second, related human sequence (Hu-Asp2) represented by a single EST (2696295). Translation of this partial cDNA sequence reveals a single DTG motif which has homology to the active site motif of a bovine aspartyl 45 protease, NM1.

BLAST searches, contig assemblies and multiple sequence alignments were performed using the bioinformatics tools provided with the LifeSeq, LifeSeqFL and LifeSeq Assembled databases from Incyte. Predicted protein motifs were identified using either the ProSite dictionary (Motifs in GCG 9) or the Pfam database.

Full-length cDNA cloning of Hu-Asp1

The open reading frame of *C. elegans* gene T18H9.2CE was used to query Incyte LifeSeq and LifeSeq-FL databases and a single electronic assembly referred to as 1863920CE1 was detected. The 5' most cDNA clone in this contig, 1863920, was obtained from Incyte and completely sequenced on both strands. Translation of the open reading frame contained within clone 1863920 revealed the presence of the duplicated aspartyl protease active site motif (DTG/DSG) but the 5' end was incomplete. The remainder of the Hu-Asp1 coding sequence was determined by 5' Marathon RACE analysis using a human placenta Marathon ready cDNA template (Clontech). A 3'-antisense oligonucleotide primer specific for the 5' end of clone 1863920 was paired with the 5'-sense primer specific for the Marathon ready cDNA synthetic adaptor in the PCR. Specific PCR products were directly sequenced by cycle sequencing and the resulting sequence assembled with the sequence of clone 1863920 to yield the complete coding sequence of Hu-Asp-1 (SEQ ID No. 1).

Several interesting features are present in the primary amino acid sequence of Hu-Asp1 (Figure 1, SEQ ID No. 2). The sequence contains a signal peptide (residues 1-20 in SEQ ID No. 2), a pro-segment, and a catalytic domain containing two copies of the aspartyl protease active site motif (DTG/DSG). The spacing between the first and second active site motifs is about 200 residues which should correspond to the expected size of a single, eukaryotic aspartyl protease domain. More interestingly, the sequence contains a predicted transmembrane domain (residues 469-492 in SEQ ID No.2) near its C-terminus which suggests that the protease is anchored in the membrane. This feature is not found in any other aspartyl protease.

Cloning of a full-length Hu-Asp-2 cDNAs:

As is described above in Example 1, genome wide scan of the *Caenorhabditis elegans* database WormPep12 for putative aspartyl proteases and subsequent mining of human EST databases revealed a human ortholog to the *C. elegans* gene T18H9.2 referred to as Hu-Asp1. The assembled contig for Hu-Asp1 was used to query for human paralogs using the BLAST search tool in human EST databases and a single significant match

(2696295CE1) with approximately 60% shared identity was found in the LifeSeq FL database. Similar queries of either gb105PubEST or the family of human databases available from TIGR did not identify similar EST clones. cDNA clone 2696295, identified by single pass sequence analysis from a human uterus cDNA library, was obtained from Incyte and completely sequence on both strands. This clone contained an incomplete 1266 bp open-reading frame that encoded a 422 amino acid polypeptide but lacked an initiator ATG on the 5' end. Inspection of the predicted sequence revealed the presence of the duplicated aspartyl protease active site motif DTG/DSG, separated by 194 amino acid residues. Subsequent queries of later releases of the LifeSeq EST database identified an additional ESTs, sequenced from a human astrocyte cDNA library (4386993), that appeared to contain additional 5' sequence relative to clone 2696295. Clone 4386993 was obtained from Incyte and completely sequenced on both strands. Comparative analysis of clone 4386993 and clone 2696295 confirmed that clone 4386993 extended the open-reading frame by 31 amino acid residues including two in-frame translation initiation codons. Despite the presence of the two in-frame ATGs, no in-frame stop codon was observed upstream of the ATG indicating that the 4386993 may not be full-length. Furthermore, alignment of the sequences of clones 2696295 and 4386993 revealed a 75 base pair insertion in clone 2696295 relative to clone 4386993 that results in the insertion of 25 additional amino acid residues in 2696295. The remainder of the Hu-Asp2 coding sequence was determined by 5' Marathon RACE analysis using a human hippocampus Marathon ready cDNA template (Clontech). A 3'-antisense oligonucleotide primer specific for the shared 5'-region of clones 2696295 and 4386993 was paired with the 5'-sense primer specific for the Marathon ready cDNA synthetic adaptor in the PCR. Specific PCR products were directly sequenced by cycle sequencing and the resulting sequence assembled with the sequence of clones 2696295 and 4386993 to yield the complete coding sequence of Hu-Asp2(a) (SEQ ID No. 3) and Hu-Asp2(b) (SEQ ID No. 5), respectively.

Several interesting features are present in the primary amino acid sequence of Hu-Asp2(a) (Figure 2 and SEQ ID No. 4) and Hu-Asp-2(b) (Figure 3, SEQ ID No. 6). Both sequences contain a signal peptide (residues 1-21 in SEQ ID No. 4 and SEQ ID No. 6), a pro-segment, and a catalytic domain containing two copies of the aspartyl protease active site motif (DTG/DSG). The spacing between the first and second active site motifs is variable due to the 25 amino acid residue deletion in Hu-Asp-2(b) and consists of 168-*versus*-194 amino acid residues, for Hu-Asp2(b) and Hu-Asp-2(a), respectively. More

5 interestingly, both sequences contains a predicted transmembrane domain (residues 455-477 in SEQ ID No.4 and 430-452 in SEQ ID No. 6) near their C-termini which indicates that the protease is anchored in the membrane. This feature is not found in any other aspartyl protease except Hu-Asp1.

10 **Example 3. Molecular cloning of mouse Asp2 cDNA and genomic DNA.**
Cloning and characterization of murine Asp2 cDNA—The murine ortholog of Hu_Asp2

was cloned using a combination of cDNA library screening, PCR, and genomic cloning.

15 Approximately 500,000 independent clones from a mouse brain cDNA library were screened using a ³²P-labeled coding sequence probe prepared from Hu_Asp2. Replicate
20 positives were subjected to DNA sequence analysis and the longest cDNA contained the entire 3' untranslated region and 47 amino acids in the coding region. PCR amplification of the same mouse brain cDNA library with an antisense oligonucleotide primer specific for the 5'-most cDNA sequence determined above and a sense primer specific for the 5' region
25 of human Asp2 sequence followed by DNA sequence analysis gave an additional 980 bp of the coding sequence. The remainder of the 5' sequence of murine Asp-2 was derived from
30 genomic sequence (see below).

Isolation and sequence analysis of the murine Asp-2 gene—A murine EST sequence encoding a portion of the murine Asp2 cDNA was identified in the GenBank EST database
35 using the BLAST search tool and the Hu-Asp2 coding sequence as the query. Clone g3160898 displayed 88% shared identity to the human sequence over 352 bp.

40 Oligonucleotide primer pairs specific for this region of murine Asp2 were then synthesized and used to amplify regions of the murine gene. Murine genomic DNA, derived from strain 129/SvJ, was amplified in the PCR (25 cycles) using various primer sets specific for murine Asp2 and the products analyzed by agarose gel electrophoresis. The primer set Zoo-1 and
45 Zoo-4 amplified a 750 bp fragment that contained approximately 600 bp of intron sequence
25 based on comparison to the known cDNA sequence. This primer set was then used to

5 screen a murine BAC library by PCR, a single genomic clone was isolated and this cloned
was confirmed contain the murine Asp2 gene by DNA sequence analysis. Shotgun DNA
sequencing of this Asp2 genomic clone and comparison to the cDNA sequences of both
10 Hu_Asp2 and the partial murine cDNA sequences defined the full-length sequence of
murine Asp2 (SEQ ID No. 7). The predicted amino acid sequence of murine Asp2 (SEQ ID
No. 8) showed 96.4% shared identity (GCG BestFit algorithm) with 18/501 amino acid
15 residue substitutions compared to the human sequence (Figure 4).

Example 4: Tissue Distribution of Expression of Hu-Asp2 Transcripts:

Materials and Methods:

20 The tissue distribution of expression of Hu-Asp-2 was determined using multiple
tissue Northern blots obtained from Clontech (Palo Alto, CA). Incyte clone 2696295 in
the vector pINCY was digested to completion with *EcoRI/NotI* and the 1.8 kb cDNA insert
purified by preparative agarose gel electrophoresis. This fragment was radiolabeled to a
25 specific activity $> 1 \times 10^9$ dpm/ μ g by random priming in the presence of [α - 32 P-dATP]
(>3000 Ci/mmol, Amersham, Arlington Heights, IL) and Klenow fragment of DNA
polymerase I. Nylon filters containing denatured, size fractionated poly A⁺ RNAs isolated
30 from different human tissues were hybridized with 2×10^6 dpm/ml probe in ExpressHyb
buffer (Clontech, Palo Alto, CA) for 1 hour at 68 °C and washed as recommended by the
manufacture. Hybridization signals were visualized by autoradiography using BioMax XR
20 film (Kodak, Rochester, NY) with intensifying screens at -80 °C.

Results and Discussion:

Limited information on the tissue distribution of expression of Hu-Asp-2 transcripts
was obtained from database analysis due to the relatively small number of ESTs detected
40 using the methods described above (< 5). In an effort to gain further information on the
expression of the Hu-Asp2 gene, Northern analysis was employed to determine both the
25 size(s) and abundance of Hu-Asp2 transcripts. PolyA⁺ RNAs isolated from a series of
peripheral tissues and brain regions were displayed on a solid support following separation
45 under denaturing conditions and Hu-Asp2 transcripts were visualized by high stringency
hybridization to radiolabeled insert from clone 2696295. The 2696295 cDNA probe
30 visualized a constellation of transcripts that migrated with apparent sizes of 3.0kb, 4.4 kb
and 8.0 kb with the latter two transcript being the most abundant.

Across the tissues surveyed, Hu-Asp2 transcripts were most abundant in pancreas and brain with lower but detectable levels observed in all other tissues examined except thymus and PBLs. Given the relative abundance of Hu-Asp2 transcripts in brain, the regional expression in brain regions was also established. A similar constellation of transcript sizes were detected in all brain regions examined [cerebellum, cerebral cortex, occipital pole, frontal lobe, temporal lobe and putamen] with the highest abundance in the medulla and spinal cord.

Example 5: Northern Blot Detection of HuAsp-1 and HuAsp-2 Transcripts in Human Cell Lines:

A variety of human cell lines were tested for their ability to produce Hu-Asp1 and Asp2 mRNA. Human embryonic kidney (HEK-293) cells, African green monkey (Cos-7) cells, Chinese hamster ovary (CHO) cells, HELA cells, and the neuroblastoma cell line IMR-32 were all obtained from the ATCC. Cells were cultured in DME containing 10% FCS except CHO cells which were maintained in α -MEM/10% FCS at 37 °C in 5% CO₂ until they were near confluence. Washed monolayers of cells (3×10^7) were lysed on the dishes and poly A⁺ RNA extracted using the Qiagen Oligotex Direct mRNA kit. Samples containing 2 μ g of poly A⁺ RNA from each cell line were fractionated under denaturing conditions (glyoxal-treated), transferred to a solid nylon membrane support by capillary action, and transcripts visualized by hybridization with random-primed labeled (³²P) coding sequence probes derived from either Hu-Asp1 or Hu-Asp2. Radioactive signals were detected by exposure to X-ray film and by image analysis with a PhosphorImager.

The Hu-Asp1 cDNA probe visualized a similar constellation of transcripts (2.6 kb and 3.5 kb) that were previously detected in human tissues. The relative abundance determined by quantification of the radioactive signal was Cos-7 > HEK 292 = HELA > IMR32.

The Hu-Asp2 cDNA probe also visualized a similar constellation of transcripts compared to tissue (3.0 kb, 4.4 kb, and 8.0 kb) with the following relative abundance: HEK 293 > Cos 7 > IMR32 > HELA.

Example 6: Modification of APP to increase A β processing for in vitro screening

Human cell lines that process A β peptide from APP provide a means to screen in cellular assays for inhibitors of β - and γ -secretase. Production and release of A β peptide into the culture supernatant is monitored by an enzyme-linked immunosorbent assay (EIA). Although expression of APP is widespread and both neural and non-neuronal cell lines

process and release A β peptide, levels of endogenous APP processing are low and difficult to detect by EIA. A β processing can be increased by expressing in transformed cell lines mutations of APP that enhance A β processing. We made the serendipitous observation that addition of two lysine residues to the carboxyl terminus of APP695 increases A β processing still further. This allowed us to create a transformed cell line that releases A β peptide into the culture medium at the remarkable level of 20,000 pg/ml.

Materials And Methods

Materials:

Human embryonic kidney cell line 293 (HEK293 cells) were obtained internally. The vector pIRES-EGFP was purchased from Clontech. Oligonucleotides for mutation using the polymerase chain reaction (PCR) were purchased from Genosys. A plasmid containing human APP695 (SEQ ID No. 9 [nucleotide] and SEQ ID No. 10 [amino acid]) was obtained from Northwestern University Medical School. This was subcloned into pSK (Stratagene) at the *NotI* site creating the plasmid pAPP695.

Mutagenesis protocol:

The Swedish mutation (K670N, M671L) was introduced into pAPP695 using the Stratagene Quick Change Mutagenesis Kit to create the plasmid pAPP695NL (SEQ ID No. 11 [nucleotide] and SEQ ID No. 12 [amino acid]). To introduce a di-lysine motif at the C-terminus of APP695, the forward primer #276 5' GACTGACCACTCGACCAGGTTC (SEQ ID No. 47) was used with the "patch" primer #274 5' CGAATTAAATTCCAGCACACTGGCTACTTCTTGTCTGCATCTCAAAGAAC (SEQ ID No. 48) and the flanking primer #275 CGAATTAAATTCCAGCACACTGGCTA (SEQ ID No. 49) to modify the 3' end of the APP695 cDNA (SEQ ID No. 15 [nucleotide] and SEQ ID No. 16 [amino acid]). This also added a BstX1 restriction site that will be compatible with the BstX1 site in the multiple cloning site of pIRES-EGFP. PCR amplification was performed with a Clontech HF Advantage cDNA PCR kit using the polymerase mix and buffers supplied by the manufacturer. For "patch" PCR, the patch primer was used at 1/20th the molar concentration of the flanking primers. PCR amplification products were purified using a QIAquick PCR purification kit (Qiagen). After digestion with restriction enzymes, products were separated on 0.8% agarose gels and then excised DNA fragments were purified using a QIAquick gel extraction kit (Qiagen).

To reassemble a modified APP695-Sw cDNA, the 5' *NotI*-*BglII* fragment of the APP695-Sw cDNA and the 3' *BglII*-*BstX1* APP695 cDNA fragment obtained by PCR were

ligated into pIRES-EGFP plasmid DNA opened at the NotI and BstXI sites. Ligations were performed for 5 minutes at room temperature using a Rapid DNA Ligation kit (Boehringer Mannheim) and transformed into Library Efficiency DH5a Competent Cells (GibcoBRL Life Technologies). Bacterial colonies were screened for inserts by PCR amplification using primers #276 and #275. Plasmid DNA was purified for mammalian cell transfection using a QIAprep Spin Miniprep kit (Qiagen). The construct obtained was designated pMG125.3 (APPSW-KK, SEQ ID No. 17 [nucleotide] and SEQ ID No. 18 [amino acid]).

Mammalian Cell Transfection:

HEK293 cells for transfection were grown to 80% confluence in Dulbecco's modified Eagle's medium (DMEM) with 10% fetal bovine serum. Cotransfections were performed using LipofectAmine (Gibco-BRL) with 3 µg pMG125.3 DNA and 9 µg pcDNA3.1 DNA per 10×10^6 cells. Three days posttransfection, cells were passaged into medium containing G418 at a concentration of 400 µg/ml. After three days growth in selective medium, cells were sorted by their fluorescence.

Clonal Selection of 125.3 cells by FACS:

Cell samples were analyzed on an EPICS Elite ESP flow cytometer (Coulter, Hialeah, FL) equipped with a 488 nm excitation line supplied by an air-cooled argon laser. EGFP emission was measured through a 525 nm band-pass filter and fluorescence intensity was displayed on a 4-decade log scale after gating on viable cells as determined by forward and right angle light scatter. Single green cells were separated into each well of one 96 well plate containing growth medium without G418. After a four day recovery period, G418 was added to the medium to a final concentration of 400 µg/ml. After selection, 32% of the wells contained expanding clones. Wells with clones were expanded from the 96 well plate to a 24 well plate and then a 6 well plate with the fastest growing colonies chosen for expansion at each passage. The final cell line selected was the fastest growing of the final six passaged. This clone, designated 125.3, has been maintained in G418 at 400 µg/ml with passage every four days into fresh medium. No loss of Aβ production or EGFP fluorescence has been seen over 23 passages.

Aβ EIA Analysis (Double Antibody Sandwich ELISA for hAβ 1-40/42):

Cell culture supernatants harvested 48 hr after transfection were analyzed in a standard Aβ EIA as follows. Human Aβ 1-40 or 1-42 was measured using monoclonal antibody (mAb) 6E10 (Senetek, St. Louis, MO) and biotinylated rabbit antiserum 162 or

164 (New York State Institute for Basic Research, Staten Island, NY) in a double antibody sandwich ELISA. The capture antibody 6E10 is specific to an epitope present on the N-terminal amino acid residues 1-16 of hA β . The conjugated detecting antibodies 162 and 164 are specific for hA β 1-40 and 1-42, respectively. Briefly, a Nunc Maxisorp 96 well immunoplate was coated with 100 μ l/well of mAb 6E10 (5 μ g/ml) diluted in 0.1M carbonate-bicarbonate buffer, pH 9.6 and incubated at 4°C overnight. After washing the plate 3x with 0.01M DPBS (Modified Dulbecco's Phosphate Buffered Saline (0.008M sodium phosphate, 0.002M potassium phosphate, 0.14M sodium chloride, 0.01 M potassium chloride, pH 7.4) from Pierce, Rockford, IL) containing 0.05% of Tween-20 (DPBST), the plate was blocked for 60 min with 200 μ l of 10% normal sheep serum (Sigma) in 0.01M DPBS to avoid non-specific binding. Human A β 1-40 or 1-42 standards (100 μ l/well (Bachem, Torrance, CA) diluted, from a 1mg/ml stock solution in DMSO, in culture medium was added after washing the plate, as well as 100 μ l/well of sample, e.g. conditioned medium of transfected cells. The plate was incubated for 2 hours at room temperature and 4°C overnight. The next day, after washing the plate, 100 μ l/well biotinylated rabbit antiserum 162 1:400 or 164 1:50 diluted in DPBST + 0.5% BSA was added and incubated at room temperature for 1hr 15 min. Following washes, 100 μ l/well neutravidin-horseradish peroxidase (Pierce, Rockford, IL) diluted 1:10,000 in DPBST was applied and incubated for 1 hr at room temperature. After the last washes 100 μ l/well of o-phenylenediamine dihydrochloride (Sigma Chemicals, St. Louis, MO) in 50mM citric acid/100mM sodium phosphate buffer (Sigma Chemicals, St. Louis, MO), pH 5.0, was added as substrate and the color development was monitored at 450nm in a kinetic microplate reader for 20 min. using Soft max Pro software. All standards and samples were run in triplicates. The samples with absorbance values falling within the standard curve were extrapolated from the standard curves using Soft max Pro software and expressed in pg/ml culture medium.

Results:

Addition of two lysine residues to the carboxyl terminus of APP695 greatly increases A β processing in HEK293 cells as shown by transient expression (Table 1).

Addition of the di-lysine motif to APP695 increases A β processing to that seen with the APP695 containing the Swedish mutation. Combining the di-lysine motif with the Swedish mutation further increases processing by an additional 2.8 fold.

Cotransformation of HEK293 cells with pMG125.3 and pcDNA3.1 allowed dual selection of transformed cells for G418 resistance and high level expression of EGFP. After clonal selection by FACS, the cell line obtained, produces a remarkable 20,000 pg A β peptide per ml of culture medium after growth for 36 hr in 24 well plates. Production of A β peptide under various growth conditions is summarized in Table 2.

TABLE 1. Release of A β peptide into the culture medium 48 hr after transient transfection of HEK293 cells with the indicated vectors containing wildtype or modified APP. Values tabulated are mean + SD and P-value for pairwise comparison using Student's t-test assuming unequal variances.

APP Construct	A β 1-40 peptide (pg/ml)	Fold Increase	P-value
pIRES-EGFP vector	147 + 28	1.0	
wt APP695 (142.3)	194 + 15	1.3	0.051
wt APP695-KK (124.1)	424 + 34	2.8	3 x 10 ⁻⁵
APP695-Sw (143.3)	457 + 65	3.1	2 x 10 ⁻³
APP695-SwKK (125.3)	1308 + 98	8.9	3 x 10 ⁻⁴

TABLE 2. Release of A β peptide from HEK125.3 cells under various growth conditions.

Type of Culture Plate	Volume of Medium	Duration of Culture	Ab 1-40 (pg/ml)	Ab 1-42 (pg/ml)
24 well plate	400 ul	36 hr	28,036	1,439

Example 7: Antisense oligomer inhibition of Abeta processing in HEK125.3 cells

The sequences of Hu-Asp1 and Hu-Asp2 were provided to Sequitur, Inc (Natick, MA) for selection of targeted sequences and design of 2nd generation chimeric antisense oligomers using proprietary technology (Sequitur Ver. D Pat pending #3002). Antisense oligomers Lot# S644, S645, S646 and S647 were targeted against Asp1. Antisense oligomers Lot# S648, S649, S650 and S651 were targeted against Asp2. Control antisense oligomers Lot# S652, S653, S655, and S674 were targeted against an irrelevant gene and antisense oligomers Lot #S656, S657, S658, and S659 were targeted against a second irrelevant gene.

For transfection with the antisense oligomers, HEK125.3 cells were grown to about 50% confluence in 6 well plates in Minimal Essential Medium (MEM) supplemented with 10% fetal calf serum. A stock solution of oligofectin G (Sequitur Inc., Natick, MA) at 2 mg/ml was diluted to 50 μ g/ml in serum free MEM. Separately, the antisense oligomer stock solution at 100 μ M was diluted to 800 nM in Opti-MEM (GIBCO-BRL, Grand

Island, NY). The diluted stocks of oligofectin G and antisense oligomer were then mixed at a ratio of 1:1 and incubated at room temperature. After 15 min incubation, the reagent was diluted 10 fold into MEM containing 10% fetal calf serum and 2 ml was added to each well of the 6 well plate after first removing the old medium. After transfection, cells were grown in the continual presence of the oligofectin G/antisense oligomer. To monitor Ab peptide release, 400 μ l of conditioned medium was removed periodically from the culture well and replaced with fresh medium beginning 24 hr after transfection. Data reported are from culture supernatants harvested 48 hr after transfection.

Results:

The 16 different antisense oligomers obtained from Sequitur Inc were transfected separately into HEK125.3 cells to determine their affect on A β peptide processing. Only antisense oligomers targeted against Asp1 & Asp2 reduced Abeta processing by HEK125.3 cells with those targeted against Asp2 having a greater inhibitory effect. Both A β (1-40) and A β (1-42) were inhibited by the same degree. In Table 3, percent inhibition is calculated with respect to untransfected cells. Antisense oligomer reagents giving greater than 50% inhibition are marked with an asterisk. Of the reagents tested, 3 of 4 antisense oligomers targeted against ASP1 gave an average 52% inhibition of A β 1-40 processing and 47% inhibition of A β 1-42 processing. For ASP2, 4 of 4 antisense oligomers gave greater than 50% inhibition with an average inhibition of 62% for A β 1-40 processing and 60% for A β 1-42 processing.

Table 3. Inhibition of A β peptide release from HEK125.3 cells treated with antisense oligomers.

Gene Targeted	Antisense Oligomer	Abeta (1-40)	Abeta (1-42)
Asp1-1	S 644	62%*	56%*
Asp1-2	S 645	41%*	38%*
Asp1-3	S646	52%*	46%*
Asp1-4	S647	6%	25%
Asp2-1	S648	71%*	67%*
Asp2-2	S649	83%*	76%*
Asp2-3	S650	46%*	50%*
Asp2-4	S651	47%*	46%*
Con1-1	S652	13%	18%
Con1-2	S653	35%	30%
Con1-3	S655	9%	18%
Con1-4	S674	29%	18%
Con2-1	S656	12%	18%
Con2-2	S657	16%	19%
Con2-3	S658	8%	35%

WO 00/17369

PCT/US99/20881

Con2-4

S659

3%

18%

5

10

15

20

25

30

35

40

45

50

55

Example 8. Demonstration of Hu-Asp2 β -Secretase Activity in Cultured Cells

Several mutations in APP associated with early onset Alzheimer's disease have been shown to alter A β peptide processing. These flank the N- and C-terminal cleavage sites that release A β from APP. These cleavage sites are referred to as the β -secretase and γ -secretase cleavage sites, respectively. Cleavage of APP at the β -secretase site creates a C-terminal fragment of APP containing 99 amino acids of 11,145 daltons molecular weight. The Swedish KM \rightarrow NL mutation immediately upstream of the β -secretase cleavage site causes a general increase in production of both the 1-40 and 1-42 amino acid forms of A β peptide. The London VF mutation (V717 \rightarrow F in the APP770 isoform) has little effect on total A β peptide production, but appears to preferentially increase the percentage of the longer 1-42 amino acid form of A β peptide by affecting the choice of γ -secretase cleavage site used during APP processing. Thus, we sought to determine if these mutations altered the amount and type of A β peptide produced by cultured cells cotransfected with a construct directing expression of Hu-Asp2.

Two experiments were performed which demonstrate Hu-Asp2 β -secretase activity in cultured cells. In the first experiment, treatment of HEK125.3 cells with antisense oligomers directed against Hu-Asp2 transcripts as described in Example 7 was found to decrease the amount of the C-terminal fragment of APP created by β -secretase cleavage (CTF99) (Figure 9). This shows that Hu-Asp2 acts directly or indirectly to facilitate β -secretase cleavage. In the second experiment, increased expression of Hu-Asp2 in transfected mouse Neuro2A cells is shown to increase accumulation of the CTF99 β -secretase cleavage fragment (Figure 10). This increase is seen most easily when a mutant APP-KK clone containing a C-terminal di-lysine motif is used for transfection. A further increase is seen when Hu-Asp2 is cotransfected with APP-Sw-KK containing the Swedish mutation KM \rightarrow NL. The Swedish mutation is known to increase cleavage of APP by the β -secretase.

5 A second set of experiments demonstrate Hu-Asp2 facilitates γ -secretase activity in
 cotransfection experiments with human embryonic kidney HEK293 cells. Cotransfection of
 Hu-Asp2 with an APP-KK clone greatly increases production and release of soluble A β 1-40
 and A β 1-42 peptides from HEK293 cells. There is a proportionately greater increase in the
 10 5 release of A β 1-42. A further increase in production of A β 1-42 is seen when Hu-Asp2 is
 cotransfected with APP-VF (SEQ ID No. 13 [nucleotide] and SEQ ID No. 14 [amino acid])
 or APP-VF-KK SEQ ID No. 19 [nucleotide] and SEQ ID No. 20 [amino acid]) clones
 15 containing the London mutation V717→F. The V717→F mutation is known to alter
 cleavage specificity of the APP γ -secretase such that the preference for cleavage at the A β 42
 20 site is increased. Thus, Asp2 acts directly or indirectly to facilitate γ -secretase processing of
 APP at the β 42 cleavage site.

Materials

Antibodies 6E10 and 4G8 were purchased from Senetek (St. Louis, MO). Antibody 369
 25 was obtained from the laboratory of Paul Greengard at the Rockefeller University.
 15 Antibody C8 was obtained from the laboratory of Dennis Selkoe at the Harvard Medical
 School and Brigham and Women's Hospital.

APP Constructs used

30 The APP constructs used for transfection experiments comprised the following

20	APP	wild-type APP695 (SEQ ID No. 9 and No. 10)
	APP-Sw	APP695 containing the Swedish KM→NL mutation (SEQ ID No. 11 and No. 12),
35	APP-VF	APP695 containing the London V→F mutation (SEQ ID No. 13 and No. 14)
25	APP-KK	APP695 containing a C-terminal KK motif (SEQ ID No. 15 and No. 16),
40	APP-Sw-KK	APP695-Sw containing a C-terminal KK motif (SEQ ID No. 17 and No. 18),
30	APP-VF-KK	APP695-VF containing a C-terminal KK motif (SEQ ID No. 19 and No. 20).

45 These were inserted into the vector pIRES-EGFP (Clontech, Palo Alto CA) between the
 NotI and BstXI sites using appropriate linker sequences introduced by PCR.

50 *Transfection of antisense oligomers or plasmid DNA constructs in HEK293 cells,
 35 HEK125.3 cells and Neuro-2A cells,*

Human embryonic kidney HEK293 cells and mouse Neuro-2a cells were transfected with expression constructs using the Lipofectamine Plus reagent from Gibco/BRL. Cells were seeded in 24 well tissue culture plates to a density of 70-80% confluence. Four wells per plate were transfected with 2 μ g DNA (3:1, APP:cotransfectant), 8 μ l Plus reagent, and 4 μ l Lipofectamine in OptiMEM. OptiMEM was added to a total volume of 1 ml, distributed 200 μ l per well and incubated 3 hours. Care was taken to hold constant the ratios of the two plasmids used for cotransfection as well as the total amount of DNA used in the transfection. The transfection media was replaced with DMEM, 10%FBS, NaPyruvate, with antibiotic/antimycotic and the cells were incubated under normal conditions (37°, 5% CO₂) for 48 hours. The conditioned media were removed to polypropylene tubes and stored at -80°C until assayed for the content of A β 1-40 and A β 1-42 by EIA as described in the preceding examples. Transfection of antisense oligomers into HEK125.3 cells was as described in Example 7.

Preparation of cell extracts, Western blot protocol

Cells were harvested after being transfected with plasmid DNA for about 60 hours. First, cells were transferred to 15-ml conical tube from the plate and centrifuged at 1,500 rpm for 5 min to remove the medium. The cell pellets were washed with PBS for one time. We then lysed the cells with lysis buffer (10 mM HEPES, pH 7.9, 150 mM NaCl, 10% glycerol, 1 mM EGTA, 1 mM EDTA, 0.1 mM sodium vanadate and 1% NP-40). The lysed cell mixtures were centrifuged at 5000 rpm and the supernatant was stored at -20°C as the cell extracts. Equal amounts of extracts from HEK125.3 cells transfected with the Asp2 antisense oligomers and controls were precipitated with antibody 369 that recognizes the C-terminus of APP and then CTF99 was detected in the immunoprecipitate with antibody 6E10. The experiment was repeated using C8, a second precipitating antibody that also recognizes the C-terminus of APP. For Western blot of extracts from mouse Neuro-2a cells cotransfected with Hu-Asp2 and APP-KK, APP-Sw-KK, APP-VF-KK or APP-VF, equal amounts of cell extracts were electrophoresed through 4-10% or 10-20% Tricine gradient gels (NOVEX, San Diego, CA). Full length APP and the CTF99 β -secretase product were detected with antibody 6E10.

Results

Transfection of HEK125.3 cells with Asp2-1 or Asp2-2 antisense oligomers reduces
production of the CTF β -secretase product in comparison to cells similarly transfected with
control oligomers having the reverse sequence (Asp2-1 reverse & Asp2-2 reverse)

In cotransfection experiments, cotransfection of Hu-Asp2 into mouse Neuro-2a cells with
the APP-KK construct increased the formation of CTF99. This was further increased if Hu-
Asp2 was coexpressed with APP-Sw-KK, a mutant form of APP containing the Swedish
KM \rightarrow NL mutation that increases β -secretase processing.

Cotransfection of Hu-Asp2 with APP has little effect on A β 40 production but
increases A β 42 production above background (Table 4). Addition of the di-lysine motif to
the C-terminus of APP increases A β peptide processing about two fold, although A β 40 and
A β 42 production remain quite low (352 pg/ml and 21 pg/ml, respectively). Cotransfection
of Asp2 with APP-KK further increases both A β 40 and A β 42 production. The stimulation
of A β 40 production by Hu-Asp2 is more than 3 fold, while production of A β 42 increases by
more than 10 fold. Thus, cotransfection of Hu-Asp2 and APP-KK constructs preferentially
increases A β 42 production.

The APP V717 \rightarrow F mutation has been shown to increase γ -secretase processing at the
A β 42 cleavage site. Cotransfection of Hu-Asp2 with the APP-VF or APP-VF-KK
constructs increased A β 42 production (a two fold increase with APP-VF and a four-fold
increase with APP-VF-KK, Table 4), but had mixed effects on A β 40 production (a slight
decrease with APP-VF, and a two fold increase with APP-VF-KK in comparison to the
pcDNA cotransfection control. Thus, the effect of Asp2 on A β 42 production was
proportionately greater leading to an increase in the ratio of A β 42/total A β . Indeed, the ratio
of A β 42/total A β reaches a very high value of 42% in HEK293 cells cotransfected with Hu-
Asp2 and APP-VF-KK.

Western blot showing reduction of CTF99 production by HEK125.3 cells transfected with antisense oligomers targeting the Hu-Asp2 mRNA. (right) Western blot showing increase in CTF99 production in mouse Neuro-2a cells cotransfected with Hu-Asp2 and APP-KK. A further increase in CTF99 production is seen in cells cotransfected with Hu-Asp2 and APP-Sw-KK.

Table 4. Results of cotransfecting Hu-Asp2 or pcDNA plasmid DNA with various APP constructs containing the V717→F mutation that modifies γ -secretase processing. Cotransfection with Asp2 consistently increases the ratio of A β 42/total A β . Values tabulated are A β peptide pg/ml.

	pcDNA Cotransfection			Asp2 Cotransfection		
	A β 40	A β 42	A β 42/Total	A β 40	A β 42	A β 42/Total
APP	192 \pm 18	<4	<2%	188 \pm 40	8 \pm 10	3.9%
APP-VF	118 \pm 15	15 \pm 19	11.5%	85 \pm 7	24 \pm 12	22.4%
APP-KK	352 \pm 24	21 \pm 6	5.5%	1062 \pm 101	226 \pm 49	17.5%
APP-VF-KK	230 \pm 31	88 \pm 24	27.7%	491 \pm 35	355 \pm 36	42%

Example 9. Bacterial expression of human Asp2L

Expression of recombinant Hu_Asp2L in E. coli.

Hu-Asp2L can be expressed in E. coli after addition of N-terminal sequences such as a T7 tag (SEQ ID No. 21 and No. 22) or a T7 tag followed by a caspase 8 leader sequence (SEQ ID No. 23 and No. 24). Alternatively, reduction of the GC content of the 5' sequence by site directed mutagenesis can be used to increase the yield of Hu-Asp2 (SEQ ID No. 25 and No. 26). In addition, Asp2 can be engineered with a proteolytic cleavage site (SEQ ID No. 27 and No. 28). To produce a soluble protein after expression and refolding, deletion of the transmembrane domain and cytoplasmic tail, or deletion of the membrane proximal region, transmembrane domain, and cytoplasmic tail is preferred.

Methods

PCR with primers containing appropriate linker sequences was used to assemble fusions of Asp2 coding sequence with N-terminal sequence modifications including a T7 tag (SEQ ID Nos. 21 and 22) or a T7-caspase 8 leader (SEQ ID Nos. 23 and 24). These constructs were cloned into the expression vector pet23a(+) [Novagen] in which a T7 promoter directs expression of a T7 tag preceding a sequence of multiple cloning sites. To clone Hu-Asp2 sequences behind the T7 leader of pet23a+, the following oligonucleotides were used for amplification of the selected Hu-Asp2 sequence:

#553=GTGGATCCACCCAGCACGGCATCCGGCTG (SEQ ID No. 35),

#554=GAAAGCTTTCATGACTCATCTGTCTGTGGAATGTTG (SEQ ID No. 36) which placed BamHI and HindIII sites flanking the 5' and 3' ends of the insert, respectively. The Asp2 sequence was amplified from the full length Asp2(b) cDNA cloned into pcDNA3.1 using the Advantage-GC cDNA PCR [Clontech] following the manufacturer's supplied protocol using annealing & extension at 68°C in a two-step PCR cycle for 25 cycles. The insert and vector were cut with BamHI and HindIII, purified by electrophoresis through an agarose gel, then ligated using the Rapid DNA Ligation kit [Boehringer Mannheim]. The ligation reaction was used to transform the E. coli strain JM109 (Promega) and colonies were picked for the purification of plasmid (Qiagen, Qiaprep minispin) and DNA sequence analysis. For inducible expression using induction with isopropyl b-D-thiogalactopyranoside (IPTG), the expression vector was transferred into E. coli strain BL21 (Statagene). Bacterial cultures were grown in LB broth in the presence of ampicillin at 100 ug/ml, and induced in log phase growth at an OD600 of 0.6-1.0 with 1 mM IPTG for 4 hour at 37°C. The cell pellet was harvested by centrifugation.

To clone Hu-Asp2 sequences behind the T7 tag and caspase leader (SEQ ID Nos. 23 and 24), the construct created above containing the T7-Hu-Asp2 sequence (SEQ ID Nos. 21 and 22) was opened at the BamHI site, and then the phosphorylated caspase 8 leader oligonucleotides #559=GATCGATGACTATCTCTGACTCTCCGCGTGAACAGGACG (SEQ ID No. 37), #560=GATCCGTCCTGTTACGCGGAGAGTCAGAGATAGTCATC (SEQ ID No. 38) were annealed and ligated to the vector DNA. The 5' overhang for each set of oligonucleotides was designed such that it allowed ligation into the BamHI site but not subsequent digestion with BamHI. The ligation reaction was transformed into JM109 as above for analysis of protein expression after transfer to E. coli strain BL21.

In order to reduce the GC content of the 5' terminus of asp2, a pair of antiparallel oligos were designed to change degenerate codon bases in 15 amino acid positions from G/C to A/T (SEQ ID Nos. 25 and 26). The new nucleotide sequence at the 5' end of asp2 did not change the encoded amino acid and was chosen to optimize E. Coli expression. The sequence of the sense linker is 5'

CGGCATCCGGCTGCCCCTGCGTAGCGGTCTGGGTGGTGCTCCACTGGGTCTGCG
TCTGCCCCGGGAGACCGACGAA G 3' (SEQ ID No. 39). The sequence of the antisense linker is : 5'

CTTCGTCCGGTCTCCCGGGGCAGACGCAGACCCAGTGGAGCACCACCCAGACCG
CTACGCAGGGGCAGCCGGATGCCG 3' (SEQ ID No. 40). After annealing the phosphorylated linkers together in 0.1 M NaCl-10 mM Tris, pH 7.4 they were ligated into unique Cla I and Sma I sites in Hu-Asp2 in the vector pTAC. For inducible expression using induction with isopropyl b-D-thiogalactopyranoside (IPTG), bacterial cultures were grown in LB broth in the presence of ampicillin at 100 ug/ml, and induced in log phase growth at an OD600 of 0.6-1.0 with 1 mM IPTG for 4 hour at 37°C. The cell pellet was harvested by centrifugation.

To create a vector in which the leader sequences can be removed by limited proteolysis with caspase 8 such that this liberates a Hu-Asp2 polypeptide beginning with the N-terminal sequence GSFV (SEQ ID Nos. 27 and 28), the following procedure was followed. Two phosphorylated oligonucleotides containing the caspase 8 cleavage site IETD, #571=5'

GATCGATGACTATCTCTGACTCTCCGCTGGACTCTGGTATCGAAACCGACG
(SEQ ID No. 41) and #572=

GATCCGTCGGTTTCGATACCAGAGTCCAGCGGAGAGTCAGAGATAGTCATC

(SEQ ID No. 42) were annealed and ligated into pET23a+ that had been opened with BamHI. After transformation into JM109, the purified vector DNA was recovered and orientation of the insert was confirmed by DNA sequence analysis. +, the following oligonucleotides were used for amplification of the selected Hu-Asp2 sequence:

#573=5'AAGGATCCTTTGTGGAGATGGTGGACAACCTG, (SEQ ID No. 43)

#554=GAAAGCTTTTCATGACTCATCTGTCTGTGGAATGTTG (SEQ ID No. 44) which placed BamHI and HindIII sites flanking the 5' and 3' ends of the insert, respectively. The Asp2 sequence was amplified from the full length Asp2 cDNA cloned into pcDNA3.1 using the Advantage-GC cDNA PCR [Clontech] following the manufacturer's supplied

5 protocol using annealing & extension at 68°C in a two-step PCR cycle for 25 cycles. The
insert and vector were cut with BamHI and HindIII, purified by electrophoresis through an
10 agarose gel, then ligated using the Rapid DNA Ligation kit [Boehringer Mannheim]. The
ligation reaction was used to transform the E. coli strain JM109 [Promega] and colonies
5 were picked for the purification of plasmid (Qiagen, Qiaprep minispin) and DNA sequence
analysis. For inducible expression using induction with isopropyl b-D-
thiogalactopyranoside (IPTG), the expression vector was transferred into E. coli strain
15 BL21 (Statagene). Bacterial cultures were grown in LB broth in the presence of ampicillin
at 100 ug/ml, and induced in log phase growth at an OD600 of 0.6-1.0 with 1 mM IPTG for
10 4 hour at 37°C. The cell pellet was harvested by centrifugation.

To assist purification, a 6-His tag can be introduced into any of the above constructs
20 following the T7 leader by opening the construct at the BamHI site and then ligating in the
annealed, phosphorylated oligonucleotides containing the six histidine sequence
#565=GATCGCATCATCACCATCACCATG (SEQ ID No. 45),
15 #566=GATCCATGGTGATGGTGATGATGC (SEQ ID No. 46). The 5' overhang for each
25 set of oligonucleotides was designed such that it allowed ligation into the BamHI site but
not subsequent digestion with BamHI.

Preparation of Bacterial Pellet:

30 36.34g of bacterial pellet representing 10.8L of growth was dispersed into a total
20 volume of 200ml using a 20mm tissue homogenizer probe at 3000 to 5000 rpm in 2M KCl,
0.1M Tris, 0.05M EDTA, 1mM DTT. The conductivity adjusted to about 193mMhos with
water.

35 After the pellet was dispersed, an additional amount of the KCl solution was added,
bringing the total volume to 500 ml. This suspension was homogenized further for about 3
25 minutes at 5000 rpm using the same probe. The mixture was then passed through a Rannie
40 high-pressure homogenizer at 10,000psi.

In all cases, the pellet material was carried forward, while the soluble fraction was
45 discarded. The resultant solution was centrifuged in a GSA rotor for 1hr. at 12,500 rpm. The
pellet was resuspended in the same solution (without the DTT) using the same tissue
30 homogenizer probe at 2,000 rpm. After homogenizing for 5 minutes at 3000 rpm, the
volume was adjusted to 500ml with the same solution, and spun for 1hr. at 12,500 rpm.
50 The pellet was then resuspended as before, but this time the final volume was adjusted to

1.5L with the same solution prior to homogenizing for 5 minutes. After centrifuging at the same speed for 30 minutes, this procedure was repeated. The pellet was then resuspended into about 150ml of cold water, pooling the pellets from the six centrifuge tubes used in the GSA rotor. The pellet has homogenized for 5 minutes at 3,000 rpm, volume adjusted to 250ml with cold water, then spun for 30 minutes. Weight of the resultant pellet was 17.75g.

Summary: Lysis of bacterial pellet in KCl solution, followed by centrifugation in a GSA rotor was used to initially prepare the pellet. The same solution was then used an additional three times for resuspension/homogenization. A final water wash/homogenization was then performed to remove excess KCl and EDTA.

Solubilization of rHuAsp2L:

A ratio of 9-10ml/gram of pellet was utilized for solubilizing the rHuAsp2L from the pellet previously described. 17.75g of pellet was thawed, and 150ml of 8M guanidine HCl, 5mM β ME, 0.1% DEA, was added. 3M Tris was used to titrate the pH to 8.6. The pellet was initially resuspended into the guanidine solution using a 20mm tissue homogenizer probe at 1000 rpm. The mixture was then stirred at 4°C for 1 hour prior to centrifugation at 12,500rpm for 1 hour in GSA rotor. The resultant supernatant was then centrifuged for 30min at 40,000 x g in an SS-34 rotor. The final supernatant was then stored at -20°C, except for 50ml.

Immobilized Nickel Affinity Chromatography of Solubilized rHuAsp2L:

The following solutions were utilized:

- A) 6M Guanidine HCl, 0.1M NaP, pH 8.0, 0.01M Tris, 5mM β ME, 0.5mM Imidazole
- A') 6M Urea, 20mM NaP, pH 6.80, 50mM NaCl
- B') 6M Urea, 20mM NaP, pH 6.20, 50mM NaCl, 12mM Imidazole
- C') 6M Urea, 20mM NaP, pH 6.80, 50mM NaCl, 300mM Imidazole

Note: Buffers A' and C' were mixed at the appropriate ratios to give intermediate concentrations of Imidazole.

The 50ml of solubilized material was combined with 50ml of buffer A prior to adding to 100-125ml Qiagen Ni-NTA SuperFlow (pre-equilibrated with buffer A) in a 5 x 10cm Bio-Rad econo column. This was shaken gently overnight at 4°C in the cold room.

Chromatography Steps:

- 1) Drained the resultant flow through.
- 2) Washed with 50ml buffer A (collecting into flow through fraction)
- 3) Washed with 250ml buffer A (wash 1)
- 4) Washed with 250ml buffer A (wash 2)
- 5) Washed with 250ml buffer A'

- 5 6) Washed with 250ml buffer B'
7) Washed with 250ml buffer A'
8) Eluted with 250ml 75mM Imidazole
9) Eluted with 250ml 150mM Imidazole (150-1)
5 10) Eluted with 250ml 150mM Imidazole (150-2)
11) Eluted with 250ml 300mM Imidazole (300-1)
10 12) Eluted with 250ml 300mM Imidazole (300-2)
13) Eluted with 250ml 300mM Imidazole (300-3)

10 Chromatography Results:

15 The rHuAsp eluted at 75mM Imidazole through 300mM Imidazole. The 75mM fraction, as well as the first 150mM Imidazole (150-1) fraction contained contaminating proteins as visualized on Coomassie Blue stained gels. Therefore, fractions 150-2 and 300-1 will be utilized for refolding experiments since they contained the greatest amount of protein (see

20 Coomassie Blue stained gel).

Refolding Experiments of rHuAsp2L:

Experiment 1:

25 Forty ml of 150-2 was spiked with 1M DTT, 3M Tris, pH 7.4 and DEA to a final concentration of 6mM, 50mM, and 0.1% respectively. This was diluted suddenly (while stirring) with 200ml of (4°C) cold 20mM NaP, pH 6.8, 150mM NaCl. This dilution gave a final Urea concentration of 1M. This solution remained clear, even if allowed to set open to the air at RT or at 4°C.

30 After setting open to the air for 4-5 hours at 4°C, this solution was then dialyzed overnight against 20mM NaP, pH 7.4, 150mM NaCl, 20% glycerol. This method effectively removes the urea in the solution without precipitation of the protein.

35 **Experiment 2:**

40 Some of the 150-2 eluate was concentrated 2x on an Amicon Centriprep, 10,000 MWCO, then treated as in Experiment 1. This material also stayed in solution, with no visible precipitation.

30

Experiment 3:

89ml of the 150-2 eluate was spiked with 1M DTT, 3M Tris, pH 7.4 and DEA to a final concentration of 6mM, 50mM, and 0.1% respectively. This was diluted suddenly (while stirring) with 445ml of (4°C) cold 20mM NaP, pH 6.8, 150mM NaCl. This solution appeared clear, with no apparent precipitation. The solution was removed to RT and stirred for 10 minutes prior to adding MEA to a final concentration of 0.1mM. This was stirred slowly at RT for 1hr. Cystamine and CuSO₄ were then added to final concentrations of 1mM and 10μM respectively. The solution was stirred slowly at RT for 10 minutes prior to being moved to the 4°C cold room and shaken slowly overnight, open to the air.

The following day, the solution (still clear, with no apparent precipitation) was centrifuged at 100,000 x g for 1 hour. Supernatants from multiple runs were pooled, and the bulk of the stabilized protein was dialyzed against 20mM NaP, pH 7.4, 150mM NaCl, 20% glycerol. After dialysis, the material was stored at -20°C.

Some (about 10ml) of the protein solution (still in 1M Urea) was saved back for biochemical analyses, and frozen at -20°C for storage.

Example 10. Expression of Hu-Asp2 and Derivatives in Insect Cells

Expression by baculovirus infection—The coding sequence of Hu-Asp2 and several derivatives were engineered for expression in insect cells using the PCR. For the full-length sequence, a 5'-sense oligonucleotide primer that modified the translation initiation site to fit the Kozak consensus sequence was paired with a 3'-antisense primer that contains the natural translation termination codon in the Hu-Asp2 sequence. PCR amplification of the pcDNA3.1(hygro)/Hu-Asp2 template (see Example 12). Two derivatives of Hu-Asp2 that delete the C-terminal transmembrane domain (SEQ ID No. 29 and No. 30) or delete the transmembrane domain and introduce a hexa-histidine tag at the C-terminus (SEQ ID No. 31 and No. 32) were also engineered using the PCR. The same 5'-sense oligonucleotide primer described above was paired with either a 3'-antisense primer that (1) introduced a translation termination codon after codon 453 (SEQ ID No. 3) or (2) incorporated a hexa-histidine tag followed by a translation termination codon in the PCR using pcDNA3.1(hygro)/Hu-Asp-2L as the template. In all cases, the PCR reactions were performed amplified for 15 cycles using *PwoI* DNA polymerase (Boehringer-Mannheim) as outlined by the supplier. The reaction products were digested to completion with *BamHI* and *NorI* and ligated to *BamHI* and *NorI* digested baculovirus transfer vector pVL1393 (Invitrogen). A portion of the ligations was used to transform competent *E. coli* DH5α cells

5 followed by antibiotic selection on LB-Amp. Plasmid DNA was prepared by standard alkaline lysis and banding in CsCl to yield the baculovirus transfer vectors pVL1393/Asp2, pVL1393/Asp2 Δ TM and pVL1393/Asp2 Δ TM(His)₆. Creation of recombinant baculoviruses and infection of sf9 insect cells was performed using standard methods.

10 5 *Expression by transfection*—Transient and stable expression of Hu-Asp2 Δ TM and Hu-Asp2 Δ TM(His)₆ in High 5 insect cells was performed using the insect expression vector pIZ/V5-His. The DNA inserts from the expression plasmids vectors pVL1393/Asp2, pVL1393/Asp2 Δ TM and pVL1393/Asp2 Δ TM(His)₆ were excised by double digestion with BamHI and NotI and subcloned into BamHI and NotI digested pIZ/V5-His using standard methods. The resulting expression plasmids, referred to as pIZ/Hu-Asp2 Δ TM and pIZ/Hu-Asp2 Δ TM(His)₆, were prepared as described above.

20 For transfection, High 5 insect cells were cultured in High Five serum free medium supplemented with 10 μ g/ml gentamycin at 27 °C in sealed flasks. Transfections were performed using High five cells, High five serum free media supplemented with 10 μ g/ml gentamycin, and InsectinPlus liposomes (Invitrogen, Carlsbad, CA) using standard methods.

30 For large scale transient transfections 1.2×10^7 high five cells were plated in a 150 mm tissue culture dish and allowed to attach at room temperature for 15-30 minutes. During the attachment time the DNA/liposome mixture was prepared by mixing 6 ml of serum free media, 60 μ g Asp2 Δ TM/pIZ (+/- His) DNA and 120 μ l of Insectin Plus and incubating at room temperature for 15 minutes. The plating media was removed from the dish of cells and replaced with the DNA/liposome mixture for 4 hours at room temperature with constant rocking at 2 rpm. An additional 6 ml of media was added to the dish prior to incubation for 4 days at 27 °C in a humid incubator. Four days post transfection the media was harvested, clarified by centrifugation at 500 x g, assayed for Asp2 expression by Western blotting. For stable expression, the cells were treated with 50 μ g/ml Zeocin and the surviving pool used to prepared clonal cells by limiting dilution followed by analysis of the expression level as noted above.

45 30 *Purification of Hu-Asp2 Δ TM and Hu-Asp2 Δ TM(His)₆*—Removal of the transmembrane segment from Hu-Asp2 resulted in the secretion of the polypeptide into the culture medium. Following protein production by either baculovirus infection or transfection, the conditioned medium was harvested, clarified by centrifugation, and dialyzed against Tris-HCl (pH 8.0). This material was then purified by successive

5 chromatography by anion exchange (Tris-HCl, pH 8.0) followed by cation exchange
chromatography (Acetate buffer at pH 4.5) using NaCl gradients. The elution profile was
monitored by (1) Western blot analysis and (2) by activity assay using the peptide substrate
described in Example 12. For the Hu-Asp2ΔTM(His)₆, the conditioned medium was
10 5 dialyzed against Tris buffer (pH 8.0) and purified by sequential chromatography on IMAC
resin followed by anion exchange chromatography.

15 Sequence analysis of the purified Hu-Asp2ΔTM(His)₆ protein revealed that the signal
peptide had been cleaved [TQHGIRLPLR].

10 Example 11. Expression of Hu-Asp2 in CHO cells

20 *Heterologous expression of Hu-Asp-2L in CHO-K1 cells*—The entire coding sequence of
Hu-Asp2 was cloned into the mammalian expression vector pcDNA3.1(+)/Hygro
(Invitrogen, Carlsbad, CA) which contains the CMV immediate early promoter and bGH
25 15 polyadenylation signal to drive over expression. The expression plasmid,
pcDNA3.1(+)/Hygro/Hu-Asp2, was prepared by alkaline lysis and banding in CsCl and
completely sequenced on both strands to verify the integrity of the coding sequence.

30 20 Wild-type Chinese hamster ovary cells (CHO-K1) were obtained from the ATCC. The
cells were maintained in monolayer cultures in α-MEM containing 10% FCS at 37°C in 5%
35 CO₂. Two 100 mm dishes of CHO-K1 cells (60% confluent) were transfected with
pcDNA3.1(+)/Hygro alone (mock) or pcDNA3.1(+)/Hygro/Hu-Asp2 using the cationic
liposome DOTAP as recommended by the supplier. The cells were treated with the plasmid
40 25 DNA/liposome mixtures for 15 hr and then the medium replaced with growth medium
containing 500 Units/ml hygromycin B. In the case of pcDNA3.1(+)/Hygro/Hu-Asp2
45 transfected CHO-K1 cells, individual hygromycin B-resistant cells were cloned by limiting
dilution. Following clonal expansion of the individual cell lines, expression of Hu-Asp2
protein was accessed by Western blot analysis using a polyclonal rabbit antiserum raised

5 against recombinant Hu-Asp2 prepared by expression in *E. coli*. Near confluent dishes of
each cell line were harvested by scraping into PBS and the cells recovered by
centrifugation. The cell pellets were resuspended in cold lysis buffer (25 mM Tris-HCl
10 (8.0)/5 mM EDTA) containing protease inhibitors and the cells lysed by sonication. The
soluble and membrane fractions were separated by centrifugation (105,000 x g, 60 min) and
15 normalized amounts of protein from each fraction were then separated by SDS-PAGE.
Following electrotransfer of the separated polypeptides to PVDF membranes, Hu-Asp-2L
protein was detected using rabbit anti-Hu-Asp2 antiserum (1/1000 dilution) and the
20 antibody-antigen complexes were visualized using alkaline phosphatase conjugated goat
anti-rabbit antibodies (1/2500). A specific immunoreactive protein with an apparent Mr
value of 65 kDa was detected in pcDNA3.1(+)/Hygro/Hu-Asp2 transfected cells and not
25 mock-transfected cells. Also, the Hu-Asp2 polypeptide was only detected in the membrane
fraction, consistent with the presence of a signal peptide and single transmembrane domain
in the predicted sequence. Based on this analysis, clone #5 had the highest expression level
30 of Hu-Asp2 protein and this production cell lines was scaled up to provide material for
purification.

35 *Purification of recombinant Hu-Asp-2L from CHO-K1/Hu-Asp2 clone #5*—In a
typical purification, clone #5 cell pellets derived from 20 150 mm dishes of confluent cells,
were used as the starting material. The cell pellets were resuspended in 50 ml cold lysis
40 buffer as described above. The cells were lysed by polytron homogenization (2 x 20 sec)
and the lysate centrifuged at 338,000 x g for 20 minutes. The membrane pellet was then
resuspended in 20 ml of cold lysis buffer containing 50 mM β -octylglucoside followed by
45 rocking at 4°C for 1hr. The detergent extract was clarified by centrifugation at 338,000 x g
for 20 minutes and the supernatant taken for further analysis.

5 The β -octylglucoside extract was applied to a Mono Q anion exchange column that
was previously equilibrated with 25 mM Tris-HCl (pH 8.0)/50 mM β -octylglucoside.
Following sample application, the column was eluted with a linear gradient of increasing
10 NaCl concentration (0-1.0 M over 30 minutes) and individual fractions assayed by Western
blot analysis and for β -secretase activity (see below). Fractions containing both Hu_Asp-
2L immunoreactivity and β -secretase activity were pooled and dialyzed against 25 mM
15 NaOAc (pH 4.5)/50 mM β -octylglucoside. Following dialysis, precipitated material was
removed by centrifugation and the soluble material chromatographed on a MonoS cation
exchange column that was previously equilibrated in 25 mM NaOAc (pH 4.5)/ 50 mM β -
20 octylglucoside. The column was eluted using a linear gradient of increasing NaCl
concentration (0-1.0 M over 30 minutes) and individual fractions assayed by Western blot
25 analysis and for β -secretase activity. Fractions containing both Hu_Asp2 immunoreactivity
and β -secretase activity were combined and determined to be >90% pure by SDS-
PAGE/Coomassie Blue staining.

30 **Example 12. Assay of Hu_Asp2 β -secretase activity using peptide substrates**

15 *β -secretase assay*— β -secretase activity was measured by quantifying the hydrolysis of a
synthetic peptide containing the APP Swedish mutation by RP-HPLC with UV detection.
35 Each reaction contained 50 mM Na-MES (pH 5.5), 1% β -octylglucoside, peptide substrate
(SEVNLDAEFR, 70 μ M) and enzyme (1-5 μ g protein). Reactions were incubated at 37 °C
40 for various times and the reaction products were resolved by RP-HPLC using a linear
gradient from 0-70 B over 30 minutes (A=0.1% TFA in water,
45 B=).1%TFA/10%water/90%AcCN). The elution profile was monitored by absorbance at
214 nm. In preliminary experiments, the two product peaks which eluted before the intact
peptide substrate, were confirmed to have the sequence DAEFR and SEVNL using both
50

5 Edman sequencing and MADLI-TOF mass spectrometry. Percent hydrolysis of the peptide
substrate was calculated by comparing the integrated peak areas for the two product
10 peptides and the starting material derived from the absorbance at 214 nm. The specificity
of the protease cleavage reaction was determined by performing the β -secretase assay in the
5 presence of a cocktail of protease inhibitors (8 μ M pepstatin A, 10 μ M leupeptin, 10 μ M
E64, and 5 mM EDTA).

15 An alternative β -secretase assay utilizes internally quenched fluorescent substrates
to monitor enzyme activity using fluorescence spectroscopy in a single sample or multiwell
format. Each reaction contained 50 mM Na-MES (pH 5.5), peptide substrate MCA-
20 EVKMDAEF[K-DNP] (BioSource International) (50 μ M) and purified Hu-Asp-2 enzyme.
These components were equilibrated to 37 °C for various times and the reaction initiated by
addition of substrate. Excitation was performed at 330 nm and the reaction kinetics were
25 monitored by measuring the fluorescence emission at 390 nm. To detect compounds that
modulate Hu-Asp-2 activity, the test compounds were added during the preincubation phase
15 of the reaction and the kinetics of the reaction monitored as described above. Activators are
scored as compounds that increase the rate of appearance of fluorescence while inhibitors
30 decrease the rate of appearance of fluorescence.

It will be clear that the invention may be practiced otherwise than as particularly
described in the foregoing description and examples.

20 Numerous modifications and variations of the present invention are possible in light of the
above teachings and, therefore, are within the scope of the invention.

The entire disclosure of all publications cited herein are hereby incorporated by reference.

Claims

5

10

15

20

25

30

35

40

45

50

55

What is claimed is:

1. Any isolated or purified nucleic acid polynucleotide that codes for a protease capable of cleaving the beta (β) secretase cleavage site of APP that contains two or more sets of special nucleic acids, where the special nucleic acids are separated by nucleic acids that code for about 100 to 300 amino acid positions, where the amino acids in those positions may be any amino acids, where the first set of special nucleic acids consists of the nucleic acids that code for the peptide DTG, where the first nucleic acid of the first special set of nucleic acids is, the first special nucleic acid, and where the second set of nucleic acids code for either the peptide DSG or DTG, where the last nucleic acid of the second set of nucleic acids is the last special nucleic acid, with the proviso that the nucleic acids disclosed in SEQ ID NO. 1 and SEQ. ID NO. 5 are not included.
2. The nucleic acid polynucleotide of claim 1 where the two sets of nucleic acids are separated by nucleic acids that code for about 125 to 222 amino acid positions, which may be any amino acids.
3. The nucleic acid polynucleotide of claim 2 that code for about 150 to 172 amino acid positions, which may be any amino acids.
4. The nucleic acid polynucleotide of claim that code for about 172 amino acid positions, which may be any amino acids.
5. The nucleic acid polynucleotide of claim 4 where the nucleotides are described in SEQ. ID. NO. 3
6. The nucleic acid polynucleotide of claim 2 where the two sets of nucleic acids are separated by nucleic acids that code for about 150 to 196 amino acid positions.
7. The nucleic acid polynucleotide of claim 6 where the two sets of nucleotides are separated by nucleic acids that code for about 196 amino acids (positions).

- 5 8. The nucleic acid polynucleotide of claim 7 where the two sets of nucleic acids are separated by the same nucleic acid sequences that separate the same set of special nucleic acids in SEQ. ID. NO. 5.
- 10 9. The nucleic acid polynucleotide of claim 4 where the two sets of nucleic acids are separated by nucleic acids that code for about 150 to 190, amino acid (positions).
- 15 10. The nucleic acid polynucleotide of claim 9 where the two sets of nucleotides are separated by nucleic acids that code for about 190 amino acids (positions).
- 10 11. The nucleic acid polynucleotide of claim 10 where the two sets of nucleotides are separated by the same nucleic acid sequences that separate the same set of special nucleotides in SEQ. ID. NO. 1.
- 20 12. Claims 1-11 where the first nucleic acid of the first special set of amino acids, that is, the first special nucleic acid, is operably linked to any codon where the nucleic acids of that codon codes for any peptide comprising from 1 to 10,000 amino acid (positions).
- 25 13. The nucleic acid polynucleotide of claims 1-12 where the first special nucleic acid is operably linked to nucleic acid polymers that code for any peptide selected from the group consisting of: any any reporter proteins or proteins which facilitate purification.
- 30 14. The nucleic acid polynucleotide of claims 1-13 where the first special nucleic acid is operably linked to nucleic acid polymers that code for any peptide selected from the group consisting of: immunoglobulin-heavy chain, maltose binding protein, glutathion S transfection, Green Fluorescent protein, and ubiquitin.
- 35 15. Claims 1-14 where the last nucleic acid of the second set of special amino acids, that is, the last special nucleic acid, is operably linked to nucleic acid polymers that code for any peptide comprising any amino acids from 1 to 10,000 amino acids.
- 40
- 45
- 50
- 55

- 5 16. Claims 1-15 where the last special nucleic acid is operably linked to any codon linked to nucleic acid polymers that code for any peptide selected from the group consisting of: any reporter proteins or proteins which facilitate purification.
- 10 17. The nucleic acid polynucleotide of claims 1-16 where the first special nucleic acid is operably linked to nucleic acid polymers that code for any peptide selected from the group consisting of: immunoglobulin-heavy chain, maltose binding protein, glutathion S transfection, Green Fluorescent protein, and ubiquitin.
- 15 18. * Any isolated or purified nucleic acid polynucleotide that codes for a protease capable of cleaving the beta secretase cleavage site of APP that contains two or more sets of special nucleic acids, where the special nucleic acids are separated by nucleic acids that code for about 100 to 300 amino acid positions, where the amino acids in those positions may be any amino acids, where the first set of special
- 20 nucleic acids consists of the nucleic acids that code for DTG, where the first nucleic acid of the first special set of nucleic acids is, the first special nucleic acid, and where the second set of nucleic acids code for either DSG or DTG, where the last nucleic acid of the second set of special nucleic acids is the last special nucleic acid, where the first special nucleic acid is operably linked to nucleic acids that code for
- 25 any number of amino acids from zero to 81 amino acids and where each of those codons may code for any amino acid.
- 30 19. The nucleic acid polynucleotide of claim 18, where the first special nucleic acid is operably linked to nucleic acids that code for any number of from 64 to 77 amino acids where each codon may code for any amino acid.
- 35 20. The nucleic acid polynucleotide of claim 19, where the first special nucleic acid is operably linked to nucleic acids that code for about 71 amino acids peptide.
- 40 21. The nucleic acid polynucleotide of claim 20, where the first special nucleic acid is operably linked to 71 amino acid peptide and where the first of those 71 amino acids is the amino acid T.
- 45 50 55

- 5 22. The nucleic acid polynucleotide of claim 21, where the polynucleotide comprises a sequence that is at least 95% identical to the same corresponding amino acids in SEQ. ID. NO. 3, that is, identical to the sequences in SEQ. ID. NO. 3 including the sequences from both the first and or the second special nucleic acids, toward the N-
10 5 Terminal, through and including 71 amino acids, see Example 10, beginning from the DTG site and including the nucleotides from that code for 71 amino acids).
- 15 23. The nucleic acid polynucleotide of claim 22, where the complete polynucleotide comprises identical to the same corresponding amino acids in SEQ. ID. NO. 3, that
10 is, identical to the sequences in SEQ. ID. NO. 3 including the sequences from both the first and or the second special nucleic acids, toward the N-Terminal, through
20 and including 71 amino acids, see Example 10, beginning from the DTG site and including the nucleotides from that code for 71 amino acids).
- 25 24. The nucleic acid polynucleotide of claim 18, where the first special nucleic acid is operably linked to nucleic acids that code for any number of from about 30 to 54 amino acids where each codon may code for any amino acid.
- 30 25. The nucleic acid polynucleotide of claim 20, where the first special nucleic acid is operably linked to 47 codons where the first those 35 or 47 amino acids is the
20 amino acid E or G.
- 35 26. The nucleic acid polynucleotide of claim 21, where the polynucleotide comprises a sequence that is at least 95% identical to the same corresponding amino acids in
25 SEQ. ID. NO. 3, that is, identical to that portion of the sequences in SEQ. ID. NO. 3 including the sequences from both the first and or the second special nucleic acids, toward the N-Terminal, through and including 35 or 47 amino acids, see
40 Example 11 for the 47 example, beginning from the DTG site and including the nucleotides from that code for the previous 35 or 47 amino acids before the DTG
45 30 site).

- 5 27. The nucleic acid polynucleotide of claim 22, where the polynucleotide comprises identical to the same corresponding amino acids in SEQ. ID. NO. 3, that is, identical to the sequences in SEQ. ID. NO. 3 including the sequences from both the first and or the second special nucleic acids, toward the N-Terminal, through and including 35 or 47 amino acids, see Example 11 for the 47 example, beginning from the DTG site and including the nucleotides from that code for the previous 35 or 47 amino acids before the DTG site).
- 10 28. * Any isolated or purified nucleic acid polynucleotide that codes for a protease capable of cleaving the beta (β) secretase cleavage site of APP that contains two or more sets of special nucleic acids, where the special nucleic acids are separated by nucleic acids that code for about 100 to 300 amino acid positions, where the amino acids in those positions may be any amino acids, where the first set of special nucleic acids consists of the nucleic acids that code for the peptide DTG, where the first nucleic acid of the first special set of amino acids is, the first special nucleic acid, and where the second set of special nucleic acids code for either the peptide DSG or DTG, where the last nucleic acid of the second set of special nucleic acids, the last special nucleic acid, is operably linked to nucleic acids that code for any number of codons from 50 to 170 codons.
- 15 29. The nucleic acid polynucleotide of claim 29 where the last special nucleic acid is operably linked to nucleic acids comprising from 100 to 170 codons.
- 20 30. The nucleic acid polynucleotide of claim 30 where the last special nucleic acid is operably linked to nucleic acids comprising from 142 to 163 codons.
- 25 31. The nucleic acid polynucleotide of claim 31 where the last special nucleic acid is operably linked to nucleic acids comprising about 142 codons.
- 30 32. The nucleic acid polynucleotide of claim 32 where the polynucleotide comprises a sequence that is at least 95% identical to SEQ. ID. # (Example 9 or 10).

- 5 33. The nucleic acid polynucleotide of claim 33, where the complete polynucleotide comprises SEQ. ID. # (Example 9 or 10).
- 10 34. The nucleic acid polynucleotide of claim 31 where the last special nucleic acid is operably linked to nucleic acids comprising about 163 codons.
- 15 35. The nucleic acid polynucleotide of claim 35 where the polynucleotide comprises a sequence that is at least 95% identical to SEQ. ID. # (Example 9 or 10).
- 20 36. The nucleic acid polynucleotide of claim 36, where the complete polynucleotide comprises SEQ. ID. # (Example 9 or 10).
- 25 37. The nucleic acid polynucleotide of claim 31 where the last special nucleic acid is operably linked to nucleic acids comprising about 170 codons.
- 30 38. Claims 1-38 where the second set of special nucleic acids code for the peptide DSG, and optionally the first set of nucleic acid polynucleotide is operably linked to a peptide purification tag.
- 35 39. Claims 1-39 where the nucleic acid polynucleotide is operably linked to a peptide purification tag which is six histidine.
- 40 40. Claims 1-40 where the first set of special nucleic acids are on one polynucleotide and the second set of special nucleic acids are on a second polynucleotide, where both first and second polynucleotides have at least 50 codons.
- 45 41. Claims 1-40 where the first set of special nucleic acids are on one polynucleotide and the second set of special nucleic acids are on a second polynucleotide, where both first and second polynucleotides have at least 50 codons where both said polynucleotides are in the same solution.
- 50 42. A vector which contains a polynucleotide described in claims 1-42.

5 43. A cell or cell line which contains a polynucleotide described in claims 1-42.

10 44. Any isolated or purified peptide or protein comprising an amino acid polymer that is a protease capable of cleaving the beta (β) secretase cleavage site of APP that
5 contains two or more sets of special amino acids, where the special amino acids are separated by about 100 to 300 amino acid positions, where each amino acid position can be any amino acid, where the first set of special amino acids consists of the peptide DTG, where the first amino acid of the first special set of amino acids is, the first special amino acid, where the second set of amino acids is selected from the
15 peptide comprising either DSG or DTG, where the last amino acid of the second set of special amino acids is the last special amino acid, with the proviso that the proteases disclosed in SEQ ID NO. 2 and SEQ. ID NO. 6 are not included.

20 45. The amino acid polypeptide of claim 45 where the two sets of amino acids are separated by about 125 to 222 amino acid positions where in each position it may be any amino acid.

25 46. The amino acid polypeptide of claim 46 where the two sets of amino acids are separated by about 150 to 172 amino acids.

30 47. The amino acid polypeptide of claim 47 where the two sets of amino acids are separated by about 172 amino acids.

35 48. The amino acid polypeptide of claim 48 where the protease is described in SEQ. ID. NO. 4

40 49. The amino acid polypeptide of claim 46 where the two sets of amino acids are separated by about 150 to 196 amino acids.

45 50. The amino acid polypeptide of claim 50 where the two sets of amino acids are separated by about 196 amino acids.

- 5 51. The amino acid polypeptide of claim 51 where the two sets of amino acids are separated by the same amino acid sequences that separate the same set of special amino acids in SEQ. ID. NO. 6.
- 10 52. The amino acid polypeptide of claim 46 where the two sets of amino acids are separated by about 150 to 190, amino acids.
- 15 53. The amino acid polypeptide of claim 53 where the two sets of nucleotides are separated by about 190 amino acids.
- 10 54. The amino acid polypeptide of claim 54 where the two sets of nucleotides are separated by the same amino acid sequences that separate the same set of special amino acids in SEQ. ID. NO. 2.
- 20 55. Claims 45-55 where the first amino acid of the first special set of amino acids, that is, the first special amino acid, is operably linked to any peptide comprising from 1 to 10,000 amino acids.
- 25 56. The amino acid polypeptide of claims 45-56 where the first special amino acid is operably linked to any peptide selected from the group consisting of: any any reporter proteins or proteins which facilitate purification.
- 30 57. The amino acid polypeptide of claims 45-57 where the first special amino acid is operably linked to any peptide selected from the group consisting of:
25 immunoglobulin-heavy chain, maltose binding protein, glutathion S transfection, Green Fluorescent protein, and ubiquitin.
- 35 58. Claims 45-58, where the last amino acid of the second set of special amino acids, that is, the last special amino acid, is operably linked to any peptide comprising any
40 amino acids from 1 to 10,000 amino acids.
- 45
- 50
- 55

59. Claims 45-59 where the last special amino acid is operably linked any peptide selected from the group consisting of any reporter proteins or proteins which facilitate purification.
60. The amino acid polypeptide of claims 45-60 where the first special amino acid is operably linked to any peptide selected from the group consisting of: immunoglobulin-heavy chain, maltose binding protein, glutathion S transfection. Green Fluorescent protein, and ubiquitin.
61. * Any isolated or purified peptide or protein comprising an amino acid polypeptide that codes for a protease capable of cleaving the beta secretase cleavage site of APP that contains two or more sets of special amino acids, where the special amino acids are separated by about 100 to 300 amino acid positions, where each amino acid in each position can be any amino acid, where the first set of special amino acids consists of the amino acids DTG, where the first amino acid of the first special set of amino acids is, the first special amino acid, D, and where the second set of amino acids is either DSG or DTG, where the last amino acid of the second set of special amino acids is the last special amino acid, G, where the first special amino acid is operably linked to amino acids that code for any number of amino acids from zero to 81 amino acid positions where in each position it may be any amino acid.
62. The amino acid polypeptide of claim 62, where the first special amino acid is operably linked to a peptide from about 30 to 77 amino acids positions where each amino acid position may be any amino acid.
63. The amino acid polypeptide of claim 63, where the first special amino acid is operably linked to a peptide of 35, 47, 71, or 77 amino acids.
64. The amino acid polypeptide of claim 63, where the first special amino acid is operably linked to the same corresponding peptides from SEQ. ID. NO. 3 that are 35, 47, 71, or 77 peptides in length, beginning counting with the amino acids on the first special sequence, DTG, towards the N-terminal of SEQ. ID. NO. 3.

5 65. The amino acid polypeptide of claim 65, where the polypeptide comprises a
sequence that is at least 95% identical to the same corresponding amino acids in
SEQ. ID. NO. 4, that is, identical to that portion of the sequences in SEQ.ID. NO. 4,
10 5 including all the sequences from both the first and or the second special nucleic
acids, toward the N- terminal, through and including 71, 47, 35 amino acids before
the first special amino acids. (Examples 10 and 11).

15 66. The amino acid polypeptide of claim 65, where the complete polypeptide comprises
10 the peptide of 71 amino acids, where the first of the amino acid is T and the second
is Q.

20 67. The amino acid polypeptide of claim 62, where the first special amino acid is
operably linked to any number of from 40 to 54 amino acids (positions) where each
15 amino acid position may be any amino acid.

25 68. The amino acid polypeptide of claim 68, where the first special amino acid is
operably linked to amino acids that code for a peptide of 47 amino acids.

30 69. The amino acid polypeptide of claim 69, where the first special amino acid is
operably linked to a 47 amino acid peptide where the first those 47 amino acids is
the amino acid E.

35 70. The amino acid polypeptide of claim 70, where the polypeptide comprises a
25 sequence that is at least 95% identical to SEQ. ID. # (Example 10).

40 71. The amino acid polypeptide of claim 71, where the complete polypeptide comprises
SEQ. ID. # (Example 10).

45 30 72. * Any isolated or purified amino acid polypeptide that is a protease capable of
cleaving the beta (β) secretase cleavage site of APP that contains two or more sets
of special amino acids, where the special amino acids are separated by about 100 to
50 300 amino acid positions, where each amino acid in each position can be any amino

acid, where the first set of special amino acids consists of the amino acids that code for DTG, where the first amino acid of the first special set of amino acids is, the first special amino acid, D, and where the second set of amino acids are either DSG or DTG, where the last amino acid of the second set of special amino acids is the last special amino acid, G, which is operably linked to any number of amino acids from 50 to 170 amino acids, which may be any amino acids.

73. The amino acid polypeptide of claim 73 where the last special amino acid is operably linked to a peptide of about 100 to 170 amino acids.

74. The amino acid polypeptide of claim 74 where the last special amino acid is operably linked to to a peptide of about 142 to 163 amino acids.

75. The amino acid polypeptide of claim 75 where the last special amino acid is operably linked to to a peptide of about about 142 amino acids.

76. The amino acid polypeptide of claim 76 where the polypeptide comprises a sequence that is at least 95% identical to SEQ. ID. # (Example 9 or 10).

77. The amino acid polypeptide of claim 75 where the last special amino acid is operably linked to a peptide of about 163 amino acids.

78. The amino acid polypeptide of claim 79 where the polypeptide comprises a sequence that is at least 95% identical to SEQ. ID. # (Example 9 or 10).

79. The amino acid polypeptide of claim 79, where the complete polypeptide comprises SEQ. ID. # (Example 9 or 10).

80. The amino acid polypeptide of claim 74 where the last special amino acid is operably linked to to a peptide of about 170 amino acids.

81. Claim 46-81 where the second set of special amino acids is comprised of the peptide with the amino acid sequence DSG.

5 82. Claims 45-82 where the amino acid polypeptide is operably linked to a peptide purification tag.

10 5 83. Claims 45-83 where the amino acid polypeptide is operably linked to a peptide purification tag which is six histidine.

15 84. Claims 45-84 where the first set of special amino acids are on one polypeptide and the second set of special amino acids are on a second polypeptide, where both first and second polypeptide have at least 50 amino acids, which may be any amino acids.

20 85. Claims 45-84 where the first set of special amino acids are on one polypeptide and the second set of special amino acids are on a second polypeptide, where both first and second polypeptides have at least 50 amino acids where both said polypeptides are in the same vessel.

25 86. A vector which contains a polypeptide described in claims 45-86.

30 87. A cell or cell line which contains a polynucleotide described in claims 45-87.

35 88. The process of making any of the polynucleotides, vectors, or cells of claims 1-44

40 89. The process of making any of the polypeptides, vectors or cells of claims 45-88

25 90. Any of the polynucleotides, polypeptides, vectors, cells or cell lines described in claims 1-88 made from the processes described in claims 89 and 90.

45 30 91. * An isolated nucleic acid molecule comprising a polynucleotide, said polynucleotide encoding a Hu-Asp polypeptide and having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:

5 (a) a nucleotide sequence encoding a Hu-Asp polypeptide selected from the group consisting of Hu-Asp1, Hu-Asp2(a), and Hu-Asp2(b), wherein said Hu-Asp1, Hu-Asp2(a) and Hu-Asp2(b) polypeptides have the complete amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, and SEQ ID No:6, respectively; and

10 5 (b) a nucleotide sequence complementary to the nucleotide sequence of (a).

15 92. The nucleic acid molecule of claim 92, wherein said Hu-Asp polypeptide is Hu-Asp1, and said polynucleotide molecule of 1(a) comprises the nucleotide sequence of SEQ ID NO:1.

20 93. The nucleic acid molecule of claim 92, wherein said Hu-Asp polypeptide is Hu-Asp2(a), and said polynucleotide molecule of 1(a) comprises the nucleotide sequence of SEQ ID NO:4.

25 94. The nucleic acid molecule of claim 92, wherein said Hu-Asp polypeptide is Hu-Asp2(b), and said polynucleotide molecule of 1(a) comprises the nucleotide sequence of SEQ ID NO:5.

30 20 95. An isolated nucleic acid molecule comprising polynucleotide which hybridizes under stringent conditions to a polynucleotide having the nucleotide sequence in (a) or (b) of claim 92.

35 96. A vector comprising the nucleic acid molecule of claim 96.

40 25 97. The vector of claim 97, wherein said nucleic acid molecule is operably linked to a promoter for the expression of a Hu-Asp polypeptide.

45 98. The vector of claim 97, wherein said Hu-Asp polypeptide is Hu-Asp1.

50 30 99. The vector of claim 97, wherein said Hu-Asp polypeptide is Hu-Asp2(a).

100. The vector of claim 97, wherein said Hu-Asp polypeptide is Hu-Asp2(b).

5 101. A host cell comprising the vector of claim 98.

10 102. A method of obtaining a Hu-Asp polypeptide comprising culturing the host cell of claim 102 and isolating said Hu-Asp polypeptide.

15 103. An isolated Hu-Asp1 polypeptide comprising an amino acid sequence at least 95% identical to a sequence comprising the amino acid sequence of SEQ ID NO:2.

20 104. An isolated Hu-Asp2(a) polypeptide comprising an amino acid sequence at least 95% identical to a sequence comprising the amino acid sequence of SEQ ID NO:4.

25 105. An isolated Hu-Asp2(a) polypeptide comprising an amino acid sequence at least 95% identical to a sequence comprising the amino acid sequence of SEQ ID NO:8.

30 106. An isolated antibody that binds specifically to the Hu-Asp polypeptide of any of claims 104-107.
sequence comprising the amino acid sequence of SEQ ID NO:8.

35 107. An isolated antibody that binds specifically to the Hu-Asp polypeptide of any of claims 104-107.

40 108. * A method to identify a cell that can be used to screen for inhibitors of β secretase activity comprising:

45 a) identifying a cell that expresses a protease capable of cleaving APP at the β secretase site,
comprising:

- 50 i) collect the cells or the supernatant from the cells to be identified
ii) measure the production of a critical peptide, where the critical
peptide is selected from the group consisting of either the APP C-terminal peptide or soluble APP,
iii) select the cells which produce the critical peptide.

5 109. The method of claim 108 where the cells are collected and the critical peptide is the APP C-terminal peptide created as a result of the β secretase cleavage.

10 110. The method of claim 108 where the supernatant is collected and the critical peptide is soluble APP where the soluble APP has a C-terminal created by β secretase cleavage.

15 111. The method of claim 108 where the cells contain any of the nucleic acids or polypeptides of claims 1-86 and where the cells are shown to cleave the β secretase site of any peptide having the following peptide structure, P2, P1, P1', P2', where P2 is K or N, 10 where P1 is M or L, where P1' is D, where P2' is A.

20 112. The method of claim 111 where P2 is K and P1 is M.

15 113. The method of claim 112 where P2 is N and P1 is L.

25 114. * Any bacterial cell comprising any nucleic acids or peptides in claims 1-86 and 92-107.

30 115. A bacterial cell of claim 114 where the bacteria is *E. coli*.

20 116. Any eukaryotic cell comprising any nucleic acids or polypeptides in claims 1-86 and 92-107.

35 117. * Any insect cell comprising any of the nucleic acids or polypeptides in claims 1-86 and 92-107. 25

40 118. A insect cell of claim 117 where the insect is sf9, or High 5.

45 119. A insect cell of claim 100 where the insect cell is High 5.

30 120. A mammalian cell comprising any of the nucleic acids or polypeptides in claims 1-86 and 92-107.

5 121 A mammalian cell of claim 120 where the mammalian cell is selected from the group consisting of, human, rodent, lagomorph, and primate.

10 122 A mammalian cell of claim 121 where the mammalian cell is selected from the group consisting of human cell.

15 123 A mammalian cell of claim 122 where the human cell is selected from the group comprising HEK293, and IMR-32.

20 124 A mammalian cell of claim 121 where the cell is a primate cell.

25 125 A primate cell of claim 124 where the primate cell is a COS-7 cell.

30 126 A mammalian cell of claim 121 where cell is selected from a rodent cells.

35 127 A rodent cell of claim 126 selected from, CHO-K1, Neuro-2A, 3T3 cells.

40 128 A yeast cell of claim 115.

45 129 An avian cell of claim 115.

50 130. * Any isoform of APP where the last two carboxy terminus amino acids of that isoform are both lysine residues.

55 131 The isoform of APP from claim 130 comprising the isoform known as APP695 modified so that its last two having two lysine residues as its last two carboxy terminus amino acids.

60 132 The isoform of claim 131 comprising SEQ. ID. 16.

65 133 The isoform variant of claim 130 comprising SEQ. ID. NO. 18, and 20.

5 134 Any eukaryotic cell line, comprising nucleic acids or polypeptides of claim 130-133.

10 135 Any cell line of claim 134 that is a mammalian cell line (HEK293, Neuro2a, are preferred plus any others.)

15 136 A method for identifying inhibitors of an enzyme that cleaves the beta secretase cleavage site of APP comprising:

- 10 a) culturing cells in a culture medium under conditions in which the enzyme causes processing of APP and release of amyloid beta-peptide into the medium and causes the accumulation of CTF99 fragments of APP in cell lysates,
- 20 b) exposing the cultured cells to a test compound; and specifically determining whether the test compound inhibits the function of the enzyme by
- 25 measuring the amount of amyloid beta-peptide released into the medium and or the amount of CTF99 fragments of APP in cell lysates;
- 30 c) identifying test compounds diminishing the amount of soluble amyloid beta peptide present in the culture medium and diminution of CTF99 fragments of APP in cell lysates as Asp2 inhibitors.

20 137 The method of claim 136 wherein the cultured cells are a human, rodent or insect cell line.

35 138 The method of claim 137 wherein the human or rodent cell line exhibits β secretase activity in which processing of APP occurs with release of amyloid beta-peptide into the

40 culture medium and accumulation of CTF99 in cell lysates.

45 139. A method as in claim 138 wherein the human or rodent cell line treated with the antisense oligomers directed against the enzyme that exhibits β secretase activity, reduces

50 release of soluble amyloid beta-peptide into the culture medium and accumulation of CTF99 in cell lysates.

5 140. A method for the identification of an agent that decreases the activity of a Hu-Asp polypeptide selected from the group consisting of Hu-Asp1, Hu-Asp2(a), and Hu-Asp2(b), the method comprising

10 (a) determining the activity of said Hu-Asp polypeptide in the presence of a test agent and in the absence of a test agent; and

(b) comparing the activity of said Hu-Asp polypeptide determined in the presence of said test agent to the activity of said Hu-Asp polypeptide determined in the absence of said test agent;

15 whereby a lower level of activity in the presence of said test agent than in the absence of said test agent indicates that said test agent has decreased the activity of said Hu-Asp polypeptide..

20 141. The nucleic acids, peptides, proteins, vectors, cells and cell lines, and assays described herein.

ATGGCCGCACTTGGCCCCGGCGCGCTGCTGTGCTCTCTGCTGGCCAGTGGCTCCTCTCCGCCCC
M G A L A R A L L L P L L A Q W L L R A

CCCCCGAGCTTGGCCCCGGCGCTTTCAGCTTGGCCCTCCCGGTGGCCGGCGCACGAAC
A P E L A P A P F T L P L R V A A A T N

GGCCTACTTTCGCCCCACCCCGGACCCCGGAGCCCTGCGGAGCGCCACGCCAGCGGCTTG
R V V A A T P T P G P G T P A E R H A D G L

GGCGTGGCCCTTGGAGCCTTGGCTTCCCGGGCGCCGCCAACTTCTTGGCCATG
A L A L E P A L A L S P A G A A N F L A M

GTAGACAACCTGCAGGGGAGACTCTGGCCGGGCTACTACCTGGAGATGCTGATCGGGACC
V D N L Q G D S G R G Y Y L E M L I G T

CCCCCGCAGAAGCTACAGATTCTGCTGACACTGGAAGCAGTAACCTTTGCGGTGGCAGGA
P P Q K L Q I L V D T G S S N F A V A G

ACCCCGCACTCCTACATAGACACGTACTTTGACACAGAGAGCTCTAGCACATACCGCTCC
T P H S Y I D T Y F D T E R S S T Y F S

AAGGGCTTTGACGTCACAGTGAAGTACACACAAGGAAGCTGGACGGGCTTCGTTGGGGAA
K G F D V T V K Y T Q G S W T G F V G E

GACCTGCTCACCAATCCCCAAGGGTTCAACTACTTCTTTTCTTGTGCAACATTTGCCACTT
D L V T I P K G G F N T S F L V N I A T I

TTTGAATCAGAGAATTTCTTTTGGCTGGGATTAAATGGGAATGGAATCTTGGCCTAGCT
T F E S E N A T T F F L P G I K W N G I L G L A

TATGCCACACTTGGCAAGCCATCAAGTTCTCTGAGACCTTCTTTCAGATCCCTGGTGACA
Y A T L A K P S S S L E T F F D S L V T

CAAGCAAAACATCCCCAACGTTTTCCTCATGTCAGATGTGTGGAGCGGGCTTCCCGCTTGGT
Q A N I P N V F S M Q M C G A G L P V A

GGATCTGGGACCAACCGAGGTAGTCTTGTCTTGGGTGGAATTGAACCAAGTTTGTATAAA
G S G T N G G S L V L G G I E P S L Y K

GGAGACATCTGGTATACCCCTATTAAAGGAAGTGGTACTACCGATGAGAATTTGTAA
G D I W Y T P I K E E W Y T Y Q I E I L K

TTGGAAATTTGGAGGCCAAAGCCCTTAATCTGGACTGACAGAGCTATAACGCAGACAAGGCC
L E T G G G Q S L N L D C R E Y N A D K A

ATCTGTGACAGTGGCACCACGCTGCTGCGCCTGCCCCAGAAGGTGTTTGATGCGGTGGT
I V D S G T T L L R L P Q K V F D A V V

GAACTGTGCCCCGCGCATCTCTGATTCCAGAATTTCTCTGATGGTTTTCGGACTGGGTCC
E A V A R A S L I P E F S D G F W T G S

CAGCTGGGTGCTGGACGAATTCGGAACACCTTGGTCTTTACTTCCCTAAATCTCCATC
Q L A C W T N S E T P W S Y F P K I S I

TACTGTAGAGATGAGAACTCCAGCAGGTCACTCCGTATCACAACTCTGCTTCAGCTTTAC
Y L R D E N S S R S F R I T I L P Q L Y

ATTACGCCCATGATGGGGCGGCGCTGAATTTGAATGTTACCGAATTTGGCATTTTCCCA
I P M M G S A G L N Y E C Y R F G I S P

TCCACAAATGCGCTGGTGTGCTGGCCACGCTGATGAGAGGGCTTCTACGTCATCTTCGAC
S T N A L V I G A T V M E G F Y V I F D

AGAGCCCGAGAAGAGGGTGGGCTTTCGAGCGAGGCCCTGTGCAGAAATTCAGGTTGCTGAC

FIGURE 1 (2)

R A Q K R V G F A A S P C A E I A G A A
GTCTCTGAAATTTCCGGGCTTTCTCAACAGAGGATGTAGCCAGCAACTGTGTCCCGCT
V S E I S G P F S T E D V A S N C V P A
CAGTCCTTGAGCGAGCCCATTTTGTGGATTGTGTCTATGCCCTCATGAGCGTCTGTGGA
Q S L S E P I L W I V S Y A L M S V C G
GCCATCCTCCTTGTCTTAATCGTCCGTGCTGCTGCGGTTCGGGTGTGAGCGTCCGCCC
A I L L V L I V L L L L P F R C Q R R P
CGTGACCCCTGAGGTCTCAATGATGAGTCTCTCTGGTCAGACATCGCTGGAATGAATA
R D P E V V N D E S S L V R H R W K
GCCAGGCTGACCTCAAGCAACCATGAATCAGCTATTAAAGAAATCACATTTCAGGGC
AGCAGCCGGGATCGATGGTGGCGCTTCTCCTGTGCCACCCGCTTCAATCTCTGTCT
GCTCCAGATGCCCTTCTAGATTCACTGTCTTTGATTCTTGATTTCAGCTTCAAAATC
CTCCCTACTTCCAAGAAAAATAATTAACAAAAAATTCATTCTAAACCAAAAAA
AAAA

FIGURE 2 (1)

ATGGCCCAAGCCCTGCCCTGGCTCCTGCTGTGGATGGGCGGGGAGTGCTGCCTGCCAC
M A Q A L P W L L L W M G A G V L P A H
GGCACCAGCACGGCATCCGGCTGCCCCCTGCGCAGCGGCCCTGGGGGGCGCCCCCTGGGG
G T Q H G I R L P L R S G L G G A P L G
CTGCGGCTGCCCGGGAGACCGACGAAGAGCCCGAGGAGCCCGGCCGGAGGGGCAGCTTT
L R L P R E T D E E P E E P G R R G S F
GTGGAGATGGTGGACAACCTGAGGGGCAAGTCGGGGCAGGGCTACTACGTGGAGATGACC
V E M V D N L R G K S G Q G Y Y V E M T
GTGGGCAGCCCCCGCAGACGCTCAACATCCTGGTGGATACAGGCAGCAGTAACCTTGCA
V G S P P Q T L N I L V D T G S S N F A
GTGGGTGCTGCCCCCACCCTTCCTGCATCGCTACTACCAGAGGCAGCTGTCCAGACA
V G A A P H P F L H R Y Y Q R Q L S S T
TACCGGGACCTCCGGAAGGGTGTGTATGTGCCCTACACCCAGGGCAAGTGGGAAGGGGAG
Y R D L R K G V Y V P Y T Q G K W E G E
CTGGGCACCGACCTGGTAAGCATCCCCCATGGCCCCAACGTCAGTGTGCGTGCCAACATT
L G T D L V S I P H G P N V T V R A N I
GCTGCCATCACTGAATCAGACAAGTTCTTCATCAACGGCTCCAAGTGGGAAGGCATCCTG
A A I T E S D K F F I N G S N W E G I L
GGGCTGGCCTATGCTGAGATTGCCAGGCTTTGTGGTGTGCTGGCTTCCCCCTCAACAGTCT
G L A Y A E I A R L C G A G F P L N Q S
GAAGTGCTGGCCTCTGTGCGAGGGAGCATGATCATTGGAGGTATCGACCACTCGCTGTAC
E V L A S V G G S M I I G G I D H S L Y
ACAGGCAGTCTCTGGTATACACCCATCCGGCGGGAGTGGTATTATGAGGTGATCATTGTG
T G S L W Y T P I R R E W Y Y E V I I V
CGGGTGAGATCAATGGACAGGATCTGAAAATGGACTGCAAGGAGTACAACCTATGACAAG
R V E I N G Q D L K M D C K E Y N Y D K
AGCATTGTGGACAGTGGCACCACCAACCTTCGTTTGCCCAAGAAAGTGTTTGAAGCTGCA
S I V D S G T T N L R L P K K V F E A A
GTCAAATCCATCAAGGCAGCCTCCTCCACGGAGAAGTTCCCTGATGGTTTCTGGCTAGGA
V K S I K A A S S T E K F P D G F W L G
GAGCAGCTGGTGTGCTGGCAAGCAGGCACCCCTTGGAACATTTTCCCAGTCACTCA
E Q I V C W Q A G T T P W N I F P V I S
CTCTACCTAATGGGTGAGGTTACCAACAGTCCTTCCGCATCACCATCCTTCCGCAGCAA
L Y L M G E V T N Q S F R I T I L P Q Q
TACCTGCGGCCAGTGGGAAGATGTGGCCACGTCCCAAGACGACTGTTACAAGTTTGCCATC

FIGURE 2 (2)

Y L R P V E D V A T S Q D D C Y K F A I
TCACAGTTCATCCACGGGCACTGTTATGGGAGCTGTTATCATGGAGGGCTTCTACGTTGTC
S Q S S T G T V M G A V I M E G F Y V V
TTTGATCGGGCCCCGAAAACGAATTGGCTTTGCTGTCAGCGCTTGCCATGTGCACGATGAG
F D R A R K R I G F A V S A C H V H D E
TTCAGGACGGCAGCGGTGGAAGGCCCTTTTGTACCTTGGACATGGAAAGCTGTGGCTAC
F R T A A V E G P F V T L D M E D C G Y
AACATTCCACAGACAGATGAGTCAACCCTCATGACCATAGCCTATGTCATGGCTGCCATC
N I P Q T D E S T L M T I A Y V M A A I
TGCGCCCTCTTCATGCTGCCACTCTGCCTCATGGTGTGTCAGTGGCGCTGCCTCCGCTGC
C A L F M L P L C L M V C Q W R C L R C
CTGCGCCAGCAGCATGATGACTTTGCTGATGACATCTCCCTGCTGAAGTGAGGAGGCCCA
L R Q Q H D D F A D D I S L L K
TGGGCAGAAGATAGAGATTCCCCTGGACCACACCTCCGTGGTTCACTTTGGTCACAAGTA
GGAGACACAGATGGCACCTGTGGCCAGAGCACCTCAGGACCCTCCCCACCCACCAATGC
CTCTGCCCTTGATGGAGAAGGAAAAGGCTGGCAAGGTGGGTCCAGGGACTGTACCTGTAG
GAAACAGAAAAGAGAAGAAAAGACACTCTGCTGGCGGGAATACTCTTGGTCACCTCAAA
TTTAAGTCGGGAAATTCTGCTGCTTGAACTTCAGCCCTGAACCTTTGTCCACCATTCCT
TTAAATTCTCCAACCCAAAGTATTCTTCTTTCTTAGTTTCAGAAGTACTGGCATCACAC
GCAGGTTACCTTGGCGTGTGTCCTGTGGTACCCTGGCAGAGAAGAGACCAAGCTTGTTT
CCCTGCTGGCCAAAGTCAGTAGGAGAGGATGCACAGTTTGCTATTTGCTTTAGAGACAGG
GACTGTATAACAAGCCTAACATTGGTGCAAAGATTGCCTCTTGAAAAAAAAAAAAA

FIGURE 3 (1)

ATGGCCCAAGCCCTGCCCTGGCTCCTGCTGTGGATGGGCGCGGGAGTGCTGCCTGCCAC
M A Q A L P W L L L W M G A G V L P A H
GGCACCAGCACGGCATCCGGCTGCCCTGCGCAGCGGCTGGGGGGCGCCCCCTGGGG
G T Q H G I R L P L R S G L G G A P L G
CTGCGGCTGCCCCGGGAGACCGACGAAGAGCCCGAGGAGCCCGGCGGAGGGGCAGCTTT
L R L P R E T D E E P E E P G R R G S F
GTGGAGATGCTCCACAACCTGAGGGGCAAGTCGGGGCAGGGCTACTACGTGGAGATGACC
V E M V D N L R G K S G Q G Y Y V E M T
GTGGGCAGCCCCCGCAGACGCTCAACATCCTGGTGGATACAGGCAGCAGTAACCTTGGCA
V G S P P Q T L N I L V D T G S S N F A
GTGGGTGCTGCCCCCACCCCTTCCTGCATCGCTACTACCAGAGGCAGCTGTCCAGCACA
V G A A P H P F L H R Y Y Q R Q L S S T
TACCGGGACCTCCGGAAGGGTGTGTATGTGCCCTACACCCAGGGCAAGTGGGAAGGGGAG
Y R D L R K G V Y V P Y T Q G K W E G E
CTGGGCACCGACCTGGTAAGCATCCCCCATGGCCCCAACGTCACTGTGCGTGCCAAACATT
L G T D L V S I P H G P N V T V R A N I
GCTGCCATCACTGAATCAGACAAGTTCTTCATCAACGGCTCCAAGTGGGAAGGCATCCTG
A A I T E S D K F F I N G S N W E G I L
GGGCTGGCCTATGCTGAGATTGCCAGGCCTGACGACTCCCTGGAGCCTTTCTTTGACTCT
G L A Y A E I A R P D D S L E P F F D S
CTGGTAAAGCAGACCCACGTTCCCAACCTCTTCTCCCTGCAGCTTTGTGGTGCTGGCTTC
L V K Q T H V F N L F S L Q L C G A G F
CCCCTCAACAGTCTGAAGTGCTGGCCTCTGTGGGAGGGAGCATGATCATTGGAGGTATC
P L N Q S E V L A S V G G S M I I G G I
GACCACTCGCTGTACACAGGCAGTCTCTGGTATACACCCATCCGGCGGGAGTGGTATTAT
D H S L Y T G S L W Y T P I R R E W Y Y
GAGGTCAATCATTGTGCGGGTGGAGATCAATGGACAGGATCTGAAAATGGACTGCAAGGAG
E V I I V R V E I N G Q D L K M D C K E
TACAACTATGACAAGAGCATTTGTGGACAGTGGCACCACCAACCTTCGTTTGGCCCAAGAAA
Y N Y D K S I V D S G T T N L R L P K K
GTGTTTGAAGCTGCAGTCAAAATCCATCAAGGCAGCCTCTCCACGGAGAAGTTCCCTGAT
V F E A A V K S I K A A S S T E K F P D

FIGURE 3 (2)

GGTTTCTGGCTAGGAGAGCAGCTGGTGTGCTGGCAAGCAGGCACCACCCCTTGGAACATT
G F W L G E Q L V C W Q A G T T P W N I
TTCCCAGTCATCTCACTCTACCTAATGGGTGAGGTTACCAACCAGTCCTTCCGCATCACC
F F V I S L Y L M G E V T N Q S F R I T
ATCCTTCCGCAGCAATACCTGCGGCCAGTGAAGATGTGGCCACGTCCTCAAGACGACTGT
I L P Q Q Y L R P V E D V A T S Q D D C
TACAAGTTTGCCATCTCACAGTCATCCACGGGCACTGTTATGGGAGCTGTTATCATGGAG
Y K F A I S Q S S T S T V M G A V I M E
GGCTTCTACGTTGTCTTTGATCGGGCCCGAAAACGAATTGGCTTTGCTGTCAGCGCTTGC
G F Y V V F D R A R K R I G F A V S A C
CATGTGCACGATGAGTTTCAGGACGGCAGCGGTGAAGGCCCTTTGTACCTTGGACATG
H V H D E F R T A A V E G P F V T L D M
GAAGACTGTGGCTACAACATTCCACAGACAGATGAGTCAACCCTCATGACCATAGCCTAT
E D C G Y N I P Q T D E S T L M T I A Y
GTCATGGCTGCCATCTGCGCCCTCTTCATGCTGCCACTCTGCCTCATGGTGTGTCAGTGG
V M A A I C A L F M L P L C L M V C Q W
CGCTGCCTCCGCTGCCTGCGCCAGCAGCATGATGACTTTGCTGATGACATCTCCCTGCTG
R C L R C L R Q Q H D D F A D D I S L L
AAGTGAGGAGGCCCATGGGCAGAAGATAGAGATTCCCCTGGACCACACCTCCGTGGTTCA
K
CTTTGGTCACAAGTAGGAGACACAGATGGCACCTGTGGCCAGAGCACCTCAGGACCCCTCC
CCACCCACCAAATGCCTCTGCCTTGATGGAGAAGGAAAAGGCTGGCAAGGTGGGTTCAG
GGACTGTACCTGTAGGAAACAGAAAAGAGAAGAAAGAAGCACTCTGCTGGCGGGAATACT
CTTGGTCACCTCAAATTTAAGTCGGGAAATTCTGCTGCTTGAAACTTCAGCCCTGAACCT
TTGTCCACCATTCCTTTAAATTCTCCAACCCAAAGTATTCTTCTTTTCTTAGTTTCAGAA
GTACTGGCATCACACGCAGGTTACCTTGGCGTGTGTCCCTGTGGTACCCTGGCAGAGAAG
AGACCAAGCTTGTTCCTGCTGGCCAAAGTCAGTAGGAGAGGATGCACAGTTTGCTATT
TGCTTTAGAGACAGGGACTGTATAACAAGCCTAACATTGGTGCAAAGATTGCCTCTTGA
ATTAAAAAAAAAAAAAAAAAAAAAAAAA

FIGURE 4

ATGSCCCAGCGCTGCACITGGCTCCTGCTATGGGTGGGCTCGGGAATGCTGCCCTGCCAG
 M A F A L H W L L L W V G S G M L P A Q
 GGAACCCATCTCGGCATCCGGCTGCCCTTCGCAGCGGCTGGCAGGGCCACCCCTGGGC
 G T H L G I R L P L R S G L A G P P L G
 CTGAGGCTGCCCCGGGAGACTGACGAGGAATCGGAGGAGCCTGGCCGGAGAGGCAGCTTT
 L R L P R E T D E E S E E P G R R G S F
 GTGAGATGCTGGACACCTCAGGGGAAAGTCCGCCCCAGGCTACTATGCTGGAGATGACC
 V E M V D N L R G K S G Q G Y Y V E M T
 GTAGGCAGCCCCCAGAGCGCTCAACATCCTGCTGGACACGGGCAGTAGTAACCTTGCA
 V G S P P Q T L N I L V D T G S S N F A
 GTGGGGCTGCCCCACACCCCTTCTGCTGCTACTACAGAGGCAGCTGTCCAGCACA
 V G A A P H P F L H R Y Y Q R C L S S T
 TATGAGACCTCCGAAAGGGTGTGTATGTGCCCTACACCCAGGGCAAGTGGGAGGGGAA
 Y R D L R K G V Y V P Y T Q G K W E G E
 CTGGCAGCGACCTGCTGAGCATCCCTCATGGCCCCAACGTCAGTGTGGCTGCCAACATT
 L G T D L V S I P H G P N V T V R A N I
 GGTGCCATCACTGAATCGGACAAAGTCTCTCATCAATGCTTCCAACTGGGAGGGCATCTTA
 A A I T E S D K F F I N G S N W E G I L
 GGGCTGGCCTATGCTGAGATTCGACGCCCGACGACTCTTTGGAGCCCTTCTTTGACTCC
 G L A Y A E I A R P D D S L E P F F D S
 CTGCTGAGCAGACCCACATTCCCAACATCTTTTCCCTGCAGCTCTGTGGCGCTGGCTTC
 L V K Q T H I P N I F S L Q L C G A G F
 CCGCTCAACCAGACCGAGGCACTGGCCTCGGTGGGAGGGAGCATGATCATTTGGTGGTATC
 P L N Q T E A L A S V G G S M I I G G I
 GACCACTGCTATACACGGGAGTCTCTGTACACACCCATCGGCGGAGTGGTATTAT
 D K S L Y T G S L W Y T P I R R E W Y Y
 GAACTGATCATTTGATCGTGGAAATCAATGGTCAAGATCTCAAGATGGACTGCAAGGAG
 E V I I V R V E I N G Q D L K M D C K E
 TACAACACGACAAGAGCATTGTGGACAGTGGGACCAACCTTGGCTTGGCCAGAAA
 Y N Y D K S I V D S G T T N L R L P K K
 GTATTTGAAGCTGCCGTCAAGTCCATCAAGGACGCTCTCGACGAGAGAGTCCCGGAT
 V F E A A V K S I K A A S S T E K F P D
 GGCTTTTGGCTAGGGGAGCAGCTGGTGTCTGCTGCAAGCAGGACGACCCCTTGGAACTT
 G F W L G E Q L V C W Q A G T T P W N I
 TTCCAGTCACTTTCACCTTACCTCATGGGTGAAGTCACCAATCAGTCTCTCCGCATCACC
 F P V I S L Y L M G E V T N Q S F R I T
 ATCCTTCTCAGCAATACCTACGCGCGGTGGAGGACGTGGCCACCTCCCAAGACGACTGT
 I L P Q Q Y L R P V E D V A T S Q D D C
 TACAAGTTTCGCTGTCTCAGAGTCATCCACGGGCACTGTTATGGGAGCCGTCATCATGGAA
 Y K F A V S Q S S T G T V M G A V I M E
 GGTTTCTATGTCTGCTTCGATCGAGCCCGAAAGCGAATTGGCTTTGCTGTACGCGCTTGC
 G F Y V V F D R A R K R I G F A V S A C
 CATGTGCACGATGAGTTCAGGACGGCGGAGTGGAAAGGTCCGTTTGTACGGCAGACATG
 H V H D E F R T A A V E G P F V T A D M
 QAAGACTGTGGCTACAACATTCCCCAGACAGATGAGTCAACACTTATGACCATAGCCCTAT
 E D C G Y N I P Q T D E S T L M T I A Y
 GTCATGGCGGCCATCTGCCGCCCTCTTCATGTTGCCACTCTGCCTCATGCTATGTCAGTGG
 V M A A I C A L F M L P L C L M V C Q W
 CGCTGCGCTGCGTTGCTGCGCCACGACGATGACTTTGCTGATGACATCTCCCTGCIC
 F C L R C L F H Q H D D F A D D I S L L
 AAGTAAGGAGGCTCGTGGGCAGATGATGGAGACGCCCCCTGGACCAATCTGGGTGGTTCC
 F
 CTTTGGTTCACATGAGTTGGAGCTATGGATGGTACTGTGGCCAGAGCACCTCAGGACCCCT
 CACCAACCTGCCAATGCTTCTGGCGTGACAGAACAGAGAAATAGGCAAGCTGATATACA
 GGGCTTGCACCTGTAGGACACAGGAGAGGGAAGGAAGCACCCCTTCTGGTGGCAGGAATAT
 CCTTAGGCACCAAACTTGAAGTTGGAATTTTGTCTGCTTGAAGCTTACGCCCTGACCCCT
 CTGCCCCAGCATCCTTTAGAGTCTCCAACTAAAGTATTCTTTATGTCCTTCCAGAAGTAC
 TGGCGTCATACTCAGGCTACCCGCGATGTGCTCCCTGTGCTACCCCTGGCAGAGAAAGGCC
 AATCTCATTTCCCTGCTGGCCAAAGTCAGCAGAAGAAGGTGAAGTTTGGCAGTTGCTTTAG
 TGAATAGGAGTGCAGACTCAGCCTACACTGGTACAAAGACTGCGTCTTGAGATAAACA
 GAA

FIGURE 5

```
1 MAQALFWLLLWMGAGVLPAGHTQHGIPLRSLGGLPLGLRLPRETDEE 50
  |||||
1 MAPALHWLLLWVGSGMLPAQGTHLGIPLRSLGAGPPLGLRLPRETDEE 50
  |||||

51 PEEPGRRGSFVEMVDNLRGKSGQGYVEMTVGSPQTLNILVDTGSSNFA 100
  |||||
51 SEEPGRRGSFVEMVDNLRGKSGQGYVEMTVGSPQTLNILVDTGSSNFA 100
  |||||

101 VGAAPHPFLHRYYQRLSSTYRDLRKGVPYPTQGWEGELGTDLVSIPI 150
  |||||
101 VGAAPHPFLHRYYQRLSSTYRDLRKGVPYPTQGWEGELGTDLVSIPI 150
  |||||

151 GPNVTVRANIAAITESDKFFINGSNWEGILGLAYAEIARPDDSLEPFDFS 200
  |||||
151 GPNVTVRANIAAITESDKFFINGSNWEGILGLAYAEIARPDDSLEPFDFS 200
  |||||

201 LVKQTHVPNLFSLQLCGAGFPLNQSEVLASVGGSMIIGGIDHSlyTGSW 250
  |||||
201 LVKQTHIPNIFSLQLCGAGFPLNQTEALASVGGSMIIGGIDHSlyTGSW 250
  |||||

251 YTPIRREWYYEVIIIVRVEINGQDLKMDCKEYNYDKSIVDSGTTNLRPKK 300
  |||||
251 YTPIRREWYYEVIIIVRVEINGQDLKMDCKEYNYDKSIVDSGTTNLRPKK 300
  |||||

301 VFEEAVKSIKAASSTEKFPDGFVLGEQLVCWQAGTTPWNIFPVISLYLMG 350
  |||||
301 VFEEAVKSIKAASSTEKFPDGFVLGEQLVCWQAGTTPWNIFPVISLYLMG 350
  |||||

351 EVTNQSFRTILPQQYLRPVEDVATSQDDCYKFAISQSSTGTVMGAVIME 400
  |||||
351 EVTNQSFRTILPQQYLRPVEDVATSQDDCYKFAVSQSSTGTVMGAVIME 400
  |||||

401 GFYVVFDRARKRIGFAVSACHVHDEFRTAAVEGPFVTLMEDCGYNIPQT 450
  |||||
401 GFYVVFDRARKRIGFAVSACHVHDEFRTAAVEGPFVTADMEDCGYNIPQT 450
  |||||

451 DESTLMTIAYVMAAICALFMLPLCLMVCQWRCLRCLRHQHDDFADDISLL 500
  |||||
451 DESTLMTIAYVMAAICALFMLPLCLMVCQWRCLRCLRHQHDDFADDISLL 500
  |||||

501 K 501
  |
501 K 501
```

FIGURE 6 (1)

ATGGCTAGCATCACTGGTGGACAGCAAATGGGTGCGGGATCCACCCAGCACGGCATCCGG
M A S M T G G Q Q M G R G S T Q H G I R

CTGCCCCCTGCGCAGCGCCCTGGGSGCGCCCCCTGGGGCTGCGGCTGCCCCGGAGACG
L P L R S G L G S A P L G L R L P R E T

GACGAAGAGCCCCGAGGAGCCCGCGGAGGGGCGAGCTTTGTGGAGATGGTGGACAACCTG
D E E P E E P G R R G S F V E M V D N L

AGGGGCAAGTCGGGGCAGGGCTACTACGTGGAGATGACCTGGGCGAGCCCCCGCAGAGG
R G K S G Q G Y Y V E M T V G S P P Q T

CTCAACATCCTGGTGGATACAGGCAGCAGTAACCTTTCAGTGGGTGCTGCCCCCACCCG
L N I L V D T G S S N F A V G A A P H P

TTCTGTCATCGCTACTACAGAGGCGAGCTGTCCAGCACATACCGGACCTCCGGAAGGGG
T L H R Y Y Q R Q L S S T Y R D L R K G

GTGTATGTGCCCTACACCCAGGCAAGTGGGAAGGGGAGCTGGGCAACCGACCTGGTAAGC
V Y V P Y T Q G K W E G E L S T D L V S

ATCCCCCATGGCCCCAACGTCACCTGTCGTGCCAACATTGCTGCCATCACTGAATCAGAC
I P H G P N V T V R A N I A A I T E S D

AAGTTCTTCAACCGCTCCAACCTGGGAAGGCATCCCGGGGCTGGCCCTATGCTGAGATT
K F F I N G S N W E G I L G L A Y A E I

GCCAGGCCTGACGACTCCCTGGAGCCCTTTCTTTGACTCTCTGTAAAGCAGACCCACGTT
A R P D C S L E P F F D S L V K Q T H V

CCCAACCTCTTCTCCCTGCAGCTTTGTGGTGTGGCTTCCCCCTCAACCACTCTGAAGTC
P N L F S L Q L C G A G F P L N Q S E V

CTGGCCTCTGTGCGAGGAGCATGATCATGGAGGTATCGACCACTCGCTGTACACAGGC
L A S V G G S M I I G G I D H S L Y T G

AGTCTCTGGTATACACCCATCCGGCGGGAGTGGTATTATGAGGTCACTATTGTGCGGGTG
S L W Y T P I E R E W Y Y E V I I V R V

GAGATCAATGCACAGGATCTGAAATGGACTGCAAGGAGTACAACTATGACAAGAGCATT
E I N G Q D L F M D C K E Y N Y D K S I

GTGGACASTGCCACCACCAACCTTCCTTTCCCAACAACTCTTTGAAGCTGCAGTCAAA
V D S G T T N L R L P K K V F E A A V K

TCCATCAAGGCAGCCTCTCCACGGAGAAGTTCCCTGATGGTTTCTGGCTAGGAGAAGAC
S I K A A S S T E K F P D G F W L G E Q

CTGSGTGTGGCAAGCAGGCACCCCTTGAACATTTTCCAGTCATCTCACTCTAC
L V C W Q A G T T P W N I F P V I S L Y

CTAATGGGTGAGGTTACCAACCAGTCCTTCCGCATCACCATCCTTCCCGAGCAATACCTG
L M G E V T N Q S F R I T I L P Q Q Y L

CGGCCAGTGAAGATGTGCCACGTCCTCCAGACGACTGTTACAAGTTTGCATCTCACAG

FIGURE 6 (2)

R P V E D V A T S Q D D C Y K F A I S Q
TCATCCACGGGCACTGTTATGGGAGCTGTTATCATGGAGGGCTTCTACGTTGTCTTTGAT
S S T G T V M G A V I M E G F Y V V F D
CGGGCCCGAAAACGAATTGGCTTTGCTGTCAGCGCTTGCCATGTGCACGATGAGTTCAGG
R A R K R I G F A V S A C H V H D E F R
ACGGCAGCGGTGGAAGGCCCTTTTGTACCTTGGACATGGAAGACTGTGGCTACAACATT
T A A V E G P F V T L D M E D C G Y N I
CCACAGACAGATGAGTCATGA
P Q T D E S *

FIGURE 7 (1)

ATGGCTAGCATGACTGGTGGACAGCAAAATGGGTGCGGATCGATGACTATCTCTGACTCT
 M A S M T G G Q Q M G R G S M T I S D S
 CCGGCTGAACAGGACGGATCCACCCAGCACGGCATCCGGCTGCCCTGCGCAGCGGCCCTG
 P R E Q D G S T Q H G I R L P L R S G L
 GGGGCGCCCCCTGGGGCTGCCGCTGCCCGGAGACCGACGAAGAGCCCGAGGAGCCC
 G G A P L G L R L P R E T D E E P E E P
 CCCCAGCGCGCAGCTTTGTGGAGATGCTGGACAACTGAGGGGCAAGTCGCGGCAGGGC
 G F R G S F V E M V D N L R G K S J Q G
 TACTACGTGGAGATGACCGTGGCGCAGCCCCCGCAGACCGCTCAACATCCTGCTGGATACA
 Y Y V E M T V G S P P Q T L N I L V D T
 GGCAGCAGTAACCTTTGAGTGGGTGCTSCCCCCACCCCTTCTGTCATCGCTACTACAG
 C S S N F A V G A A P H P F L H R Y Y Q
 AGGCAGCTGTCCAGCACATACCGGGACCTCCGGAAGGGCGTGTATGTGCCCTACACCCAG
 R Q L S S T Y E D L R K G V Y V P Y T Q
 GGCAAGTGGGAAGGGAGCTGGGCACCGACCTGGTAAGCATCCCCATGGCCCCAACGTC
 G H W E G E L G T D L V S I P H G P N V
 ACTGTGCGTGCCAACATTGCTGCCATCACTGAATCAGACAAGTTCTTCATCAACGGCTCC
 T V R A N I A A I T E S D K F F I N C S
 AACTGGGAAGGCATCCTGGGGCTGGCCTATGCTGAGATTGCCAGGCCCTGACGACTCCCTG
 N W E G I L G L A Y A E I A R P D D S L
 GAGCCTTTCTTTGACTCTCTGGTAAAGCAGACCCACGTTCCCAACCTCTTCTCCCTGCAG
 E P F F D S L V K Q T H V P N L F S L Q
 CTTCGTGGTGGCTTCCCCCTCAACCAGTCTGAAGTGTGGCCTCTGTGGAGGGAGC
 L C G A G F P L N Q S E V L A S V G G S
 ATGATCATTTGGAGGTATCGACCACTCGCTGTACACAGGCAGTCTCTGGTATACACCCATC
 M I I G G I D H S L Y T G S L W Y T P I
 CGGCGGGAGTGGTATTATGAGGTCATCATTGTGCGGCTGGAGATCAATGGACAGGATCTG
 R R E W Y Y E V I I V R V E I N G Q D L
 AAAATGGACTGCAAGGAGTACAACATATGACAAGAGCATTTGTGGACAGTGGCACCACCAAC
 K M D C K E Y N Y D K S I V D S G T T N
 CTTCGTTTGGCCCAAGAAAGTCTTTGAAGCTGCAGTCAAATCCATCAAGGCAGCCCTCCTCC
 L E L P K K V F E A A V K S I K A A S S
 ACGGAGAAGTTCCTGTATGGTTTCTGGCTAGGAGAGCAGCTGGTGTGTGGTAAGCAGGC
 T E K F P D G F W L G E Q L V C W Q A G
 ACCACCCCTTGGAAACATTTTCCAGTCATCTCACTCTACSTAATGGGTGAGGTACCAAC
 T T P W N I F P V I S L Y L M G E V T N

FIGURE 7 (2)

CAGTCCTTCGGCATCACCATCCTTCGGCAGCAATACCTGGGGCCAGTGGGAAGATGTGGCC
Q S F R I T I L P Q Q Y L R P V E D V A
ACGTCCCAAGACGACTGTACAAAGTTTGGCATCTCACAGTCATCCACGGGCACTGTATG
T S Q D D C Y K F A I S Q S S I G T V M
GGAGCTGTTATCATGGAGGGCTTCTACGTTGTCTTTGATCGGGCCGAAAACGAATTGGC
G A V I M E G F Y V V F D R A R K R I G
TTTCCTGTCAGCGCTTGGCATGTGCACGATGAGTTCAGGACGGCAGCGGTGGAAGGCCCT
F A V S A C H V H D E F R T A A V E G P
TTTGTACATTGGACATGGAAGACTGTGGCTACAACATTCCACAGACAGATGAGTCATGA
F V T L D M E D C G Y N I P Q T D E S *

FIGURE 8 (1)

ATGACTCAGCATGCTATTCGTCTGCTCACTGCGTAGCGGTCTGGGTGGTGTCCACTGGGT
 M T Q H G I R L F L R S G L G G A P L G -
 CTGGCTCTGCCCCGGGAGACCGACCAAGAGCCCCGAGGAGCCCCGGGAGGGGCGAGCTTT
 L R L P R E T D E E P E E P G R R G S F -
 CTGGAGATGGTGACAACCTGAGGGGCAAGTCGGGGCAGGGCTACTACGTGGAGATGACC
 V E M V D N L R G K S G Q G Y Y V E M T -
 GTGGGCAGCCCCCGCAGACGCTCAACATCCTGGTGATACAGGCAACAGTAACCTTTGCA
 V G S P P Q T L N I L V D T G S S N F A -
 GTGGGTGCTGCCCCCACCCTTCTGATCGCTACTACAGAGGCGAGCTGTCCAGGACA
 V G A A P H P F L H R Y Y Q R Q L S S T -
 TACCGGACCTTCGGGAAGGGCTGTATGTGCTTACATCCAGGGCAAGTGGGAAGGGGAG
 Y R D L R K G V Y V P Y T Q G K W E G E -
 CTGGCCACCGACTTGTGAAGCATGCCCCATGCCCCCAAGCTCACTGTGCGTGGCAACATT
 L G T D L V S I P H G P N V T V R A N I -
 GCTGCCATCACTGAATCAGACAACTTCTTCATCAACGGCTCCCACTGGGAAGGCATCCTG
 A A I T E S D K F F I N G S N W E G I L -
 GGGCTGGCCTATGCTGAGATTGGCCAGGCTGACGACTCCCTGGAGCCTTTCTTTGACTCT
 G L A Y A E I A R P D D S L E P F F D S
 CTGATAAAGCAGACCCACCTTCCCAACCTCTTCTCCCTGCACTTTCTGCTGCTGGCTTC
 L V K Q T H V P N L F S L Q L C G A G F -
 CCCCCTCAACCACTGTGAAGTGTGCGCTCTGTGCGGAGGAGCATGATCATTGGAGGTATC
 P L N Q S E V L A S V G G S M I I G G I -
 GACCACTCGGTGTACACAGGCACTCTCTGTATACACCATCCGGCGGGAGTGGTATTAT
 D H S L Y T G S L W Y T P I R R E W Y Y -
 GAGGTCACTATTGTGCGGGTGGAGATCAATGACAGGATCTGAAAATGGACTGCAAGGAG
 E V I I V R V E I N G Q D L K M D C K E
 TACAACTATGACAAGAGCATTGTGACAGTGGCACCCACCAACCTTCGTTTGGCCCAAGAAA
 Y N Y D K S I V D S G T T N L R L P K K -
 GTGTTTGAAGCTGCAGTCAAAATCCATCAAGCAGCCTCCCTCCACGGAGAAGTTCCTGTAT
 V F E A A V K S I K A A S S T E K F P D -
 GGTTTCTGGCTAGGAGAGCAGCTGGTGTGCTGGCAAGCAGGCAACCCCTTGGAAACATT
 G F W L G E Q L V C W Q A S T T P W N I -
 TTCCAGTCATCTCACTCTACCTAATGGGTGAGGTTACCAACCAGTCCTTTTGGCATCACC
 F P V I S L Y L M G E V T N Q S F R I T -
 ATCCTTCCGCAGCAATACCTGCGGTCAGTGGGAAGATGTGSCCAGTCCCAAGACGACTGT
 I L P Q Q Y L R P V E D V A T S Q D D C -

FIGURE 8 (2)

TACAAGTTTGCCATCTCACAGTCATCCACGGGCACTGTTATGGGAGCTGTTATCATGGAG
Y K F A I S Q S S T G T V M G A V I M E -
GGCTTCTACGTTGTCCTTGATCGGGCCCGAAAACGAATTGGCTTTGCTGTCAGCGCTTGC
G F Y V V F D R A R K R I G F A V S A C -
CATTAG
H

FIGURE 9




FIGURE 10




FIGURE 11

MAOALPWLLLLWMGAGVLPAGHTQHGIRLPLRSGLGGAFLGLRLPRETDEE
PEEPGRRGSFVEMVDNLRGKSGQGYVEMTVGSPQTINILVDTGSSNFA
VGAAPHFPLHRYYQRQLSSTYRDLRKGVYVPYTQGWEGELGTDLVSIPIH
GPNVTVRANIAAITESDKFFINGSNWEGILGLAYAEIARPDSDLPPFFDS
LVKQTHVPNLFSLQLCGAGFPLNQSEVLASVGGSMIIGGIDHSLYTGLW
YTPIRREWYVEVILVRVEINGQDLKMDCKEYNDKSIIVDSGTTNLRPLPK
VFEAAVKSIIKAASSTEKFPDGFVLGEQLVCWQAGTTPWNIFPVISLYLMG
EVTNQSFRTITLPQQYLRPVEDVATSQDDCYKFAISQSSTGTVMGAVIME
GFYVVFDRARKKRIGFAVSACHVHDEFRTAAVEGPFVTLDMEDCGYNIPQT
DES

FIGURE 12

MAQALPWLLWMCAGVLPAGHTQHGIRLPLRSGLGAPLGLRLPRETDEE
PEEPGRRGSFVEMVDNLRGKSGQGYVEMTVGSPPQTLNVLVDTGSSNFA
VGAAPHFPLHRYYQRQLSSTYRDLRKGVYVPYTQGWEGELGTDLVSIPIH
GPNVTVRANIAAITESDKFFINGSNWEGILGLAYAEIARPDDSLEPFFDS
LVKQTHVPNLFSLQLCGAGFPLNQSEVLASVGGSMIIGGIDHSLYTGSLW
YTPIRREWYEVIIIVRVEINGQDLKMDCKEYNYDKSIVDSGTTNLRPLPKK
VFEAAVKSIIKAASSTEKFPDGFWLGEQLVCWQAGTTPWNIFPVISLYLMG
EVTNQSFRTILPQQYLRPVEDVATSQDDCYKFAISQSSTGTVMGAVIME
GFYVVVDRARKRIGFAVSACHVHDEFRTAAVEGPFVTLDMEDCGYNIPQT
DESHHHHHH

SEQUENCE LISTING

<110> Gurney, Mark E.

Bienkowski, Michael J.

Heinrikson, Robert L.

Parodi, Luis A.

Yan, Riqiang

Pharmacia & Upjohn Company

<120> Alzheimer's Disease Secretase

<130> 6177.P CP

<140>

<141>

<150> 60/101,594

<151> 1998-09-24

<160> 49

<170> PatentIn Ver. 2.0

<210> 1

<211> 1804

<212> DNA

<213> Homo sapiens

<400> 1

atggggcgac tggcccgggc gctgctgctg cctctgctgg cccagtggat cctgcgcgcc 60

ggcccgaggc tggcccccgc gcccttcagc ctgccctccc ggggtggcgc ggccacgaac 120

cgcgtagttg cgcacacccc gggacccggg acccctgcgg agcgccacgc cgaagggttg 180
 gcgcctgcgc tggagccctgc cctggcgtec ccggcgggcg ccgcccaactt cttggccatg 240
 gtagacaaac tgcaggggga ctctggcggc ggctactacc tggagatgct gatcggggac 300
 ccccgacaga agctacagat tctggttgac actggaagca gtaactttgc cgtggcagga 360
 acccgcactt cctacataga cactactttt gacacagaga ggtctagcan ataccgctcc 420
 aagggttttg acgtacacgt gaagtacaca caagggaagct ggaagggttt cgttggggaa 480
 gacctcgtca ccatcccaaa agccttcaat acttcttttc ttgtcaacat tgcactatt 540
 ttgaatcag agaatttttt tttgcctggg attaaatgga atggaatact tggcctagct 600
 tatgccacac ttgccaaagg atcaagttct ctggagacct tcttcgactc cttggtgaca 660
 gaagcaaaac tcccaacgt tttctccatg cagatgtgtg gaggcgctt gcccgttgct 720
 gcatctggga ccaacggagg tagtcttgc ttgggtggaa ttgaaccaag ttgtataaa 780
 cgagacatct ggtatacccc tattaaggaa gagtggtaact accagataga aattctgaaa 840
 ttggaatttg gaggccaaag ccttaactct gactgcagag agtataacgc agacaaggcc 900
 atcttggaac gtggcaccac gctgctgcgc ctgccccaga aggtgtttga tgcggtggtg 960
 gaagctgttg cccgcgcac tctgattcca gaattctctg atggtttctg gactgggttc 1020
 cagctggcgt gctggacgaa ttcggaacaa ccttggctct acttccctaa aatctccac 1080
 tacctgagag atgagaactc cagcaggtca ttcgtatca caactctgac taagctttac 1140
 attcagccca tgatgggggc cggcctgaat tatgaatgtt accgattcgg catttccca 1200
 tccacaaatg cgtggtgat cgggtgccac gtgatggagg gcttctacgt catcttcgac 1260
 agagcccaga agaggggtgg ctctgcagcg agccctgtg cagaaattgc aggtgctgca 1320
 gtgtctgaaa ttctcgggac ttctcacaac gaggatgtag ccagcaactg tgcctccgct 1380
 cagtctttga gcgagcccat tttgtggatt gtgtctatg cgtctatgag cgtctgtgga 1440
 gccatctctc ttgtcttaat cgtctgtctg ctgctgcctt tccggtgtca gctctgcctc 1500
 ggtgacctg aggtctgcaa tgatgagtc tctctggta gacatcgtg gaaatgaata 1560
 gccagggctg acctcaagca accatgaact cagctattaa gaaaatcaca ttccagggc 1620
 agcagccggg atcgatggtg gcctttctc ctgtgccac cgtcttcaa tctctgtct 1680
 gctccagat gcctcttaca ttaactgtct ttgattctt gatttcaag ctttcaaat 1740
 ctccctactt ccaagaaaaa taattaaaaa aaaaacttca ttctaaacca aaaaaaaaaa 1800
 aaaa

1804

<210> 2

<211> 518

<212> PRT

<213> Homo sapiens

<400> 2

Met Gly Ala Leu Ala Arg Ala Leu Leu Leu Pro Leu Leu Ala Gln Trp

1 5 10 15

Leu Leu Arg Ala Ala Pro Gln Leu Ala Pro Ala Pro Phe Thr Leu Pro

20 25 30

Leu Arg Val Ala Ala Ala Thr Asn Arg Val Val Ala Pro Thr Pro Gly

35 40 45

Pro Gly Thr Pro Ala Glu Arg His Ala Asp Gly Leu Ala Leu Ala Leu

50 55 60

Glu Pro Ala Leu Ala Ser Pro Ala Gly Ala Ala Asn Phe Leu Ala Met

65 70 75 80

Val Asp Asn Leu Gln Gly Asp Ser Gly Arg Gly Tyr Tyr Leu Glu Met

85 90 95

Leu Ile Gly Thr Pro Pro Gln Lys Leu Gln Ile Leu Val Asp Thr Gly

100 105 110

Ser Ser Asn Phe Ala Val Ala Gly Thr Pro His Ser Tyr Ile Asp Thr

115 120 125

Tyr Phe Asp Thr Glu Arg Ser Ser Thr Tyr Arg Ser Lys Gly Phe Asp

130 135 140

Val Thr Val Lys Tyr Thr Gln Gly Ser Trp Thr Gly Phe Val Gly Glu

145 150 155 160

Asp Leu Val Thr Ile Pro Lys Gly Phe Asn Thr Ser Phe Leu Val Asn

165 170 175

Ile Ala Thr Ile Phe Glu Ser Glu Asn Phe Phe Leu Pro Gly Ile Lys

180 185 190

Trp Asn Gly Ile Leu Gly Leu Ala Tyr Ala Thr Leu Ala Lys Pro Ser

195 200 205

Ser Ser Leu Glu Thr Phe Phe Asp Ser Leu Val Thr Gln Ala Asn Ile

210 215 220

Pro Asn Val Phe Ser Met Gln Met Cys Gly Ala Gly Leu Pro Val Ala

225 230 235 240

Gly Ser Gly Thr Asn Gly Gly Ser Leu Val Leu Gly Gly Ile Glu Pro

245 250 255

Ser Leu Tyr Lys Gly Asp Ile Trp Tyr Thr Pro Ile Lys Glu Glu Trp

260 265 270

Tyr Tyr Gln Ile Glu Ile Leu Lys Leu Glu Ile Gly Gly Gln Ser Leu

275 280 285

Asn Leu Asp Cys Arg Glu Tyr Asn Ala Asp Lys Ala Ile Val Asp Ser

290 295 300

Gly Thr Thr Leu Leu Arg Leu Pro Gln Lys Val Phe Asp Ala Val Val
 305 310 315 320
 Glu Ala Val Ala Arg Ala Ser Leu Ile Pro Glu Phe Ser Asp Gly Phe
 325 330 335
 Trp Thr Gly Ser Gln Leu Ala Cys Trp Thr Asn Ser Glu Thr Pro Trp
 340 345 350
 Ser Tyr Phe Pro Lys Ile Ser Ile Tyr Leu Arg Asp Glu Asn Ser Ser
 355 360 365
 Arg Ser Phe Arg Ile Thr Ile Leu Pro Gln Leu Tyr Ile Gln Pro Met
 370 375 380
 Met Gly Ala Gly Leu Asn Tyr Glu Cys Tyr Arg Phe Gly Ile Ser Pro
 385 390 395 400
 Ser Thr Asn Ala Leu Val Ile Gly Ala Thr Val Met Glu Gly Phe Tyr
 405 410 415
 Val Ile Phe Asp Arg Ala Gln Lys Arg Val Gly Phe Ala Ala Ser Pro
 420 425 430
 Cys Ala Glu Ile Ala Gly Ala Ala Val Ser Glu Ile Ser Gly Pro Phe
 435 440 445
 Ser Thr Glu Asp Val Ala Ser Asn Cys Val Pro Ala Gln Ser Leu Ser
 450 455 460
 Glu Pro Ile Leu Trp Ile Val Ser Tyr Ala Leu Met Ser Val Cys Gly

465 470 475 480

Ala Ile Leu Leu Val Leu Ile Val Leu Leu Leu Leu Pro Phe Arg Cys

 485 490 495

Gln Arg Arg Pro Arg Asp Pro Glu Val Val Asn Asp Glu Ser Ser Leu

 500 505 510

Val Arg His Arg Trp Lys

 515

<210> 3

<211> 2070

<212> DNA

<213> Homo sapiens

<400> 3

atggcccaag cctgccctg gctcctgctg tggatggggc eggagtgct gctgcaccac 60

ggcaccacag acggcatcog gctgccctg cgcagcggcc tgggggggc cccctgggg 120

tggggctgc cccgggagac cgaagagag cccgaggagc cggccggag gggcagcttt 180

gtggagatgg tggacaacct gaggggcaag tggggcagg gctactaagt ggagatgacc 240

gtgggcagcc cccgcagac gctcaacatc ctgctgata cagccacag taacttttca 300

gtgggtgctg ccccccacc ctctctgcat cgtactaac agaggcagct gtccagcaca 360

taccgggacc tccggaaggg tgtgtatgt cctacacc agggcaagt ggaaggggag 420

gtggccacag acctggttag cctcccccac ggcccacag tcactgtgg tgccaacatt 480

gctgcacaca ctgaatcaga caagttcttc atcaacggct ccaactggga aggcacccct 540

gggtggcct atgtgagat tgcaggcct gacgactcc tggagccttc ctttgactc 600

ctggtaaagg agaccacgt tcccaacctc ttctccctgc acctttgtgg tctggcttc 660

ccctcaacc agtctgaagt gctggcctc tggggaggga gcatgacat tggaggtatc 720

gaccactcgc tgtacacagg cagtctctgg talacacca tccggcggga gtggtattat 780

gaggtcatca ttgtgggggt ggagatcaat ggacaggate tgaacatgga ctgcaaggag 840
 tacaactatg acaagagcat tgtggacagt ggcaccacca accttcqtti gcccaadaaa 900
 gtgtttgaag ctgcagtcaa atccatcaag gcagctctct ccacggagaa gtteectgat 960
 ggtttctggc taggagagca gctggtgtgc tggcaagcag gcaaccaccc ttggaacatt 1020
 ttccagtea tctcactcta cctaattgggt gaggttaaca accagctctt cccatcacc 1080
 atccttcgcg agcaatacct gcggccagtg gaagatgtgg caagtcaca agacgactgt 1140
 tacaagtttg ccatctcaca gtcatccacg ggcactqta tggagctctt tatcatqga 1200
 ggctttctag ttgttttga tggggccga aaacgaattg gctttgtgt cagcgttgc 1260
 catgtgcag atgagttcag gacggcagcg gtggaaggcc cttttgtcac ctggacatg 1320
 gaagactgtg gctacaacat tccacagaca gatgagtcaa ccttcatgac catagcctat 1380
 gtcatggctg ccatctgcgc cctcttcctg ctgccactct gctcatgggt gtgtcagtgg 1440
 cgtctctcc gctgcctgcg ccagcagcat gatgactttg ctgatgacat ctccctgctg 1500
 agtgaggag gcccatgggc agaagataga gattcccttg gaccacactt ccgtgggtta 1560
 ctttggtaac aagtaggaga cacagatggc acctgtggcc agagcacctc aggacctcc 1620
 ccaccacca atgctctctg ccttgatgga gaaggaaaag gctggcaagg tgggttcag 1680
 ggaactgtac tgtaggaaac agaaaagaga agaaagaagc actctgctgg cgggaatact 1740
 ttggttacc tcaattttaa gtgggaaat tctgtgtgtt gaaacttcag cctgaacct 1800
 ttgtccacca tctctttaa ttctccaacc caaagtattt tcttttctt agtttcagaa 1860
 gtactggcat cacacgcagg ttactttggc gtgtgtctct gtggtacctt ggcagagaag 1920
 agaccaagct tgtttctctg ctggccaaag tcagtaggag aggatgcaca gtttctatt 1980
 tgccttagag acagggactg tataaacaag cctaacattg gtgcaagat tgcctcttga 2040
 attaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2070

<210> 4

<211> 501

<212> PRT

<213> Homo sapiens

<400> 4

Met Ala Gln Ala Leu Pro Trp Leu Leu Leu Trp Met Gly Ala Gly Val

1

5

10

15

Leu Pro Ala His Gly Thr Gln His Gly Ile Arg Leu Pro Leu Arg Ser

20

25

30

Gly Leu Gly Gly Ala Pro Leu Gly Leu Arg Leu Pro Arg Glu Thr Asp

35

40

45

Glu Glu Pro Glu Glu Pro Gly Arg Arg Gly Ser Phe Val Glu Met Val

50

55

60

Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val Glu Met Thr

65

70

75

80

Val Gly Ser Pro Pro Gln Thr Leu Asn Ile Leu Val Asp Thr Gly Ser

85

90

95

Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe Leu His Arg Tyr

100

105

110

Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg Lys Gly Val

115

120

125

Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu Leu Gly Thr Asp

130

135

140

Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val Arg Ala Asn Ile

145

150

155

160

Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly Ser Asn Trp

165

170

175

Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile Ala Arg Pro Asp Asp

180

185

190

Ser Leu Glu Pro Phe Phe Asp Ser Leu Val Lys Gln Thr His Val Pro

195

200

205

Asn Leu Phe Ser Leu His Leu Cys Gly Ala Gly Phe Pro Leu Asn Gln

210

215

220

Ser Glu Val Leu Ala Ser Val Gly Gly Ser Met Ile Ile Gly Gly Ile

225

230

235

240

Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr Pro Ile Arg Arg

245

250

255

Glu Trp Tyr Tyr Glu Val Ile Ile Val Arg Val Glu Ile Asn Gly Gln

260

265

270

Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp Lys Ser Ile Val

275

280

285

Asp Ser Gly Thr Thr Asn Leu Arg Leu Pro Lys Lys Val Phe Glu Ala

290

295

300

Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu Lys Phe Pro Asp

305

310

315

320

Gly Phe Trp Leu Gly Glu Gln Leu Val Cys Trp Gln Ala Gly Thr Thr

325

330

335

Pro Trp Asn Ile Phe Pro Val Ile Ser Leu Tyr Leu Met Gly Glu Val

```

      340              345              350
      "
      "
      Thr Asn Gln Ser Phe Arg Ile Thr Ile Leu Pro Gln Gln Tyr Leu Arg
      355              360              365
      "
      "
      Pro Val Glu Asp Val Ala Thr Ser Gln Asp Asp Cys Tyr Lys Phe Ala
      370              375              380
      "
      "
      Ile Ser Gln Ser Ser Thr Gly Thr Val Met Gly Ala Val Ile Met Glu
      385              390              395              400
      "
      "
      Gly Phe Tyr Val Val Phe Asp Arg Ala Arg Lys Arg Ile Gly Phe Ala
      405              410              415
      "
      "
      Val Ser Ala Cys His Val His Asp Glu Phe Arg Thr Ala Ala Val Glu
      420              425              430
      "
      "
      Gly Pro Phe Val Thr Leu Asp Met Glu Asp Cys Gly Tyr Asn Ile Pro
      435              440              445
      "
      "
      Gln Thr Asp Glu Ser Thr Leu Met Thr Ile Ala Tyr Val Met Ala Ala
      450              455              460
      "
      "
      Ile Cys Ala Leu Phe Met Leu Pro Leu Cys Leu Met Val Cys Gln Trp
      465              470              475              480
      "
      "
      Arg Cys Leu Arg Cys Leu Arg Gln Gln His Asp Asp Phe Ala Asp Asp
      485              490              495
      "
      "
      Ile Ser Leu Leu Lys
      500
  
```

<210> 5

<211> 1977

<212> DNA

<213> Homo sapiens

<400> 5

```

atggcccaag cctgcccctg gctcccctg tggatgggcg cgggagtgtt gctgcccac 60
ggcaccacag acggcaccct gctgcccctg cgcagcggcc tggggggcgc cccctgggg 120
ctggggctgc cccgggagac cgcagcagag cccgaggagc ccggccggag gggcagcttt 180
gtggagatgg tggacaacct gaggggcaag tggggcagg gctactactt ggagatgac 240
gtgggcagcc ccccgagac gctcaacatc ctggtggata caggcagcag taactttgca 300
gtgggtgctg ccccccaccc ctctctgat cgtactacc agaggcagct gtccagcaca 360
taccgggacc tccggaaggc tgtgtatgtg cctacaccc agggcaagtg ggaaggggag 420
gtgggcaccc acctggtaag cctcccccct ggccccaccg tcatgtgtcg tgccaacatt 480
gctgccatca ctcaatcaga caactctctt atcaacggtt ccaactggga aggcacctg 540
gggtgggctt atgtgagat tgccaggctt tgtgtgtg gcttccccct caaccagtct 600
gaagtgtctg cctctgtcg agggagcatg atcattggag gtatcgacca ctgctgtac 660
acaggcagtc tctgtatcc acccaccctg cgggagtgtt attatgaggt gaccattgtg 720
gggtgggaga tcaatggaca gcatctgaaa atggactgca aggagtacaa ctatgacaag 780
agcattgttg acagtggcac caccacccct cgtttgccc agaaagtgtt tgaagctgca 840
gtcaaatcca tcaaggcagc ctctccacag gagaagttcc ctgaaggctt ctggctagga 900
gagcagctgg tctgtggca agcaggcacc accccttggg acattttccc agtcattcca 960
ctctacctaa tgggtgaggt taccacaccg tcttccgca taccatctt tccgcagcaa 1020
tactcgggct cagtggaga tgtggcacc tcccaagacg actgttaccg gtttgcacc 1080
tcacagtcac ccacgggac tgttatggga cctgttatca tggagggtt ctacgttgc 1140
tttgatcggg cccgaaaacg aattggcttt gctgtacgcg ctgcccattt gcagatgag 1200
tccaggacgg cagcggtgga agccctttt gtcaccttg acatggaaga ctgtggtac 1260
aacattccac agacagatga gtcaaccctc atgaccatag cctatgtcat ggctgccac 1320
tgcgccctct tcatgtgac actctgcctc atggtgtgtc agtggcgtg cctccgtgc 1380

```

ctggccagc agcatgatga ctttctgat gacatctccc tgetgaagtg aggaggccca 1440
 tgggcagaag atagagatgc ccttggacca cactcctgtg gttcactttg gtcacaagta 1500
 ggagacacag atggcacctg tggccagagc acctcaggac cctccccacc caccaaatgc 1560
 ctctgcttgc atggagaagg aaaaggctgg caaggctggg tccaggggact gtacctgtag 1620
 gaaacagaaa agagaagaaa gaagcacctc gctggcggga atactcttgg tcacctcaaa 1680
 ttttaagtcg gaaattctgc tgcctgaaa ttcagccctg aacctttgtc caccattcct 1740
 ttaaattctc caaccacaag tatctctctt ttcttagttt cagaagtact ggcattcac 1800
 gcaggttacc ttggcgtgtg tccctgttgt acctggcag agaagagacc aagcttggtt 1860
 ccttgcctgg caaagtcagt aggagaggtt gcacagtctg ctatttgctt tagagacagg 1920
 gactgtataa acagcctaa cattgggtga aagattgctt ctgcaaaaa aaaaaa 1977

<210> 6

<211> 476

<212> PRT

<213> Homo sapiens

<400> 6

Met Ala Gln Ala Leu Pro Trp Leu Leu Leu Trp Met Gly Ala Gly Val

1 5 10 15

Leu Pro Ala His Gly Thr Gln His Gly Ile Arg Leu Pro Leu Arg Ser

20 25 30

Gly Leu Gly Gly Ala Pro Leu Gly Leu Arg Leu Pro Arg Glu Thr Asp

35 40 45

Glu Glu Pro Glu Glu Pro Gly Arg Arg Gly Ser Phe Val Glu Met Val

50 55 60

Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val Glu Met Thr

65 70 75 80

Val Gly Ser Pro Pro Gln Thr Leu Asn Ile Leu Val Asp Thr Gly Ser

85

90

95

Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe Leu His Arg Tyr

100

105

110

Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg Lys Gly Val

115

120

125

Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu Leu Gly Thr Asp

130

135

140

Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val Arg Ala Asn Ile

145

150

155

160

Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly Ser Asn Trp

165

170

175

Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile Ala Arg Leu Cys Gly

180

185

190

Ala Gly Phe Pro Leu Asn Gln Ser Glu Val Leu Ala Ser Val Gly Gly

195

200

205

Ser Met Ile Ile Gly Gly Ile Asp His Ser Leu Tyr Thr Gly Ser Leu

210

215

220

Trp Tyr Thr Pro Ile Arg Arg Glu Trp Tyr Tyr Glu Val Ile Ile Val

225

230

235

240

Arg Val Glu Ile Asn Gly Gln Asp Leu Lys Met Asp Cys Lys Glu Tyr

245 250 255

Asn Tyr Asp Lys Ser Ile Val Asp Ser Gly Thr Thr Asn Leu Arg Leu

260 265 270

Pro Lys Lys Val Phe Glu Ala Ala Val Lys Ser Ile Lys Ala Ala Ser

275 280 285

Ser Thr Glu Lys Phe Pro Asp Gly Phe Trp Leu Gly Glu Gln Leu Val

290 295 300

Cys Trp Gln Ala Gly Thr Thr Pro Trp Asn Ile Phe Pro Val Ile Ser

305 310 315 320

Leu Tyr Leu Met Gly Glu Val Thr Asn Gln Ser Phe Arg Ile Thr Ile

325 330 335

Leu Pro Gln Gln Tyr Leu Arg Pro Val Glu Asp Val Ala Thr Ser Gln

340 345 350

Asp Asp Cys Tyr Lys Phe Ala Ile Ser Gln Ser Ser Thr Gly Thr Val

355 360 365

Met Gly Ala Val Ile Met Glu Gly Phe Tyr Val Val Phe Asp Arg Ala

370 375 380

Arg Lys Arg Ile Gly Phe Ala Val Ser Ala Cys His Val His Asp Glu

385 390 395 400

Phe Arg Thr Ala Ala Val Glu Gly Pro Phe Val Thr Leu Asp Met Glu

405 410 415
 Asp Cys Gly Tyr Asn Ile Pro Gln Thr Asp Glu Ser Thr Leu Met Thr
 420 425 430

Ile Ala Tyr Val Met Ala Ala Ile Cys Ala Leu Phe Met Leu Pro Leu
 435 440 445

Cys Leu Met Val Cys Gln Trp Arg Cys Leu Arg Cys Leu Arg Gln Gln
 450 455 460

His Asp Asp Phe Ala Asp Asp Ile Ser Leu Leu Lys
 465 470 475

<210> 7

<211> 2043

<212> DNA

<213> Mus musculus

<400> 7

atggcccag cgtgcactg gctcctgcta tgggtgggt cgggaatgt gctgcccag 60
 ggaaccata tcgccatccg gctgccctt cgcagcgcc tggcagggcc acccctgggc 120
 ctgaggetgc cccgggagac tgacgaggaa tcggaggagc ctgjcggag aggcagcttt 180
 gtggagatgg tggacaacct gaggggaaag tcggccagg gctactatgt ggagatgacc 240
 gtaggcagcc ccccacagac gctcaacatc ctggtggaca cgggcagtag taactttgca 300
 gtgggggctg ccccacacc ttctctgcat cgtactacc agaggeagct gtcagcaca 360
 tatcgagacc tcgaaaaggq tqtqtatqt cctacacc agggcaagt ggagggggaa 420
 ctgggcaccg acctggtgag catcctcat ggccccacg tcaactgtcg tgccaacatt 480
 gctgccatca ctgaatggga caagttcttc atcaatggtt ccaactggga gggcacccta 540
 gggctggct atgtgagat tcccagccc gacgactct tggagcctt ctttgactcc 600

ctggtgaagc agaccacat tcccaacatc tttccctgc agctctgtgg cgtgggttc 660
 cccctcaacc agaccgagga actggctctg gtgggagggg gcctgctcat tgggtgtatc 720
 gaccactcgc tatcacggg cagtctctgg tacacaccca tccggcggga gtggtatcat 780
 gaagtgatea ttgtactgtt ggaatcaat ggtaagatc tcaagatgga ctgcaaggag 840
 tacaactacg acaagagcat tctggacagt ggtaccacca accttcgctt gcccaagaaa 900
 gtatttgaag ctgcctcaa gtccatcaag gcagctctct cgaaggagaa gtcccgcat 960
 ggtttttggc taggggagca gctgggtgtc tggcaagcag gcacgacccc ttggaacatt 1020
 ttccagtcac tttaacttta cctcatgggt gaagtcacca atcagtcctt ccgcatcacc 1080
 atctctcttc agcaatacct acggcgggtg gaggacgttg ccacgtccca agacgactgt 1140
 tacaagttcg ctgtctcaca gtcatccag ggcactgtta tgggagccgt catcatggaa 1200
 agttttctat tctctctcga tggacccga aagcgaattg gctttgctgt cagcgtttgc 1260
 catgtgcacg atgagttcag gacggcgcca ggggaaggtc cgtttgttac ggcagacatg 1320
 aaactctgtg gctacaacat tcccagaca gatgagtcac cacttatgac catagcctat 1380
 gtcatggggg ccctctgcgc cctcttcctg ttgcactct gcctcatggt atgtcagttg 1440
 cgtgtctctg gttgcctgag ccaccagcac gatgactttt ctgatgacat ctccctgctc 1500
 aagtaaggag gctcgtgggc agatgatgga gacgcctct gaccacatct ggggtggttc 1560
 ctttggtcac atgagttgga gctatggatg gtacctgttg ccagagccac tcaggacct 1620
 caccacctg ccaatgcttc tggcgtgaca gaacagagaa atcaggcaag ctggattaca 1680
 gggcttgcat ctgtaggaca caggagaggg aaggagcag cgtttctggt gcaggatat 1740
 ccttaggcac caccaccttg agttggaaat ttgtctgtt gaagcttcag cctgacct 1800
 ctgcccagca tcttttagag tctccaacct aagttattct ctatgtctt ccagaagtac 1860
 tggcgtcata ctcaggctac ccggcatgtg tccctgtggt acctggcag agaaagggcc 1920
 catctcattc cctgtgggc aaagtcagca gaagaagggt aagtttgca gttgctttag 1980
 tgatagggac tgcagactca agctacact ggtacaaaga ctgcgtcttg agataaaca 2040
 gaa 2043

<210> 8

<211> 501

<212> PRT

<213> Mus musculus

<400> 8

Met Ala Pro Ala Leu His Trp Leu Leu Leu Trp Val Gly Ser Gly Met

1 5 10 15

Leu Pro Ala Gln Gly Thr His Leu Gly Ile Arg Leu Pro Leu Arg Ser

20 25 30

Gly Leu Ala Gly Pro Pro Leu Gly Leu Arg Leu Pro Arg Glu Thr Asp

35 40 45

Glu Glu Ser Glu Glu Pro Gly Arg Arg Gly Ser Phe Val Glu Met Val

50 55 60

Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val Glu Met Thr

65 70 75 80

Val Gly Ser Pro Pro Gln Thr Leu Asn Ile Leu Val Asp Thr Gly Ser

85 90 95

Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe Leu His Arg Tyr

100 105 110

Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg Lys Gly Val

115 120 125

Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu Leu Gly Thr Asp

130 135 140

Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val Arg Ala Asn Ile

145 150 155 160

Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly Ser Asn Trp

165 170 175

Glu Gly Ile Leu Gly Leu Ala Tyr Ala Gln Ile Ala Arg Pro Asp Asp

180 185 190

Ser Leu Glu Pro Phe Phe Asp Ser Leu Val Lys Gln Thr His Ile Pro

195 200 205

Asn Ile Phe Ser Leu Gln Leu Cys Gly Ala Gly Phe Pro Leu Asn Gln

210 215 220

Thr Glu Ala Leu Ala Ser Val Gly Gly Ser Met Ile Ile Gly Gly Ile

225 230 235 240

Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr Pro Ile Arg Arg

245 250 255

Glu Trp Tyr Tyr Glu Val Ile Ile Val Arg Val Glu Ile Asn Gly Gln

260 265 270

Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp Lys Ser Ile Val

275 280 285

Asp Ser Gly Thr Thr Asn Leu Arg Leu Pro Lys Lys Val Phe Glu Ala

290 295 300

Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu Lys Phe Pro Asp

305 310 315 320

Gly Phe Trp Leu Gly Glu Gln Leu Val Cys Trp Gln Ala Gly Thr Thr

325	330	335
Pro Trp Asn Ile Phe Pro Val Ile Ser Leu Tyr Leu Met Gly Glu Val		
340	345	350
Thr Asn Gln Ser Phe Arg Ile Thr Ile Leu Pro Gln Gln Tyr Leu Arg		
355	360	365
Pro Val Glu Asp Val Ala Thr Ser Gln Asp Asp Cys Tyr Lys Phe Ala		
370	375	380
Val Ser Gln Ser Ser Thr Gly Thr Val Met Gly Ala Val Ile Met Glu		
385	390	395
Gly Phe Tyr Val Val Phe Asp Arg Ala Arg Lys Arg Ile Gly Phe Ala		
405	410	415
Val Ser Ala Cys His Val His Asp Glu Phe Arg Thr Ala Ala Val Glu		
420	425	430
Gly Pro Phe Val Thr Ala Asp Met Glu Asp Cys Gly Tyr Asn Ile Pro		
435	440	445
Gln Thr Asp Glu Ser Thr Leu Met Thr Ile Ala Tyr Val Met Ala Ala		
450	455	460
Ile Cys Ala Leu Phe Met Leu Pro Leu Cys Leu Met Val Cys Gln Trp		
465	470	475
Arg Cys Leu Arg Cys Leu Arg His Gln His Asp Asp Phe Ala Asp Asp		
485	490	495

Ile Ser Leu Leu Lys

500

<210> 9

<211> 2088

<212> DNA

<213> Homo sapiens

<400> 9

atgctgcccg gtttggcact gctcctgctg gccgcctgga cgctcggggc gctggaggta 60
 gccactgatg gtaatgctgg cctgctggct gaaccccaga tggccatgtr ctgtggcaga 120
 ctgaacatgc acatgaatgt ccagaatggg aagtgggatt cagatccatc aggcacccaa 180
 ccctgcattg ataccaagga aggcacccctg cagtattgac aagaagtcta cctgaactg 240
 cagatcacca atgtggtaga agccacccaa ccagtaccca tccagaactg gtgcaagcgg 300
 gccgcgaagc agtgcgaagc ccctcccccac ttgtgtatgc cctaccgctg cttagtgtgt 360
 cagtttgtlaa gtgatgcctt tctgttctt gacaagtgc aattcttaca ccaggagagg 420
 tgggatgttt gcgaaactca tcttcaactg cacacgctg ccacagagac atgcagttag 480
 aagaglacca acttgcatga ctaaggcatg ttgtgcctt gccgaattga caagttccga 540
 cgggttagagt ttgtgtgttg ccactggct gaagaaagt acaatgtgga ttctgtgat 600
 gccgaggagg atgactcgga tgtctgggtg gccggagcag acacagacta tgcagatggg 660
 aqtqaagaca aagtagtaga agtagcagag gaggaagaag tggctgaggt ggaacaagaa 720
 gaagcagatg atgacgagga cgatgaggat ggtgatgagg tagaggaaga ggcagaggaa 780
 ccctacgaag aagccacaga gagaaccacc agcattgcc aaccaccac caccaccaca 840
 gagtctgtgg aagaggtggt tccagtctct acaacagcag ccagtacccc tgatgccgtt 900
 gacaagtatc tccagacacc tggggatgag aatgaacatg cccatttcca gaaagccaaa 960
 gagaggcttg aggcacagca ccgagagaga atgtcccagg tcatgagaga atgggaagag 1020
 gcagaacgta aagcaaagaa cttgcctaaa gctgataaga aggcagttat ccagcatttc 1080
 caggagaaag tggaatcttt ggaacaggaa gcagccaccc agagacagca gctggttgag 1140
 acacacatgg ccagagtggg agccatgctc aatgaccgcc gccgcctggc cctggagaa 1200

tacatcaacg ctctgcaggc tgttctctct eggcctctgc acgtgttcaa tatgctaaag 1260
 aagtatgtcc ggcgagaaca gaaggacaga cagcacaccc taaagcattt cgagcatgtg 1320
 cgcattgttg atccccagaa agccgcctag atccggctcc aggttatgac acacctcctt 1380
 gtgattctat aggcctatgaa tcagctctct tccctgtctt acaacctgac tgcagtggcc 1440
 uaggagattc aggatgaagt tgatgagctg ctccagaaaag agcaaaacta ttcagatgac 1500
 gtcttggcca acatgattag tgaaccaagg atcagttacg gaaacgatgc tctcatgcca 1560
 tctttgacgg aaacgaaaac caccgtggag ctctctcccg tgaatggaga gtccagcctg 1620
 gacgatcttc agccgtggca ttcttttggg gctgactctg tgcagccaa cacagaaaac 1680
 gaagttagc ctgttgatgc ccgcctgtct gccgaccgag gactgaccac tgcaccaggt 1740
 tctgggttga caaatatcaa gacggaggag atctctgaag tgaagatgga tgcagaatc 1800
 cgacatgast caggatatga agttcatcat caaaaattgg tttcttttgc agaagatgtg 1860
 ggttcaatca zaggtgcaat cattggctc atggggggcg gtgttgtcat agcgacagtg 1920
 atcgctatca ccttgggtgat gctgaagaag aaacagtaca catccattca tcatggtgtg 1980
 ggggaggttg acgcctgtgt caccacagag gaggccacc tgtccaagat gcagcagaac 2040
 ggtacgaaa atccaacct caagttcttt gacgagatgc agaactag 2088

<210> 10

<211> 696

<212> PRT

<213> Homo sapiens

<400> 10

Met Leu Pro Gly Leu Ala Leu Leu Leu Leu Ala Ala Trp Thr Ala Arg

1 5 10 15

Ala Leu Glu Val Pro Thr Asp Gly Asn Ala Gly Leu Leu Ala Glu Pro

20 25 30

Gln Ile Ala Met Phe Cys Gly Arg Leu Asn Met His Met Asn Val Gln

35 40 45

Asn Gly Lys Trp Asp Ser Asp Pro Ser Gly Thr Lys Thr Cys Ile Asp

50

55

60

Thr Lys Glu Gly Ile Leu Gln Tyr Cys Gln Glu Val Tyr Pro Glu Leu

65

70

75

80

Gln Ile Thr Asn Val Val Glu Ala Asn Gln Pro Val Thr Ile Gln Asn

85

90

95

Trp Cys Lys Arg Gly Arg Lys Gln Cys Lys Thr His Pro His Phe Val

100

105

110

Ile Pro Tyr Arg Cys Leu Val Gly Glu Phe Val Ser Asp Ala Leu Leu

115

120

125

Val Pro Asp Lys Cys Lys Phe Leu His Gln Glu Arg Met Asp Val Cys

130

135

140

Glu Thr His Leu His Trp His Thr Val Ala Lys Glu Thr Cys Ser Glu

145

150

155

160

Lys Ser Thr Asn Leu His Asp Tyr Gly Met Leu Leu Pro Cys Gly Ile

165

170

175

Asp Lys Phe Arg Gly Val Glu Phe Val Cys Cys Pro Leu Ala Glu Glu

180

185

190

Ser Asp Asn Val Asp Ser Ala Asp Ala Glu Glu Asp Asp Ser Asp Val

195

200

205

Trp Trp Gly Gly Ala Asp Thr Asp Tyr Ala Asp Gly Ser Glu Asp Lys

210	215	220	
Val Val Glu Val Ala Glu Glu Glu Glu Val Ala Glu Val Glu Glu Glu			
225	230	235	240
Glu Ala Asp Asp Asp Glu Asp Asp Glu Asp Gly Asp Glu Val Glu Glu			
	245	250	255
Glu Ala Glu Glu Pro Tyr Glu Glu Ala Thr Glu Arg Thr Thr Ser Ile			
	260	265	270
Ala Thr Thr Thr Thr Thr Thr Thr Glu Ser Val Glu Glu Val Val Arg			
	275	280	285
Val Pro Thr Thr Ala Ala Ser Thr Pro Asp Ala Val Asp Lys Tyr Leu			
	290	295	300
Glu Thr Pro Gly Asp Glu Asn Glu His Ala His Phe Gln Lys Ala Lys			
305	310	315	320
Glu Arg Leu Glu Ala Lys His Arg Glu Arg Met Ser Gln Val Met Arg			
	325	330	335
Glu Trp Glu Glu Ala Glu Arg Gln Ala Lys Asn Leu Pro Lys Ala Asp			
	340	345	350
Lys Lys Ala Val Ile Gln His Phe Gln Glu Lys Val Glu Ser Leu Glu			
	355	360	365
Gln Glu Ala Ala Asn Glu Arg Gln Gln Leu Val Glu Thr His Met Ala			
	370	375	380

Arg Val Glu Ala Met Leu Asn Asp Arg Arg Arg Leu Ala Leu Glu Asn

385 390 395 400

Tyr Ile Thr Ala Leu Gln Ala Val Pro Pro Arg Pro Arg His Val Phe

405 410 415

Asn Met Leu Lys Lys Tyr Val Arg Ala Glu Gln Lys Asp Arg Gln His

420 425 430

Thr Leu Lys His Phe Glu His Val Arg Met Val Asp Pro Lys Lys Ala

435 440 445

Ala Gln Ile Arg Ser Gln Val Met Thr His Leu Arg Val Ile Tyr Glu

450 455 460

Arg Met Asn Gln Ser Leu Ser Leu Leu Tyr Asn Val Pro Ala Val Ala

465 470 475 480

Glu Glu Ile Gln Asp Glu Val Asp Glu Leu Leu Gln Lys Glu Gln Asn

485 490 495

Tyr Ser Asp Asp Val Leu Ala Asn Met Ile Ser Glu Pro Arg Ile Ser

500 505 510

Tyr Gly Asn Asp Ala Leu Met Pro Ser Leu Thr Glu Thr Lys Thr Thr

515 520 525

Val Glu Leu Leu Pro Val Asn Gly Glu Phe Ser Leu Asp Asp Leu Gln

530 535 540

Pro Trp His Ser Phe Gly Ala Asp Ser Val Pro Ala Asn Thr Glu Asn

545 550 555 560

Glu Val Glu Pro Val Asp Ala Arg Pro Ala Ala Asp Arg Gly Leu Thr

565 570 575

Thr Arg Pro Gly Ser Gly Leu Thr Asn Ile Lys Thr Glu Glu Ile Ser

580 585 590

Glu Val Lys Met Asp Ala Glu Phe Arg His Asp Ser Gly Tyr Glu Val

595 600 605

His His Gln Lys Leu Val Phe Phe Ala Glu Asp Val Gly Ser Asn Lys

610 615 620

Gly Ala Ile Ile Gly Leu Met Val Gly Gly Val Val Ile Ala Thr Val

625 630 635 640

Ile Val Ile Thr Leu Val Met Leu Lys Lys Lys Gln Tyr Thr Ser Ile

645 650 655

His His Gly Val Val Glu Val Asp Ala Ala Val Thr Pro Glu Glu Arg

660 665 670

His Leu Ser Lys Met Gln Gln Asn Gly Tyr Glu Asn Pro Thr Tyr Lys

675 680 685

Phe Phe Glu Gln Met Gln Asn

690 695

<210> 11

<211> 2088

<212> DNA

<213> Homo sapiens

<400> 11

atgctgcccg gtttcgcact gctcctcctg uccgcctgga cggctcgggc gctggaggta 60
 cccactgatg gtaatgctgg cctgctgggt gaaccccaga ttgcacgtt ctgtggcaga 120
 ctgaacatgc acatgaatgt ccagaatggg aagtgggatt cagatccatc agggacccaa 180
 acctgcattg ataccaagga aggcaccttg cagtattgca aagaagtcta cctgaactg 240
 cagatcccca atgtggtaga agccaaacaa ccagtgaaca tccagaactg gtcgaaggcg 300
 ggccgcaage agtgcaagac ccaccccac ttgtgcatc cctaccctg cttagtgggt 360
 gagtttgtaa gtgagcctt tctgttctt gacaagtga aattcttaca ccaggagagg 420
 atggatgttt gggaaacta tcttcaactg cacaccctg ccaagagac atgcagttag 480
 aagagtacca acctgcctga ctacggcatg ttgctgccct gcggaattga caagttccga 540
 ggggttagat ttgtgtgtg cccactggct gaagaaagty acaatgtgga ttctgtgat 600
 gcggaggagg atgaactgga tgtctgggtg ggccggagag acacagacta tgcagatggg 660
 agtgaagaca aatgtatata agtagcagag gaggaagaag tggctgaggt ggaagaagaa 720
 gaagccgatg atgacgagga ccatgaggat ggtgatgagg tagaggaaga ggtcaggaa 780
 cctacgaag aagccacaga gagaaccacc agcattgcca ccaccaccac caccaccaca 840
 gagtctgttg aagaggtggt tccagtctct acaacagcag ccagtacccc tgatgccgtt 900
 gacaagtatc tccagacacc tggggatgag aatgaacatg cccatttcca gaaagccaaa 960
 gagaggcttg aggccaaaga ccgagagaga atgtcccagg tcatagada atqqaagag 1020
 gcagaacgtc aagcaaagaa cttgcctaaa gctgataaga aggcagttat ccagcatttc 1080
 caggagaaag tggaaacttt ggaacaggaa gcagccaacg agagacagca gctggtygag 1140
 acacacatgg ccagagtggg agccatgctc aatgaccgcc gcgcctggc cctgggagaa 1200
 tacatcccg ctctgcaggc tgttctctct cggctctgct acgtgttcaa tatgctaaag 1260
 aagtatgtcc gcgcagaaca gaaggacaga cagcacccc taaagcattt ccagcatgtc 1320
 cgcattggtg atcccaagaa agccgtcag atccggtccc agyttatgac acacctccgt 1380
 ttgatttatg agcccatgaa tcaatctctc tccctctctc acaacgtgca tgcagtggcc 1440
 gaggagatc aggangaagt tgatgagctg ctccagaaag agcaaaacta ttcagatgac 1500

gtcttggcca acatgattag tgaaccaagg atcagttacg gaaacgatgc tctcatgcca 1580
 tctttgaccc aaacgaaaaa caccctcagc ctctcttccc tgaatggaga gtccagcctg 1620
 gacgatctcc agccttgcca ttcttttggg gctgactctg tgcacgccaa cacagaaaaa 1660
 gaagttgagc ctgttgatgc ccgccttgcg gcgcacagag gactgaaccac tgcaccaggt 1740
 tctgggttga caaatatcaa gacggaggag atctctjaag tgaatctgga tgcagaattc 1800
 cgcacatgact caggatatga agttcactat caaaaattgg tctcttttgc agaagatctg 1860
 ggttcaaaac aaqgttcaat catttquact atggtggggc gtgtttgtcat agcagacagt 1920
 atcgtcacta ccttggtgat gctgaagaag naacagtaca catccattca tcatggtgtg 1980
 gtggaggttg accgcctgtt caccacagag gaggcgcacc tgtccaagat gcagcagaac 2040
 ggctacgaaa atccaacctt caagttcttt gaggcagatgc agaactag 2088

<210> 12

<211> 695

<212> PRT

<213> Homo sapiens

<400> 12

Met Leu Pro Gly Leu Ala Leu Leu Leu Ala Ala Trp Thr Ala Arg

1 5 10 15

Ala Leu Glu Val Pro Thr Asp Gly Asn Ala Gly Leu Leu Ala Gln Pro

20 25 30

Gln Ile Ala Met Phe Cys Gly Arg Leu Asn Met His Met Asn Val Gln

35 40 45

Asn Gly Lys Trp Asp Ser Asp Pro Ser Gly Thr Lys Thr Cys Ile Asp

50 55 60

Thr Lys Glu Gly Ile Leu Gln Tyr Cys Gln Glu Val Tyr Pro Glu Leu

65 70 75 80

Gln Ile Thr Asn Val Val Glu Ala Asn Gln Pro Val Thr Ile Gln Asn

85

90

95

Trp Cys Lys Arg Gly Arg Lys Gln Cys Lys Thr His Pro His Phe Val

100

105

110

Ile Pro Tyr Arg Cys Leu Val Gly Glu Phe Val Ser Asp Ala Leu Leu

115

120

125

Val Pro Asp Lys Cys Lys Phe Leu His Gln Glu Arg Met Asp Val Cys

130

135

140

Glu Thr His Leu His Trp His Thr Val Ala Lys Glu Thr Cys Ser Glu

145

150

155

160

Lys Ser Thr Asn Leu His Asp Tyr Gly Met Leu Leu Pro Cys Gly Ile

165

170

175

Asp Lys Phe Arg Gly Val Glu Phe Val Cys Cys Pro Leu Ala Glu Glu

180

185

190

Ser Asp Asn Val Asp Ser Ala Asp Ala Glu Glu Asp Asp Ser Asp Val

195

200

205

Trp Trp Gly Gly Ala Asp Thr Asp Tyr Ala Asp Gly Ser Glu Asp Lys

210

215

220

Val Val Glu Val Ala Glu Glu Glu Glu Val Ala Glu Val Glu Glu Glu

225

230

235

240

Glu Ala Asp Asp Asp Glu Asp Asp Glu Asp Gly Asp Glu Val Glu Glu

245 250 255

Glu Ala Glu Glu Pro Tyr Glu Glu Ala Thr Glu Arg Thr Thr Ser Ile

260 265 270

Ala Thr Thr Thr Thr Thr Thr Thr Glu Ser Val Glu Glu Val Val Arg

275 280 285

Val Pro Thr Thr Ala Ala Ser Thr Pro Asp Ala Val Asp Lys Tyr Leu

290 295 300

Glu Thr Pro Gly Asp Glu Asn Glu His Ala His Phe Gln Lys Ala Lys

305 310 315 320

Glu Arg Leu Glu Ala Lys His Arg Glu Arg Met Ser Gln Val Met Arg

325 330 335

Glu Trp Glu Glu Ala Glu Arg Gln Ala Lys Asn Leu Pro Lys Ala Asp

340 345 350

Lys Lys Ala Val Ile Gln His Phe Gln Glu Lys Val Glu Ser Leu Glu

355 360 365

Gln Glu Ala Ala Asn Glu Arg Gln Gln Leu Val Glu Thr His Met Ala

370 375 380

Arg Val Glu Ala Met Leu Asn Asp Arg Arg Arg Leu Ala Leu Glu Asn

385 390 395 400

Tyr Ile Thr Ala Leu Gln Ala Val Pro Pro Arg Pro Arg His Val Pro

405	410	415	
Asn Met Leu Lys Lys Tyr Val Arg Ala Glu Gln Lys Asp Arg Gln His			
420	425	430	
Thr Leu Lys His Phe Glu His Val Arg Met Val Asp Pro Lys Lys Ala			
435	440	445	
Ala Gln Ile Arg Ser Gln Val Met Thr His Leu Arg Val Ile Tyr Glu			
450	455	460	
Arg Met Asn Gln Ser Leu Ser Leu Leu Tyr Asn Val Pro Ala Val Ala			
465	470	475	480
Glu Gln Ile Gln Asp Glu Val Asp Glu Leu Leu Gln Lys Glu Gln Asn			
485	490	495	
Tyr Ser Asp Asp Val Leu Ala Asn Met Ile Ser Glu Pro Arg Ile Ser			
500	505	510	
Tyr Gly Asn Asp Ala Leu Met Pro Ser Leu Thr Glu Thr Lys Thr Thr			
515	520	525	
Val Glu Leu Leu Pro Val Asn Gly Glu Phe Ser Leu Asp Asp Leu Gln			
530	535	540	
Pro Trp His Ser Phe Gly Ala Asp Ser Val Pro Ala Asn Thr Glu Asn			
545	550	555	560
Glu Val Glu Pro Val Asp Ala Arg Pro Ala Ala Asp Arg Gly Leu Thr			
565	570	575	

Thr Arg Pro Gly Ser Gly Leu Thr Asn Ile Lys Thr Glu Glu Ile Ser

580

585

590

Glu Val Asn Leu Asp Ala Glu Phe Arg His Asp Ser Gly Tyr Glu Val

595

600

605

His His Gln Lys Leu Val Phe Phe Ala Glu Asp Val Gly Ser Asn Lys

610

615

620

Gly Ala Ile Ile Gly Leu Met Val Gly Gly Val Val Ile Ala Thr Val

625

630

635

640

Ile Val Ile Thr Leu Val Met Leu Lys Lys Lys Gln Tyr Thr Ser Ile

645

650

655

His His Gly Val Val Glu Val Asp Ala Ala Val Thr Pro Glu Glu Arg

660

665

670

His Leu Ser Lys Met Gln Gln Asn Gly Tyr Glu Asn Pro Thr Tyr Lys

675

680

685

Phe Phe Glu Gln Met Gln Asn

690

695

<210> 13

<211> 2088

<212> DNA

<213> Homo sapiens

<400> 13

atgctgcccg gtttggcaat gctctgtctg gcgccttggc cggctcgggc gctggaggta 60
 cccactgatg gtaatgctgg cctgctggct gaaccccaga ttgccatgtt ctgtggcaga 120
 ctgaacatgc acatgaatgt ccagaatggg aagtgggall cagatccatc agggaccaaa 180
 acctgcattg ataccaagga aggcacccct cagtattgcc aagaagtcta cctgaactg 240
 cagatccaca atgtggtaga agccaaccaa ccagtgacca tccagaactg gtgcaagcgg 300
 ggccgcaage agtgcaagac ccaccccaa tttgtgattc cctaccgtg cttagtgtgt 360
 gagtctgtaa gtgatgcct tctgttctt gacaagtcca aattcttaca ccaggagagg 420
 atggtgtttt gcgaaactca tcttcactg caccacgtcg ccaagagac atgcagttag 480
 aagagtacaa acttgcattg ctacggcatg ttgctgcctt ggggaattga caagttccga 540
 ggggtagagt ttgtgtgttg cccactggct gaagaagtg acaatgtgga ttctgtgat 600
 ggggaggagg atgactcggg tctctggctg gggggagcag acacagacta tgcagatggg 660
 agtgagaca aagtagtaga agtagcagag gaggaagaag tggctgaggt ggaagaagaa 720
 gaagccgatg atgacgagga cgtataggat ggtgatgagg tagaggaaga ggcagaggaa 780
 cctacgaag aagccacaga gagaaccacc agcattgcca ccaccaccac caccaccaca 840
 gagtctgttg aagaggtggc tcgatttctt acaacagcag ccagtacccc tgatgcggtt 900
 gacaagtalc tcgagacacc tggggtatg aatgaacatg cccatttcca gaaagccaaa 960
 gagaggcttg aggccaaaga ccgagagaga atgtcccagg tcatgagaga atgggaagag 1020
 gcagaacgtc aagcaagaa cttgcctaaa gctgataaga aggcagttat ccagcatttc 1080
 caggagaaaag tggaaatcttt ggaacaggaa gcagccaacg agagacagca gctggtggag 1140
 acacacatgg ccagagtggc agccatgttc aatgacggcc gcgccttggc cctggagaa 1200
 tacatcaccg ctctgcaygc tgttctctct cggcctcgtc acgtgttcaa tatgttaaag 1260
 aagtatgttc gcgcagaaca gaaggacaga cagcacacc taaagcattt cgagcatgtg 1320
 tgcctggttg atcccaagaa agccgtcag atccggtccc aggttatgac acacctccgt 1380
 gtgatttatg agcgcataaa tcagtctctc tccctgtctt acaacgtgcc tgcagtggcc 1440
 gaggagatcc aggatgaagt tgatgagctg ctccagaaag agcaaaacta ttcagatgac 1500
 gtcttggcca acatgattag tgaaccaagg atcagttacg gaaacgatgc tctcatgcca 1560
 tctttgaccg aaacgaaaac caccgtggag ctcttctccg tgaatggaga gttcaycctg 1620
 gacgatctcc agcgtggcca ttcttttggg gctgaactct tgcagccaa cacagaaaac 1680
 gaagttgagc ctgttgatgc ccgccctgt gcgcacgag gactgaccac tgcaccaggt 1740
 tctgggttga caaatatcaa gacggaggag atctctgaag tgaagatgga tgcagatc 1800

```

cgacatgact caggatataga agttcatcat caaaaattgg ttttttttc agaagatgtg 1860
ggttcaaaaca aagggtgcaat cattggactc atgggtggcg gtgttgteat agcgacagtg 1920
atcttcatca ccttggtgat gtgaagaag aaacagtaca catccattca tcatggtgtg 1980
gtggaggttg acgcgctgt caccacagag gagegcacc tgtccaagat gcagcagaac 2040
ggctacgaaa atccaaccta caagttcttt gagcagatgc agaactag 2088

```

<210> 14

<211> 695

<212> PRT

<213> Homo sapiens

<400> 14

Met Leu Pro Gly Leu Ala Leu Leu Leu Ala Ala Trp Thr Ala Arg

1 5 10 15

Ala Leu Glu Val Pro Thr Asp Gly Asn Ala Gly Leu Leu Ala Glu Pro

20 25 30

Gln Ile Ala Met Phe Cys Gly Arg Leu Asn Met His Met Asn Val Gln

35 40 45

Asn Gly Lys Trp Asp Ser Asp Pro Ser Gly Thr Lys Thr Cys Ile Asp

50 55 60

Thr Lys Glu Gly Ile Leu Gln Tyr Cys Gln Glu Val Tyr Pro Glu Leu

65 70 75 80

Gln Ile Thr Asn Val Val Glu Ala Asn Gln Pro Val Thr Ile Gln Asn

85 90 95

Trp Cys Lys Arg Gly Arg Lys Gln Cys Lys Thr His Pro His Phe Val

100	105	110	
Ile Pro Tyr Arg Cys Leu Val Gly Glu Phe Val Ser Asp Ala Leu Leu			
115	120	125	
Val Pro Asp Lys Cys Lys Phe Leu His Gln Glu Arg Met Asp Val Cys			
130	135	140	
Glu Thr His Leu His Trp His Thr Val Ala Lys Glu Thr Cys Ser Glu			
145	150	155	160
Lys Ser Thr Asn Leu His Asp Tyr Gly Met Leu Leu Pro Cys Gly Ile			
165	170	175	
Asp Lys Phe Arg Gly Val Glu Phe Val Cys Cys Pro Leu Ala Glu Glu			
180	185	190	
Ser Asp Asn Val Asp Ser Ala Asp Ala Glu Glu Asp Asp Ser Asp Val			
195	200	205	
Trp Trp Gly Gly Ala Asp Thr Asp Tyr Ala Asp Gly Ser Glu Asp Lys			
210	215	220	
Val Val Glu Val Ala Glu Glu Glu Glu Val Ala Glu Val Glu Glu Glu			
225	230	235	240
Glu Ala Asp Asp Asp Glu Asp Asp Glu Asp Gly Asp Glu Val Glu Glu			
245	250	255	
Glu Ala Glu Glu Pro Tyr Glu Glu Ala Thr Glu Arg Thr Thr Ser Ile			
260	265	270	

Ala Thr Thr Thr Thr Thr Thr Thr Glu Ser Val Glu Glu Val Val Arg

275

280

285

Val Pro Thr Thr Ala Ala Ser Thr Pro Asp Ala Val Asp Lys Tyr Leu

290

295

300

Glu Thr Pro Gly Asp Glu Asn Glu His Ala His Phe Gln Lys Ala Lys

305

310

315

320

Glu Arg Leu Glu Ala Lys His Arg Glu Arg Met Ser Gln Val Met Arg

325

330

335

Glu Trp Glu Glu Ala Glu Arg Gln Ala Lys Asn Leu Pro Lys Ala Asp

340

345

350

Lys Lys Ala Val Ile Gln His Phe Gln Glu Lys Val Glu Ser Leu Glu

355

360

365

Gln Glu Ala Ala Asn Glu Arg Gln Gln Leu Val Glu Thr His Met Ala

370

375

380

Arg Val Glu Ala Met Leu Asn Asp Arg Arg Arg Leu Ala Leu Glu Asn

385

390

395

400

Tyr Ile Thr Ala Leu Gln Ala Val Pro Pro Arg Pro Arg His Val Phe

405

410

415

Asn Met Leu Lys Lys Tyr Val Arg Ala Glu Gln Lys Asp Arg Gln His

420

425

430

Thr Leu Lys His Phe Glu His Val Arg Met Val Asp Pro Lys Lys Ala

435

440

445

Ala Gln Ile Arg Ser Gln Val Met Thr His Leu Arg Val Ile Tyr Glu

450

455

460

Arg Met Asn Gln Ser Leu Ser Leu Leu Tyr Asn Val Pro Ala Val Ala

465

470

475

480

Glu Gln Ile Gln Asp Glu Val Asp Glu Leu Leu Gln Lys Glu Gln Asn

485

490

495

Tyr Ser Asp Asp Val Leu Ala Asn Met Ile Ser Glu Pro Arg Ile Ser

500

505

510

Tyr Gly Asn Asp Ala Leu Met Pro Ser Leu Thr Glu Thr Lys Thr Thr

515

520

525

Val Glu Leu Leu Pro Val Asn Gly Glu Phe Ser Leu Asp Asp Leu Gln

530

535

540

Pro Trp His Ser Phe Gly Ala Asp Ser Val Pro Ala Asn Thr Glu Asn

545

550

555

560

Glu Val Glu Pro Val Asp Ala Arg Pro Ala Ala Asp Arg Gly Leu Thr

565

570

575

Thr Arg Pro Gly Ser Gly Leu Thr Asn Ile Lys Thr Glu Glu Ile Ser

580

585

590

Glu Val Lys Met Asp Ala Glu Phe Arg His Asp Ser Gly Tyr Glu Val

```

      595              600              605
~
~
His His Gln Lys Leu Val Phe Phe Ala Glu Asp Val Gly Ser Asn Lys
      610              615              620
~
~
Gly Ala Ile Ile Gly Leu Met Val Gly Gly Val Val Ile Ala Thr Val
      625              630              635              640
~
~
Ile Phe Ile Thr Leu Val Met Leu Lys Lys Lys Gln Tyr Thr Ser Ile
      645              650              655
~
~
His His Gly Val Val Glu Val Asp Ala Ala Val Thr Pro Glu Glu Arg
      660              665              670
~
~
His Leu Ser Lys Met Gln Gln Asn Gly Tyr Glu Asn Pro Thr Tyr Lys
      675              680              685
~
~
Phe Phe Glu Gln Met Gln Asn
      690              695
~

```

<210> 15

<211> 2094

<212> DNA

<213> Homo sapiens

<400> 15

```

atgctgcccg gtttggcact gctcctgctg gcagcctgga cggctcgggc gctggaggta 60
ccaactgatg gtaatgctgg cctgctggct gaaccccaga ttgccatgtt ctgtggcaga 120
ctgaacatgc acatgaatgt ccagaatggg aactgggatt cagatccatc aqggacccaa 180
acctgcattg ataccaagga aggcacccctg cagtattgcc aagaagtcta cctgaactg 240

```

cagatcaca atgtggtaga agccaacca ccagtgacca tccagaactg gtgcaagcgg 300
 qcccgcaagc agtacaagac ccacccccc tttgtcattc cctaccctc cttagtctgt 360
 gagtttgtaa gtgatgccct tctcgttctt gacaagtga aattcttaca ccaggagagg 420
 atggtatgtt gcgaaactca tcttcaactg cacaccytcg ccaagagagc atgcagtga 480
 aagagtacaa acttgcctga ctacggcctg ttgctgccct gcggaattga caagttccga 540
 ggggttagagt ttgtgtgttg cccactggct gaagaaagtg acaatgtgga ttctgtgat 600
 ccqaaqagg atgactcga tctctctctg qccggaqca acacagacta tgcagatqgg 660
 agtgaagaca aagtagtaga agtagcagag gaggaagaag tggctgaggt ggaagaagaa 720
 gaagccgatg atgacgagga cgtatgagat ggtgatgagg tagaygaaga ggtgagga 780
 cctacgaag aagccacaga gagaaccacc agcattgcca ccacaccac caccaccaca 840
 gagctgttg aagaggtggg tccagtctct acaacagcag ccagtacccc tgatgcctt 900
 acaagtatc tccagacacc tggggatgag aatgaacatg cccatttcca gaaagccaaa 960
 gagaggttg agccaagca ccagagagaga atgtcccagg tcatgagaga atgggaagag 1020
 gcagaacgtc aagcaaagaa cttgcctaaa gctgataaga aggcagttat ccagcatttc 1080
 caggagaaa tggaattctt ggaacaggaa gcagccaacg agagacagca gctggtggag 1140
 acacacatg ccagagtga agccatgctc aatgacggc gccgctggc cctggagaa 1200
 tacatcacg ctctcagc tytctctct cggcctctc acgtgttcaa tatgtctaa 1260
 aagtatgtc gccgagaaca gaaggacaga cagcacacc taaagcattt ccagcatgtg 1320
 cgcattggtg atcccaagaa agccgctcag atccggctcc aggttatgac acacctccgt 1380
 gtgatttatg agcccatgaa tccgtctct cccctgtct acaacgtgc tgcagtggc 1440
 gagagatc aggatgaag tgatgagctg ctccagaaag agcaaaacta ttcagatgac 1500
 gtcttggcca acatgattag tgaaccaagg atcagttacg gaaacgatgc tctcatgcca 1560
 tctttgaccg aaacgaaaac caccgtggag ctctctccg tgaatggaga gttcagcctg 1620
 gacgatctc agccgtggca ttcttttggg gctgactctg tgcagccaa cacagaaaac 1680
 gaagtgtgc ctgttgatgc ccgcccgtg gccgaccgag gactgaccac tgcaccaggt 1740
 ctggtgtga caaatatcaa gacggaggag atctctgaag tgaagatgga tgcagaatc 1800
 cgacatgact caggatatga agttcctcat caaaaattgg tgttctttgc agaagatgtg 1860
 ggttcaaca aaggtgcaat cattggactc atggtggggc gtgttgcct agcgacagt 1920
 atcgtcatca ccttggtgat gctgaagaag aaacagtaca catccattca tcatggtgtg 1980
 atggaggttg acgcgctgt caccocagag gacgcacc cgttcaagat gcagcagaa 2040
 ggtacgaaa atccaaacta caagttcttt gaccagatgc agaacaagaa gtag 2094

<210> 16

<211> 697

<212> PRT

<213> Homo sapiens

<400> 16

Met Leu Pro Gly Leu Ala Leu Leu Leu Leu Ala Ala Trp Thr Ala Arg

1 5 10 15

Ala Leu Glu Val Pro Thr Asp Gly Asn Ala Gly Leu Leu Ala Glu Pro

20 25 30

Gln Ile Ala Met Phe Cys Gly Arg Leu Asn Met His Met Asn Val Gln

35 40 45

Asn Gly Lys Trp Asp Ser Asp Pro Ser Gly Thr Lys Thr Cys Ile Asp

50 55 60

Thr Lys Glu Gly Ile Leu Gln Tyr Cys Gln Glu Val Tyr Pro Glu Leu

65 70 75 80

Gln Ile Thr Asn Val Val Glu Ala Asn Gln Pro Val Thr Ile Gln Asn

85 90 95

Trp Cys Lys Arg Gly Arg Lys Gln Cys Lys Thr His Pro His Phe Val

100 105 110

Ile Pro Tyr Arg Cys Leu Val Gly Glu Phe Val Ser Asp Ala Leu Leu

115 120 125

Val Pro Asp Lys Cys Lys Phe Leu His Gln Glu Arg Met Asp Val Cys

130 135 140

Glu Thr His Leu His Trp His Thr Val Ala Lys Glu Thr Cys Ser Glu

145 150 155 160

Lys Ser Thr Asn Leu His Asp Tyr Gly Met Leu Leu Pro Cys Gly Ile

165 170 175

Asp Lys Phe Arg Gly Val Glu Phe Val Cys Cys Pro Leu Ala Glu Glu

180 185 190

Ser Asp Asn Val Asp Ser Ala Asp Ala Glu Glu Asp Asp Ser Asp Val

195 200 205

Trp Trp Gly Gly Ala Asp Thr Asp Tyr Ala Asp Gly Ser Glu Asp Lys

210 215 220

Val Val Glu Val Ala Glu Glu Glu Glu Val Ala Glu Val Glu Glu Glu

225 230 235 240

Glu Ala Asp Asp Asp Glu Asp Asp Glu Asp Gly Asp Glu Val Glu Glu

245 250 255

Glu Ala Glu Glu Pro Tyr Glu Glu Ala Thr Glu Arg Thr Thr Ser Ile

260 265 270

Ala Thr Thr Thr Thr Thr Thr Thr Glu Ser Val Glu Glu Val Val Arg

275 280 285

Val Pro Thr Thr Ala Ala Ser Thr Pro Asp Ala Val Asp Lys Tyr Leu

290	295	300	
Glu Thr Pro Gly Asp Glu Asn Glu His Ala His Phe Gln Lys Ala Lys			
305	310	315	320
Glu Arg Leu Glu Ala Lys His Arg Glu Arg Met Ser Gln Val Met Arg			
325	330	335	
Glu Trp Glu Glu Ala Glu Arg Gln Ala Lys Asn Leu Pro Lys Ala Asp			
340	345	350	
Lys Lys Ala Val Ile Gln His Phe Gln Glu Lys Val Glu Ser Leu Glu			
355	360	365	
Gln Glu Ala Ala Asn Glu Arg Gln Gln Leu Val Glu Thr His Met Ala			
370	375	380	
Arg Val Glu Ala Met Leu Asn Asp Arg Arg Arg Leu Ala Leu Glu Asn			
385	390	395	400
Tyr Ile Thr Ala Leu Gln Ala Val Pro Pro Arg Pro Arg His Val Phe			
405	410	415	
Asn Met Leu Lys Lys Tyr Val Arg Ala Glu Gln Lys Asp Arg Gln His			
420	425	430	
Thr Leu Lys His Phe Glu His Val Arg Met Val Asp Pro Lys Lys Ala			
435	440	445	
Ala Gln Ile Arg Ser Gln Val Met Thr His Leu Arg Val Ile Tyr Glu			
450	455	460	

Arg Met Asn Gln Ser Leu Ser Leu Leu Tyr Asn Val Pro Ala Val Ala

465 470 475 480

Glu Glu Ile Gln Asp Glu Val Asp Glu Leu Leu Gln Lys Glu Gln Asn

485 490 495

Tyr Ser Asp Asp Val Leu Ala Asn Met Ile Ser Glu Pro Arg Ile Ser

500 505 510

Tyr Gly Asn Asp Ala Leu Met Pro Ser Leu Thr Glu Thr Lys Thr Thr

515 520 525

Val Glu Leu Leu Pro Val Asn Gly Glu Phe Ser Leu Asp Asp Leu Gln

530 535 540

Pro Trp His Ser Phe Gly Ala Asp Ser Val Pro Ala Asn Thr Glu Asn

545 550 555 560

Glu Val Glu Pro Val Asp Ala Arg Pro Ala Ala Asp Arg Gly Leu Thr

565 570 575

Thr Arg Pro Gly Ser Gly Leu Thr Asn Ile Lys Thr Glu Glu Ile Ser

580 585 590

Glu Val Lys Met Asp Ala Glu Phe Arg His Asp Ser Gly Tyr Glu Val

595 600 605

His His Gln Lys Leu Val Phe Phe Ala Glu Asp Val Gly Ser Asn Lys

610 615 620

Gly Ala Ile Ile Gly Leu Met Val Gly Gly Val Val Ile Ala Thr Val

625 630 635 640

Ile Val Ile Thr Leu Val Met Leu Lys Lys Lys Gln Tyr Thr Ser Ile

645 650 655

His His Gly Val Val Glu Val Asp Ala Ala Val Thr Pro Glu Glu Arg

660 665 670

His Leu Ser Lys Met Gln Gln Asn Gly Tyr Glu Asn Pro Thr Tyr Lys

675 680 685

Phe Phe Glu Gln Met Gln Asn Lys Lys

690 695

<210> 17

<211> 2094

<212> DNA

<213> Homo sapiens

<400> 17

atgctgcccg gtttggcaat gctctgtgtg gccgcctgga cggctcgggc gctggaggta 60
 cccactgatg gtaatgctgg cctgtgggt gaaccccaga ttgccatgtt ctgtggcaga 120
 ctgaacatgc acatgaatgt ccagaatggg aagtgggatt cagatccatc agggacaaaa 180
 acctgcattg ataccaagga aggcatactg cagtattgcc aagaagtcta cctgaactg 240
 cagatcacca atgtggtaga agccaaacaa ccagtgaaca tccagaactg gtgcaagcgg 300
 ggccgcaagc agtjcaagac ccattcccac tttgtgattc cctaccgctg cttagttggt 360
 jagtttgtaa gtgatgcctt tctgttctct gacaagtgea aattcttaca ccaggagagg 420
 atggatgttt gcgaaaacta tcttcaactg cacacgtctg ccaaagagac atgcagttag 480
 aagagtacca accttgcattg ctacggcatg ttgctgcctt gcggaattga caagttccga 540

ggggtagagt ttgtgtgttg cccactggct gaagaaagtg acaatgtgga ttctgctgat 600
 qccgaggagg atgactcgga tgtctggtgg ggcggagcag acacagacta tgcagatggg 660
 agtgaagana aagtagtaga agtagcagag gaggaagaag tggctgaggt ggaagaagaa 720
 gaagccgatg atgacgagga ccatyaggat ggtgatgagg tayaggaaga ggtgaggyaa 780
 ccttacgaag aagccacaga gagaaccacc agcattgcca ccaccaccac caccaccaca 840
 gagtctgttg aagaggtggt tcgagtctct acaacagcag ccagtacccc tgatgcctgt 900
 qacaagtatc tcgagacacc tggggatgag aatgaacatg cccatttcca qaaaaccaaa 960
 gagaggtctg aggcacaagca ccgagagaga atgtcccagg tcatgagaga atgggaagag 1020
 gcagaacgtc aagcaagaa cttgctataa gctgataaga aggcagttat ccagcatttc 1080
 caggagaaag tggaatcttt ggaacaggaa gcagccaacy agagacagca gctgjtggag 1140
 acacacatgg ccagagtggg agccatgctc aatgaccgcc gcgcctgggc cctggagaac 1200
 cacatcccg ctctgcaggg tcttctctct cggctctgtc acgtgttcaa tatgctaaag 1260
 aagtatgtcc gccagaaaca gaaggacaga cagcacacc ctaaagcattt ccagcatgtg 1320
 cgcattggtg atcccaagaa agccgctcag atccggtccc aggttatgac acacctccgt 1380
 gtgatttatg agcgcattga tcagtctctc tccctgctct acaacgtgcc tgcagtggcc 1440
 qaggagatcc aggatgaagt tgatgagctg ctccagaaaag agcaaaaata ttcagatgac 1500
 gtcttggcca acatgallag tgaaccaagg atcagttacg gaaacgatgc tctcatycca 1560
 cctttgaccg aaacgaaaac caccgtggag ctccctcccg tgaatggaga gttcagcctg 1620
 gacgatctcc agccgtggca ttcttttggg gctgactctg tgccagccaa cacagaaaac 1680
 gaagtggagc ctgttgatgc cccccctgct gccgaccgag gactgaccac tcgaccaggt 1740
 tctgggttga caaatatcaa gacggaggag atctctgaag tgaatctgga tgcagaatto 1800
 cgacatgact caggatatga agttcatcat caaaaattgg tgttctttgc agaagatgtg 1860
 qgttcaaaaca aaggtgcaat cattggactc atgggtggcg gtgttgtcat agcgacagtg 1920
 ategtcacca ccttggtgat gctgaagaag aaacagtaca catccatca tcatgggtgtg 1980
 ggggaggttg acgcgctgt cccccagag gagcgccacc tgtccaagat gcagcagaaac 2040
 agctacgaaa atccaaccha caagttcttt gagcagatgc agaacaagaa gtat 2094

<210> 18

<211> 697

<212> PRT

<213> Homo sapiens

<400> 18

Met Leu Pro Gly Leu Ala Leu Leu Leu Leu Ala Ala Trp Thr Ala Arg

1 5 10 15

Ala Leu Glu Val Pro Thr Asp Gly Asn Ala Gly Leu Leu Ala Glu Pro

20 25 30

Gln Ile Ala Met Phe Cys Gly Arg Leu Asn Met His Met Asn Val Gln

35 40 45

Asn Gly Lys Trp Asp Ser Asp Pro Ser Gly Thr Lys Thr Cys Ile Asp

50 55 60

Thr Lys Glu Gly Ile Leu Gln Tyr Cys Gln Glu Val Tyr Pro Glu Leu

65 70 75 80

Gln Ile Thr Asn Val Val Glu Ala Asn Gln Pro Val Thr Ile Gln Asn

85 90 95

Trp Cys Lys Arg Gly Arg Lys Gln Cys Lys Thr His Pro His Phe Val

100 105 110

Ile Pro Tyr Arg Cys Leu Val Gly Glu Phe Val Ser Asp Ala Leu Leu

115 120 125

Val Pro Asp Lys Cys Lys Phe Leu His Gln Glu Arg Met Asp Val Cys

130 135 140

Glu Thr His Leu His Trp His Thr Val Ala Lys Glu Thr Cys Ser Glu

145 150 155 160

Lys Ser Thr Asn Leu His Asp Tyr Gly Met Leu Leu Pro Cys Gly Ile

165

170

175

Asp Lys Phe Arg Gly Val Glu Phe Val Cys Cys Pro Leu Ala Glu Glu

180

185

190

Ser Asp Asn Val Asp Ser Ala Asp Ala Glu Glu Asp Asp Ser Asp Val

195

200

205

Trp Trp Gly Gly Ala Asp Thr Asp Tyr Ala Asp Gly Ser Glu Asp Lys

210

215

220

Val Val Glu Val Ala Glu Glu Glu Glu Val Ala Glu Val Glu Glu Glu

225

230

235

240

Glu Ala Asp Asp Asp Glu Asp Asp Glu Asp Gly Asp Glu Val Glu Glu

245

250

255

Glu Ala Glu Glu Pro Tyr Glu Glu Ala Thr Glu Arg Thr Thr Ser Ile

260

265

270

Ala Thr Thr Thr Thr Thr Thr Thr Glu Ser Val Glu Glu Val Val Arg

275

280

285

Val Pro Thr Thr Ala Ala Ser Thr Pro Asp Ala Val Asp Lys Tyr Leu

290

295

300

Glu Thr Pro Gly Asp Glu Asn Glu His Ala His Phe Gln Lys Ala Lys

305

310

315

320

Glu Arg Leu Glu Ala Lys His Arg Glu Arg Met Ser Gln Val Met Arg

325 330 335

Glu Trp Glu Glu Ala Glu Arg Gln Ala Lys Asn Leu Pro Lys Ala Asp

340 345 350

Lys Lys Ala Val Ile Gln His Phe Gln Glu Lys Val Glu Ser Leu Glu

355 360 365

Gln Glu Ala Ala Asn Glu Arg Gln Gln Leu Val Glu Thr His Met Ala

370 375 380

Arg Val Glu Ala Met Leu Asn Asp Arg Arg Arg Leu Ala Leu Glu Asn

385 390 395 400

Tyr Ile Thr Ala Leu Gln Ala Val Pro Pro Arg Pro Arg His Val Phe

405 410 415

Asn Met Leu Lys Lys Tyr Val Arg Ala Glu Gln Lys Asp Arg Gln His

420 425 430

Thr Leu Lys His Phe Glu His Val Arg Met Val Asp Pro Lys Lys Ala

435 440 445

Ala Gln Ile Arg Ser Gln Val Met Thr His Leu Arg Val Ile Tyr Glu

450 455 460

Arg Met Asn Gln Ser Leu Ser Leu Leu Tyr Asn Val Pro Ala Val Ala

465 470 475 480

Glu Glu Ile Gln Asp Glu Val Asp Glu Leu Leu Gln Lys Glu Gln Asn

485	490	495	
Tyr Ser Asp Asp Val Leu Ala Asn Met Ile Ser Glu Pro Arg Ile Ser			
500	505	510	
Tyr Gly Asn Asp Ala Leu Met Pro Ser Leu Thr Glu Thr Lys Thr Thr			
515	520	525	
Val Glu Leu Leu Pro Val Asn Gly Glu Phe Ser Leu Asp Asp Leu Gln			
530	535	540	
Pro Trp His Ser Phe Gly Ala Asp Ser Val Pro Ala Asn Thr Glu Asn			
545	550	555	560
Glu Val Glu Pro Val Asp Ala Arg Pro Ala Ala Asp Arg Gly Leu Thr			
565	570	575	
Thr Arg Pro Gly Ser Gly Leu Thr Asn Ile Lys Thr Glu Glu Ile Ser			
580	585	590	
Glu Val Asn Leu Asp Ala Glu Phe Arg His Asp Ser Gly Tyr Glu Val			
595	600	605	
His His Gln Lys Leu Val Phe Phe Ala Glu Asp Val Gly Ser Asn Lys			
610	615	620	
Gly Ala Ile Ile Gly Leu Met Val Gly Gly Val Val Ile Ala Thr Val			
625	630	635	640
Ile Val Ile Thr Leu Val Met Leu Lys Lys Lys Gln Tyr Thr Ser Ile			
645	650	655	

His His Gly Val Val Glu Val Asp Ala Ala Val Thr Pro Glu Glu Arg

660

665

670

His Leu Ser Lys Met Gln Gln Asn Gly Tyr Glu Asn Pro Thr Tyr Lys

675

680

685

Phe Phe Glu Gln Met Gln Asn Lys Lys

690

695

<210> 19

<211> 2094

<212> DNA

<213> Homo sapiens

<400> 19

atgctgcccc gtttggcaat gctcctgctg gcggcctgga cggtcggggc gctggaggta 60
 cccactgatg gtaactgctg cctgctgggt gaaccaccaga ttgccatgtt ctgtggcaga 120
 ctgaacatgc acatgaatgt ccaqaatggg aatgqgatt cagatccatc agggaccaaa 180
 acctgcattg ataccaagga aggcatactg cagtattgac aagaagtota cctgaactg 240
 cagatcacca atgtggtaga agccaaccaa ccagtgaaca tccagaactg gtgcaagcgg 300
 ggccgcaagc agtgcaagac ccataccacac tttgtgatc cctaccgctg cttagtgtgt 360
 gagtttgtaa gtgatgcctt tctcgttctt gacaagtga aattcttaca ccaggagagg 420
 atggatgttt gcgaaactca tcttcactgg cacaccgtcg ccaaagagac atgcagtgag 480
 aagagtacca acctgcatga ctacggcatg ttgctgcctt gcggaaatga caagtccga 540
 ggggttagagt ttgtgtgttg cccactgggt gaagaaagtg acaatgtgga ttctgtgat 600
 gcggaggagg atgactcgga tgcctggttg gcgggagcag acacagacta tgcagatggg 660
 agtgaagaca aagtagtaga agtagcagag gaggaagaag tggtgaggt ggaagaagaa 720
 gaagccgatg atgacagga cgtatgaggat gctgatgagg tagaagaaqa ggctgaggaa 780
 cctacgaag aagccacaga gagaaccacc agcattgcca ccaccaccac caccaccaca 840

gagtctgtgg aagaggtggt tegagttcct acaacagcag ccagtacccc tgatgcggtt 900
 gacaagtatc tccagacacc tggggatgag aatgaacatg cccatttcca gaaagcnaaa 960
 gagaggtctt aggcnaagca ccgagagaga atgtcccagg tcatgagaga atgggaagag 1020
 ccagaacgtc aagcaaaaga cttgcctaaa gctgataaga aggcagttat ccagcatttc 1080
 caggagaaaag tggaaatctt ggaacaggaa gcacccaacy agagacagca gctgggtggag 1140
 ccacacatgg ccagagtggc agccatgctc aatgaacgca gcgcctggc cctggagaa 1200
 tacatcacag ctctgcagga tgttctctct aggcctctgc acgtgttcaa tatgctaaag 1260
 aagtatgtcc gcgtagaaca gaaggacaga cagcacaacc taaagcattt ccagcagtgt 1320
 ccgctgtgtg atcccaagaa agccgctcag atccggtccc aggttatgac acacctccgt 1380
 gtgatttatg agcgcctgaa tcagtctctc tccctgctct acaacgtgcc tgcagtggcc 1440
 gaggagattc aggatgaagt tgatgagctg ctccagaaaag agcaaaaacta ttcagatgac 1500
 ctcttggcca acatgattag tgaaccaagg atcagttacg gaaacgatgc tctcatgcca 1560
 tctttgaccg aaacgaaaac caccgtggag ctcttctccg tgaatggaga gtccagcctg 1620
 gacgatctcc agccgtggca ttcttttggg gctgactctg tgcagccaa cacagaaaac 1680
 gaagttgagc ctgttgatgc ccgcctgct gcgcacagag gactgaaccac tgcaccaggt 1740
 tctgggttga caaatatcaa gacggaggag atctctgaag tgaagatgga tgcagaatc 1800
 ccacatgact caggatatca agttcatcat caaaaattg tgttctttgc aqaagatgtg 1860
 ggttcaaaaa aaggtgcaat cattggactc atgggtggcg gtgttctcat agcgacagt 1920
 atcttcatca ccttgggtgat gctgaagaag aaacagtaca catccattca tcatggtgtg 1980
 ctggagggtt acgcctgtgt caccacagag gagcgccacc tgtccaagat gcagcagaac 2040
 ggctacgaaa atccaaccta caagttcttt gacgagatgc agaacaagaa gtag 2094

<210> 20

<211> 607

<212> PRT

<213> Homo sapiens

<400> 20

Met Leu Pro Gly Leu Ala Leu Leu Leu Ala Ala Trp Thr Ala Arg

1 5 10 15

Ala Leu Glu Val Pro Thr Asp Gly Asn Ala Gly Leu Leu Ala Glu Pro

20

25

30

Gln Ile Ala Met Phe Cys Gly Arg Leu Asn Met His Met Asn Val Gln

35

40

45

Asn Gly Lys Trp Asp Ser Asp Pro Ser Gly Thr Lys Thr Cys Ile Asp

50

55

60

Thr Lys Glu Gly Ile Leu Gln Tyr Cys Gln Glu Val Tyr Pro Glu Leu

65

70

75

80

Gln Ile Thr Asn Val Val Glu Ala Asn Gln Pro Val Thr Ile Gln Asn

85

90

95

Trp Cys Lys Arg Gly Arg Lys Gln Cys Lys Thr His Pro His Phe Val

100

105

110

Ile Pro Tyr Arg Cys Leu Val Gly Glu Phe Val Ser Asp Ala Leu Leu

115

120

125

Val Pro Asp Lys Cys Lys Phe Leu His Gln Glu Arg Met Asp Val Cys

130

135

140

Glu Thr His Leu His Trp His Thr Val Ala Lys Glu Thr Cys Ser Glu

145

150

155

160

Lys Ser Thr Asn Leu His Asp Tyr Gly Met Leu Leu Pro Cys Gly Ile

165

170

175

Asp Lys Phe Arg Gly Val Glu Phe Val Cys Cys Pro Leu Ala Glu Glu

180	185	190	
Ser Asp Asn Val Asp Ser Ala Asp Ala Glu Glu Asp Asp Ser Asp Val			
195	200	205	
Trp Trp Gly Gly Ala Asp Thr Asp Tyr Ala Asp Gly Ser Glu Asp Lys			
210	215	220	
Val Val Glu Val Ala Glu Glu Glu Glu Val Ala Glu Val Glu Glu Glu			
225	230	235	240
Glu Ala Asp Asp Asp Glu Asp Asp Glu Asp Gly Asp Glu Val Glu Glu			
245	250	255	
Glu Ala Glu Glu Pro Tyr Glu Glu Ala Thr Glu Arg Thr Thr Ser Ile			
260	265	270	
Ala Thr Thr Thr Thr Thr Thr Thr Glu Ser Val Glu Glu Val Val Arg			
275	280	285	
Val Pro Thr Thr Ala Ala Ser Thr Pro Asp Ala Val Asp Lys Tyr Leu			
290	295	300	
Glu Thr Pro Gly Asp Glu Asn Glu His Ala His Phe Gln Lys Ala Lys			
305	310	315	320
Glu Arg Leu Glu Ala Lys His Arg Glu Arg Met Ser Gln Val Met Arg			
325	330	335	
Glu Trp Glu Glu Ala Glu Arg Gln Ala Lys Asn Leu Pro Lys Ala Asp			
340	345	350	

Lys Lys Ala Val Ile Gln His Phe Gln Glu Lys Val Glu Ser Leu Glu

355 360 365

Gln Glu Ala Ala Asn Glu Arg Gln Gln Leu Val Glu Thr His Met Ala

370 375 380

Arg Val Glu Ala Met Leu Asn Asp Arg Arg Arg Leu Ala Leu Glu Asn

385 390 395 400

Tyr Ile Thr Ala Leu Gln Ala Val Pro Pro Arg Pro Arg His Val Phe

405 410 415

Asn Met Leu Lys Lys Tyr Val Arg Ala Glu Gln Lys Asp Arg Gln His

420 425 430

Thr Leu Lys His Phe Glu His Val Arg Met Val Asp Pro Lys Lys Ala

435 440 445

Ala Gln Ile Arg Ser Gln Val Met Thr His Leu Arg Val Ile Tyr Glu

450 455 460

Arg Met Asn Gln Ser Leu Ser Leu Leu Tyr Asn Val Pro Ala Val Ala

465 470 475 480

Glu Glu Ile Gln Asp Glu Val Asp Glu Leu Leu Gln Lys Glu Gln Asn

485 490 495

Tyr Ser Asp Asp Val Leu Ala Asn Met Ile Ser Glu Pro Arg Ile Ser

500 505 510

Tyr Gly Asn Asp Ala Leu Met Pro Ser Leu Thr Glu Thr Lys Thr Thr

515

520

525

Val Glu Leu Leu Pro Val Asn Gly Glu Phe Ser Leu Asp Asp Leu Gln

530

535

540

Pro Trp His Ser Phe Gly Ala Asp Ser Val Pro Ala Asn Thr Glu Asn

545

550

555

560

Glu Val Glu Pro Val Asp Ala Arg Pro Ala Ala Asp Arg Gly Leu Thr

565

570

575

Thr Arg Pro Gly Ser Gly Leu Thr Asn Ile Lys Thr Glu Glu Ile Ser

580

585

590

Glu Val Lys Met Asp Ala Glu Phe Arg His Asp Ser Gly Tyr Glu Val

595

600

605

His His Gln Lys Leu Val Phe Phe Ala Glu Asp Val Gly Ser Asn Lys

610

615

620

Gly Ala Ile Ile Gly Leu Met Val Gly Gly Val Val Ile Ala Thr Val

625

630

635

640

Ile Phe Ile Thr Leu Val Met Leu Lys Lys Lys Gln Tyr Thr Ser Ile

645

650

655

His His Gly Val Val Glu Val Asp Ala Ala Val Thr Pro Glu Glu Arg

660

665

670

His Leu Ser Lys Met Gln Gln Asn Gly Tyr Glu Asn Pro Thr Tyr Lys

675

680

685

Phe Phe Glu Gln Met Gln Asn Lys Lys

690

695

<210> 21

<211> 1341

<212> DNA

<213> Homo sapiens

<400> 21

atgggtatga ttactggtgg aaagcaaatg ggtcggggat caaccacgaa cggcatccgg 60
 ctgctcctgc gcagcggcct gggggggccc cccctggggc tgcggtgccc cggggagacc 120
 gacgaagagc ccgaggagcc cggcgggagg ggcagctttg tggagatggt ggacaacctg 180
 aggggcaagt cggggcaggg ctactacgtg gagatgaccg tgggcagccc ccgcagagc 240
 tcaacatccc tgytggatac aggcagcagt aactttgcag tgggtgctgc cccccaccc 300
 ttcctgcacc gctactacca gaggcagctg tccagccat accgggacct ccggaagggt 360
 gtgtatgtgc cctacacca gggcaagtgg gaaggggagc tgggcaccca cctggtaagc 420
 atcccacatg gcccacagt cactgtgcgt gccaacattg ctgccatcac tgaatcagac 480
 aagttcttca tcaacggctc caactgggaa ggcacccctg ggtcggccta tgtgagatt 540
 gccaggcctg accactccct ggagccttcc ttgactctc tggtaaagca gaccacggt 600
 cccaacctct tctcctgca cctttgtggt gctggcttcc cctcaacca gtctgaagt 660
 ctggcctctg tcggaggag catgatcatt ggaggtatcg accactcgt gtacacagc 720
 agtctctggt atacacccat ccggcgggag tgggtattatg aggtcatcat tgtcgggtg 780
 gagatcaatg gacaggatct gaaaatggac tgcaaggagt acaactatga caagagcatt 840
 gtggacagtg gcaccaccaa ccttcgtttg cccaagaaag tgtttgaagc tgcagtcaa 900
 tccatcaagg cagcctctc caaggagaag ttcctgatg gttctgggt aggagagcag 960
 ctggtgtgct ggcaagcagg caccacccct tggaacattt tcccagtcac ctactctac 1020
 ctatggggtg aggttaccac ccagtccttc cgcacacca tcttccgca gcaataacct 1080
 cggccagtgg aagatgtgga cagtcacca gacgactgtt acaagttgc catctcacg 1140

tcatccacgg gcactgttat gggagctgtt atcatggagg gcttctacgt tgtctttgat 1200
 cgggcccga aacgaattgg ctttgcgtgc accccttccc atgtgcacga tgatttcagg 1260
 agggcagcgg tgggaaggccc ttttgcacc ttggacatgg aagactgtgg ctacaacatt 1320
 ctacaqacag atgagtcatg a 1341

<210> 23

<211> 446

<212> PRT

<213> Homo sapiens

<400> 22

Met Ala Ser Met Thr Gly Gly Gln Gln Met Gly Arg Gly Ser Thr Gln

1 5 10 15

His Gly Ile Arg Leu Pro Leu Arg Ser Gly Leu Gly Gly Ala Pro Leu

20 25 30

Gly Leu Arg Leu Pro Arg Glu Thr Asp Glu Glu Pro Glu Glu Pro Gly

35 40 45

Arg Arg Gly Ser Phe Val Glu Met Val Asp Asn Leu Arg Gly Lys Ser

50 55 60

Gly Gln Gly Tyr Tyr Val Glu Met Thr Val Gly Ser Pro Pro Gln Thr

65 70 75 80

Leu Asn Ile Leu Val Asp Thr Gly Ser Ser Asn Phe Ala Val Gly Ala

85 90 95

Ala Pro His Pro Phe Leu His Arg Tyr Tyr Gln Arg Gln Leu Ser Ser

100 105 110

Thr Tyr Arg Asp Leu Arg Lys Gly Val Tyr Val Pro Tyr Thr Gln Gly

115

120

125

Lys Trp Glu Gly Glu Leu Gly Thr Asp Leu Val Ser Ile Pro His Gly

130

135

140

Pro Asn Val Thr Val Arg Ala Asn Ile Ala Ala Ile Thr Glu Ser Asp

145

150

155

160

Lys Phe Phe Ile Asn Gly Ser Asn Trp Glu Gly Ile Leu Gly Leu Ala

165

170

175

Tyr Ala Glu Ile Ala Arg Pro Asp Asp Ser Leu Glu Pro Phe Phe Asp

180

185

190

Ser Leu Val Lys Gln Thr His Val Pro Asn Leu Phe Ser Leu His Leu

195

200

205

Cys Gly Ala Gly Phe Pro Leu Asn Gln Ser Glu Val Leu Ala Ser Val

210

215

220

Gly Gly Ser Met Ile Ile Gly Gly Ile Asp His Ser Leu Tyr Thr Gly

225

230

235

240

Ser Leu Trp Tyr Thr Pro Ile Arg Arg Glu Trp Tyr Tyr Glu Val Ile

245

250

255

Ile Val Arg Val Glu Ile Asn Gly Gln Asp Leu Lys Met Asp Cys Lys

260

265

270

Glu Tyr Asn Tyr Asp Lys Ser Ile Val Asp Ser Gly Thr Thr Asn Leu

275

280

285

Arg Leu Pro Lys Lys Val Phe Glu Ala Ala Val Lys Ser Ile Lys Ala

290

295

300

Ala Ser Ser Thr Glu Lys Phe Pro Asp Gly Phe Trp Leu Gly Glu Gln

305

310

315

320

Leu Val Cys Trp Gln Ala Gly Thr Thr Pro Trp Asn Ile Phe Pro Val

325

330

335

Ile Ser Leu Tyr Leu Met Gly Glu Val Thr Asn Gln Ser Phe Arg Ile

340

345

350

Thr Ile Leu Pro Gln Gln Tyr Leu Arg Pro Val Glu Asp Val Ala Thr

355

360

365

Ser Gln Asp Asp Cys Tyr Lys Phe Ala Ile Ser Gln Ser Ser Thr Gly

370

375

380

Thr Val Met Gly Ala Val Ile Met Glu Gly Phe Tyr Val Val Phe Asp

385

390

395

400

Arg Ala Arg Lys Arg Ile Gly Phe Ala Val Ser Ala Cys His Val His

405

410

415

Asp Glu Phe Arg Thr Ala Ala Val Glu Gly Pro Phe Val Thr Leu Asp

420

425

430

Met Glu Asp Cys Gly Tyr Asn Ile Pro Gln Thr Asp Glu Ser

435

440

445

<210> 23

<211> 1380

<212> DNA

<213> Homo sapiens

<400> 23

atggctagca tgactgggtg acagcaaatg ggtcgcggtat cgatgactat ctctgactct 60
 ccgcgtgaac aggaacgata caaccagcac ggcacccggt tgcacctgag cagcggcctg 120
 gggggcgccc cctgggggtt ggggtgccc cgggagaccg accaagagcc cgaggagccc 180
 ggccggaggg gcagctttgt ggagatgggt gacaacctga ggggcaagtc ggggcagggc 240
 tactacgtgg agatgacctt gggcagccc ccgcagagcc tcacctctct ggtggataca 300
 ggcagcagta actttgcagt ggggtgtgcc cccacacctt cctgtcatcg ctactaccag 360
 aggcagctgt ccagcacata ccgggacctc cggaagggtg tgtatgtgac ctacaccag 420
 ggcaagtggg aaggggagct gggcaccgac ctggttaagc tccccatgg ccccaacgtc 480
 actgtgcgtg ccaacattgc tgcctacctt gaatcagaca agttctctat caacggctcc 540
 aactgggaag gcacctggg gctggcctat gctgagattg ccaggcctga cgaacctctg 600
 gagctttctt ttgactctct ggtaaacgag acccacttcc ccaacctctt ctccctgac 660
 ctttgtgggt ctggcttccc cccaaccag ttggaagtgc tggcctctgt cggaggggagc 720
 atgatacttg gaggtatcga ccactcgtg tacacaggca gtctctggta tacaccatc 780
 cggcgggaat ggtattatga ggtcatcatt gtcgggggtg agatcaatgg acaggatctg 840
 aaaatggact gcaaggagta caactatgac aagagcattg tggacagtgg caccaccaac 900
 cttcgtttgc ccaagaaagt gtttyaagct gcagtcacat ccatcaaggc agcctctctc 960
 acggagaagt tccctgatgg tttctggcta ggagagcagc tgggtgtgct gcaagcagga 1020
 accacccctt ggaacatttt ccagtccttc tcaactctac taatgggtga ggttaaccaac 1080
 caatccttcc gcataccat ccttcgcgac caatacctgc qccactgga agatgtggcc 1140
 acgtcccaag acgactgtta caagtttgc atctcacagt catccacggg cactgtttatg 1200
 ggagctgcta tcatggaggg cttctacgtt gtctttgac gggcccgaaa acgaattggc 1260
 tttgtgtca gccttgcca tgtgcacgat gatttcagga cggcagcggg ggaaggccct 1320

tttgtcactt tggacatgga agactgtgga tacaacattc cacagacaga tgagtcatga 1360

<310> 24

<311> 459

<312> PRT

<313> Homo sapiens

<400> 24

Met Ala Ser Met Thr Gly Gly Gln Gln Met Gly Arg Gly Ser Met Thr

1 5 10 15

Ile Ser Asp Ser Pro Arg Glu Gln Asp Gly Ser Thr Gln His Gly Ile

20 25 30

Arg Leu Pro Leu Arg Ser Gly Leu Gly Gly Ala Pro Leu Gly Leu Arg

35 40 45

Leu Pro Arg Glu Thr Asp Glu Glu Pro Glu Glu Pro Gly Arg Arg Gly

50 55 60

Ser Phe Val Glu Met Val Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly

65 70 75 80

Tyr Tyr Val Glu Met Thr Val Gly Ser Pro Pro Gln Thr Leu Asn Ile

85 90 95

Leu Val Asp Thr Gly Ser Ser Asn Phe Ala Val Gly Ala Ala Pro His

100 105 110

Pro Phe Leu His Arg Tyr Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg

115 120 125

Asp Leu Arg Lys Gly Val Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu

130

135

140

Gly Glu Leu Gly Thr Asp Leu Val Ser Ile Pro His Gly Pro Asn Val

145

150

155

160

Thr Val Arg Ala Asn Ile Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe

165

170

175

Ile Asn Gly Ser Asn Trp Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu

180

185

190

Ile Ala Arg Pro Asp Asp Ser Leu Glu Pro Phe Phe Asp Ser Leu Val

195

200

205

Lys Gln Thr His Val Pro Asn Leu Phe Ser Leu His Leu Cys Gly Ala

210

215

220

Gly Phe Pro Leu Asn Gln Ser Glu Val Leu Ala Ser Val Gly Gly Ser

225

230

235

240

Met Ile Ile Gly Gly Ile Asp His Ser Leu Tyr Thr Gly Ser Leu Trp

245

250

255

Tyr Thr Pro Ile Arg Arg Glu Trp Tyr Tyr Glu Val Ile Ile Val Arg

260

265

270

Val Glu Ile Asn Gly Gln Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn

275

280

285

Tyr Asp Lys Ser Ile Val Asp Ser Gly Thr Thr Asn Leu Arg Leu Pro

290

295

300

Lys Lys Val Phe Glu Ala Ala Val Lys Ser Ile Lys Ala Ala Ser Ser

305

310

315

320

Thr Glu Lys Phe Pro Asp Gly Phe Trp Leu Gly Glu Gln Leu Val Cys

325

330

335

Trp Gln Ala Gly Thr Thr Pro Trp Asn Ile Phe Pro Val Ile Ser Leu

340

345

350

Tyr Leu Met Gly Glu Val Thr Asn Gln Ser Phe Arg Ile Thr Ile Leu

355

360

365

Pro Gln Gln Tyr Leu Arg Pro Val Glu Asp Val Ala Thr Ser Gln Asp

370

375

380

Asp Cys Tyr Lys Phe Ala Ile Ser Gln Ser Ser Thr Gly Thr Val Met

385

390

395

400

Gly Ala Val Ile Met Glu Gly Phe Tyr Val Val Phe Asp Arg Ala Arg

405

410

415

Lys Arg Ile Gly Phe Ala Val Ser Ala Cys His Val His Asp Glu Phe

420

425

430

Arg Thr Ala Ala Val Glu Gly Pro Phe Val Thr Leu Asp Met Glu Asp

435

440

445

Cys Gly Tyr Asn Ile Pro Gln Thr Asp Glu Ser

450

455

<210> 25

<211> 1302

<212> DNA

<213> Homo sapiens

<100> 25

atgaactcagc atgggtattcg tatgcacactg cgtagcgggtc tgggtgggtgc tccactgggtc 60
 ctgcgtctgc cccgggagac cgaagaagag ccggaggagc ccggccggag gggcagcttt 120
 gtggacatgg tggacaacct gaggggcaag tcggggcagg gctactacgt ggagatgacc 180
 gtgggcagcc ccccgagac gctcaacatc ctggtggata caggcagcag taactttgca 240
 gtgggtgctg cccccaccc cttcttgcac cgtactacc agaggcagct gtccagcaca 300
 taccgggacc tcgggaaggg tgtglatgtg cctacaccc agggcaagtg ggaaggggag 360
 ctgggcaccc acctggtaag catcccccac ggcaccaacg tcactgtgcg tggcaacatt 420
 gctgccatca ctgaatcaga caagttcttc atcaacggct ccaactggga aggcacctcg 480
 gggttggtct atgctgagat tgcaggcct gaagactccc tggagccttt ctttgactct 540
 ctggttaage agacccaagt tcccaacctc ttctccctgc acctttgtgg tcttggttc 600
 cccctcaacc actctqaaqt gctggcctct gtccggaggga gcatgatcat tggagggtac 660
 gaccactcgc tgtacacagg cagtctcttg tatacaccca tcgggcggga gtggtattat 720
 gaggtcatca ttgtgcgggt ggagatcaat ggacaggatc tgaatatgga ctgcaaggag 780
 tacaactatg acaagagcat tgtggacagt ggcaccacca accttcgttt gcccaagaaa 840
 gtgtttgaag ctgcagtcaa atccatcaag gcagcctcct ccacggagaa gttccctgat 900
 ggtttctggc taggagagca gctggtgtgc tggcaaggag gcaccacccc ttggaacatt 960
 ttcacagcca tctcactcta cctaattgggt gaggttatca accagtcctt ccgcacacc 1020
 atctttccgc agcaatacct ggggccagtg gaagatgtgg ccacgtccca agacgactgt 1080
 tacaagtttg ccattccaca gtcacccacg ggcactgtta tgggagctgt tatcatggag 1140
 ggcttctacg ttgtctttga tcgggcccga aaacgaattg gctttgtgtg cagcgcttgc 1200
 tatgtgcacg atgagttcag gacggcagcg gtggaaggcc cttttgtcac cttggacatg 1260
 gaagactgtg gctacaacat tccacagaca gatgagtcac ga 1302

<210> 26

<211> 433

<212> PRT

<213> Homo sapiens

<400> 26

Met Thr Gln His Gly Ile Arg Leu Pro Leu Arg Ser Gly Leu Gly Gly

1 5 10 15

Ala Pro Leu Gly Leu Arg Leu Pro Arg Glu Thr Asp Glu Glu Pro Glu

20 25 30

Glu Pro Gly Arg Arg Gly Ser Phe Val Glu Met Val Asp Asn Leu Arg

35 40 45

Gly Lys Ser Gly Gln Gly Tyr Tyr Val Glu Met Thr Val Gly Ser Pro

50 55 60

Pro Gln Thr Leu Asn Ile Leu Val Asp Thr Gly Ser Ser Asn Phe Ala

65 70 75 80

Val Gly Ala Ala Pro His Pro Phe Leu His Arg Tyr Tyr Gln Arg Gln

85 90 95

Leu Ser Ser Thr Tyr Arg Asp Leu Arg Lys Gly Val Tyr Val Pro Tyr

100 105 110

Thr Gln Gly Lys Trp Glu Gly Glu Leu Gly Thr Asp Leu Val Ser Ile

115 120 125

Pro His Gly Pro Asn Val Thr Val Arg Ala Asn Ile Ala Ala Ile Thr

130 135 140

Glu Ser Asp Lys Phe Phe Ile Asn Gly Ser Asn Trp Glu Gly Ile Leu

145 150 155 160

Gly Leu Ala Tyr Ala Glu Ile Ala Arg Pro Asp Asp Ser Leu Glu Pro

165 170 175

Phe Phe Asp Ser Leu Val Lys Gln Thr His Val Pro Asn Leu Phe Ser

180 185 190

Leu His Leu Cys Gly Ala Gly Phe Pro Leu Asn Gln Ser Glu Val Leu

195 200 205

Ala Ser Val Gly Gly Ser Met Ile Ile Gly Gly Ile Asp His Ser Leu

210 215 220

Tyr Thr Gly Ser Leu Trp Tyr Thr Pro Ile Arg Arg Glu Trp Tyr Tyr

225 230 235 240

Glu Val Ile Ile Val Arg Val Glu Ile Asn Gly Gln Asp Leu Lys Met

245 250 255

Asp Cys Lys Glu Tyr Asn Tyr Asp Lys Ser Ile Val Asp Ser Gly Thr

260 265 270

Thr Asn Leu Arg Leu Pro Lys Lys Val Phe Glu Ala Ala Val Lys Ser

275 280 285

Ile Lys Ala Ala Ser Ser Thr Glu Lys Phe Pro Asp Gly Phe Trp Leu

290 295 300
Gly Glu Gln Leu Val Cys Trp Gln Ala Gly Thr Thr Pro Trp Asn Ile
305 310 315 320
Phe Pro Val Ile Ser Leu Tyr Leu Met Gly Glu Val Thr Asn Gln Ser
325 330 335
Phe Arg Ile Thr Ile Leu Pro Gln Gln Tyr Leu Arg Pro Val Glu Asp
340 345 350
Val Ala Thr Ser Gln Asp Asp Cys Tyr Lys Phe Ala Ile Ser Gln Ser
355 360 365
Ser Thr Gly Thr Val Met Gly Ala Val Ile Met Glu Gly Phe Tyr Val
370 375 380
Val Phe Asp Arg Ala Arg Lys Arg Ile Gly Phe Ala Val Ser Ala Cys
385 390 395 400
His Val His Asp Glu Phe Arg Thr Ala Ala Val Glu Gly Pro Phe Val
405 410 415
Thr Leu Asp Met Glu Asp Cys Gly Tyr Asn Ile Pro Gln Thr Asp Glu
420 425 430
Ser

<210> 27

<211> 1278

<212> DNA

<213> Homo sapiens

<400> 27

```

atggctagca tgactggtgg acagcaaatg ggtcgcgga cgtgactat ctctgactct 60
ccgctggact ctggtatcga aaccgacgga tcctttgtgg agatggtgga caacctgagg 120
ggcaagtctg ggcagggcta ctactggag atgacctgg gcagccccc gcagacgctc 180
aacatctctg tggatagagg cagcagtaac ttgcagtg gtyctcccc ccaccccttc 240
ctgcctcgt actaccagag gcagctgtcc agcacatacc gggacctccg gaagggtgtg 300
tatgtgcctt acacccaggg caagtgggaa ggggagctgg ccaccgacct ggttaagcat 360
ccccatggcc ccaacgtcac tgtcgtgccc aacattctct ccatcactga atcagacaag 420
ttcttcctca accgctccaa ctgggaagga atctctgggc tggcctatgc tgagattgcc 480
aggcctgacg actccctgga gcctttcttt gactctctgg taaagcagac ccacgttccc 540
aacctcttct cctcgacct ttgtggtgct ggttccccc tcaaccagtc tgaagtgtct 600
gcctctgttg gagggagcat gatcattgga ggtatcgacc actcgtgtga cacaggcagt 660
ctctggtata caccatccg gcgggagtggt tattatgagg tcatcattgt gcgggtggag 720
atcaatggac aggatctgaa aatggactgc aaggagtaca actatgacaa gagcattgtg 780
gacagtggca cccccaacct tcgtttgccc aagaaagtgt ttgaagctgc agtcaaatcc 840
atcaaggcag cctcctccac gcagaagttc cctgatggtt tctggctagg aagacagctg 900
gtgtgctggc aagcaggcac cccccctgg aacattttcc cagtcatctc actctaccta 960
atgggtgagg ttaccaacca gtccctccgc atcaccatcc ttccgcagca atacctgcgg 1020
tcagtggaag atgtggccac gtcccaagac gactgttaca agtttgccat ctacagtc 1080
cccaagggca ctgttatggg agctgttata atggagggtt tctacgttgt ctttgatcgg 1140
gcccgaaaac gaattggctt tctgtcagc gcttgccatg tgcacgatga gttcaggacg 1200
gcagcgggtg aaggcccttt tctaccttg gacatggaag actgtggcta caacattcca 1260
cagacagatg agtcctga

```

1278

<210> 28

<211> 425

<212> PRT

<213> Homo sapiens

<400> 28

Met Ala Ser Met Thr Gly Gly Gln Gln Met Gly Arg Gly Ser Met Thr

1 5 10 15

Ile Ser Asp Ser Pro Leu Asp Ser Gly Ile Glu Thr Asp Gly Ser Phe

20 25 30

Val Glu Met Val Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr

35 40 45

Val Glu Met Thr Val Gly Ser Pro Pro Gln Thr Leu Asn Ile Leu Val

50 55 60

Asp Thr Gly Ser Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe

65 70 75 80

Leu His Arg Tyr Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu

85 90 95

Arg Lys Gly Val Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu

100 105 110

Leu Gly Thr Asp Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val

115 120 125

Arg Ala Asn Ile Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn

130 135 140

Gly Ser Asn Trp Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile Ala

145	150	155	160
Arg Pro Asp Asp Ser Leu Glu Pro Phe Phe Asp Ser Leu Val Lys Gln			
	165	170	175
Thr His Val Pro Asn Leu Phe Ser Leu His Leu Cys Gly Ala Gly Phe			
	180	185	190
Pro Leu Asn Gln Ser Glu Val Leu Ala Ser Val Gly Gly Ser Met Ile			
	195	200	205
Ile Gly Gly Ile Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr			
	210	215	220
Pro Ile Arg Arg Glu Trp Tyr Tyr Glu Val Ile Ile Val Arg Val Glu			
	225	230	235
Ile Asn Gly Gln Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp			
	245	250	255
Lys Ser Ile Val Asp Ser Gly Thr Thr Asn Leu Arg Leu Pro Lys Lys			
	260	265	270
Val Phe Glu Ala Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu			
	275	280	285
Lys Phe Pro Asp Gly Phe Trp Leu Gly Glu Gln Leu Val Cys Trp Gln			
	290	295	300
Ala Gly Thr Thr Pro Trp Asn Ile Phe Pro Val Ile Ser Leu Tyr Leu			
	305	310	315
			320

Met Gly Glu Val Thr Asn Gln Ser Phe Arg Ile Thr Ile Leu Pro Gln

325

330

335

Gln Tyr Leu Arg Pro Val Glu Asp Val Ala Thr Ser Gln Asp Asp Cys

340

345

350

Tyr Lys Phe Ala Ile Ser Gln Ser Ser Thr Gly Thr Val Met Gly Ala

355

360

365

Val Ile Met Glu Gly Phe Tyr Val Val Phe Asp Arg Ala Arg Lys Arg

370

375

380

Ile Gly Phe Ala Val Ser Ala Cys His Val His Asp Glu Phe Arg Thr

385

390

395

400

Ala Ala Val Glu Gly Pro Phe Val Thr Leu Asp Met Glu Asp Cys Gly

405

410

415

Tyr Asn Ile Pro Gln Thr Asp Glu Ser

420

425

<210> 29

<211> 1362

<212> DNA

<213> Homo sapiens

<400> 29

atggcccaag cctgcccctg gctcctgctg tygatgggcy cyggagtgcg gctgcccac 60

ggcaccacagc acggcatcgc gctgcccctg cgcagcggcc tggggggggc cccctgggg 120

ctggcgctgc ccggggagac cgacgaagag ccggaggaga ccggccggag gggcagcttt 180
 gtggagatgg tggacaacct gaggggcaag tgggggcagg gctactacgt ggaatgacc 240
 gtgggcagcc cccgcagac gctcaacatc ctgggtggata caggcagcag taactttgca 300
 gtgggtgctg cccccaccc ctctctgcat cgtactacc agaggcagct gtccagaca 360
 taacgggacc tccggaaggg tjtgtatgtg cctacaccc agggcaagtg ggaaggggag 420
 ctgggcacgg acctggtaag catcccccac ggcccacacg taactgtgctg tgccacatt 480
 gctgcatca ctgaatcaga caagtctctc atcaacggct ccaactggga agccatcttg 540
 gggttggtct atgttgatg tggcaggcct gacgaatccc tggagccttt ctctgactct 600
 ctggtaaaag agacccacgt tcccaacctc ctctccctgc accttggttg tctggcttc 660
 tccctcaac agtgtgaagt gctggcctct gtgggagga gcatgacat tggaggtatc 720
 caccactcgc tgtacacagg cagtctctgg tataacccc tccggcgga gtjgtattat 780
 caggctcatc ttgtgcgggt ggagatcat ggacaggatc tgaatatgga ctgcaaggag 840
 tacaactatg acaagagcat tgtggacagt ggcaccacca accttcgttt gcccagaaa 900
 gtgtttgaag ctgcagtcac atccatcaag gcagctctct ccacggagaa gttccctgat 960
 ggtttctggc taggagagca gctggtgtgc tggcaagcag gcaccacccc ttggaacatt 1020
 ctccagtcac tctactctc cctaatgggt gaggttaaca accagtcctt ccgcatcacc 1080
 atctctccgc agcaatacct cgggcccagtg gaagatgttg ccaagtccca agacagctgt 1140
 tacaagtttg ccatctcaca gtcctccacg ggcactgtta tgggagctgt tatcatggag 1200
 gctctctacg ttgtctttga tgggcccga aaacguattg gctttgtgt cagcgttgc 1260
 catgtgcacg atgagttcag cagggcagcg gtggaaggcc ctcttgctac ctggacatg 1320
 gaagactgtg cctacaacat tccacagaca gatgagtcac ga 1362

<210> 30

<211> 453

<212> PRT

<213> Homo sapiens

<400> 30

Met Ala Gln Ala Leu Pro Trp Leu Leu Leu Trp Met Gly Ala Gly Val

1

5

10

15

Leu Pro Ala His Gly Thr Gln His Gly Ile Arg Leu Pro Leu Arg Ser
 ~~~~~ 20 25 30  
 Gly Leu Gly Gly Ala Pro Leu Gly Leu Arg Leu Pro Arg Glu Thr Asp  
 ~~~~~ 35 40 45  
 Glu Glu Pro Gln Glu Pro Gly Arg Arg Gly Ser Phe Val Glu Met Val
 ~~~~~ 50 55 60  
 Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val Glu Met Thr  
 ~~~~~ 65 70 75 80  
 Val Gly Ser Pro Pro Gln Thr Leu Asn Ile Leu Val Asp Thr Gly Ser
 ~~~~~ 85 90 95  
 Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe Leu His Arg Tyr  
 ~~~~~ 100 105 110  
 Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg Lys Gly Val
 ~~~~~ 115 120 125  
 Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu Leu Gly Thr Asp  
 ~~~~~ 130 135 140  
 Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val Arg Ala Asn Ile
 ~~~~~ 145 150 155 160  
 Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly Ser Asn Trp  
 ~~~~~ 165 170 175  
 Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile Ala Arg Pro Asp Asp
 ~~~~~

180	185	190	
Ser Leu Glu Pro Phe Phe Asp Ser Leu Val Lys Gln Thr His Val Pro			
195	200	205	
Asn Leu Phe Ser Leu Gln Leu Cys Gly Ala Gly Phe Pro Leu Asn Gln			
210	215	220	
Ser Glu Val Leu Ala Ser Val Gly Gly Ser Met Ile Ile Gly Gly Ile			
225	230	235	240
Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr Pro Ile Arg Arg			
245	250	255	
Glu Trp Tyr Tyr Glu Val Ile Ile Val Arg Val Glu Ile Asn Gly Gln			
260	265	270	
Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp Lys Ser Ile Val			
275	280	285	
Asp Ser Gly Thr Thr Asn Leu Arg Leu Pro Lys Lys Val Phe Glu Ala			
290	295	300	
Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu Lys Phe Pro Asp			
305	310	315	320
Gly Phe Trp Leu Gly Glu Gln Leu Val Cys Trp Gln Ala Gly Thr Thr			
325	330	335	
Pro Trp Asn Ile Phe Pro Val Ile Ser Leu Tyr Leu Met Gly Glu Val			
340	345	350	

Thr Asn Gln Ser Phe Arg Ile Thr Ile Leu Pro Gln Gln Tyr Leu Arg

355

360

365

Pro Val Glu Asp Val Ala Thr Ser Gln Asp Asp Cys Tyr Lys Phe Ala

370

375

380

Ile Ser Gln Ser Ser Thr Gly Thr Val Met Gly Ala Val Ile Met Glu

385

390

395

400

Gly Phe Tyr Val Val Phe Asp Arg Ala Arg Lys Arg Ile Gly Phe Ala

405

410

415

Val Ser Ala Cys His Val His Asp Glu Phe Arg Thr Ala Ala Val Glu

420

425

430

Gly Pro Phe Val Thr Leu Asp Met Glu Asp Cys Gly Tyr Asn Ile Pro

435

440

445

Gln Thr Asp Glu Ser

450

<210> 31

<211> 1380

<212> DNA

<213> Homo sapiens

<400> 31

atggcccaag cctgcccgtg gctcctgctg tggatggggc egggagtget gctgcccac 60

ggcaccacagc acggcatcgg gctgcccctg cgcaggggac tggggggcgc cccctgggg 120

ctggcgcttc cccgggagac cgcggaagag ccgaggagac cgggcgggag gggcagcttc 180  
 gtggagatgg tggacaacct gaggggcaag tcggggcagg gctactacgt gtagatgacc 240  
 gtgggcagac cccgcagac gctcaacatc ctggtggata caggcagcag taactttgca 300  
 gtgggtgctg cccccccc ctctctgcat cgtactacc agaggcagct gtccagcaca 360  
 taacgggacc tcgggaaggg tgggtatgtg ccctacacc agggcaagtg ggaaggggag 420  
 ctgggcacgg acctggttaag catcccccac ggcaccaacg tcaactgtgcg tgccaacatt 480  
 gctgcacatc ctgaatcaga caagttcttc atcaacggct ccaactggga aggcacctctg 540  
 cggctggcct atgctgagat tgcagacct gacgactcc cggagcttt ctctgaactct 600  
 ctggtaaacg agatccacgt tcccaacctc tctccctgca acctttgtgg tctgtgcttc 660  
 cccctcaacc agtctgaagt gctggcctct gcgggaggga gcctgatcat tggaggtatc 720  
 caccactcgc tgtacacagg cagtctctgg tatacaccca tccggcggga gtggtattat 780  
 gaggctatca ttgtgcgggt ggagatcaat ggacaggatc tgaaaatgga ctgcaaggag 840  
 tacaactatg acaagagcat tgtggacagt ggcaccacca accttcgttt gcccaagaaa 900  
 gtgtttgaag ctgcagtcac atccatcang gcagctctct ccaaggagaa gtctccctgat 960  
 cgtttctgga taggagagca gctggtgtgc tggcaagcag ccaccacccc ttgcaacatt 1020  
 tccacagtc cctcaactct cctaatgggt gagggtacac accagtcctt ccgcacacc 1080  
 atctctccgc agcaatacct gcggccagtg gaagatgtgg ccagctccca agacgactgt 1140  
 tacaagtttg ccctctcaca gtcctccacg ggcactgtta tgggagctgt tatcatggag 1200  
 ggctctacg ttgtcttga tcgggcccga aaacgaattg gctttgctgt cagcgtttgc 1260  
 catgtgcacg atgagttcag gacggcagcg gtggaaggcc cttttgctac ctggacatg 1320  
 gaagactgtg gctacaacat tccacagaca gatgagtcac agcagcagca gcagcagtga 1380

<110> 32

<111> 459

<112> PRT

<113> Homo sapiens

<400> 32

Met Ala Gln Ala Leu Pro Trp Leu Leu Leu Trp Met Gly Ala Gly Val

1

5

10

15

Leu Pro Ala His Gly Thr Gln His Gly Ile Arg Leu Pro Leu Arg Ser

20

25

30

Gly Leu Gly Gly Ala Pro Leu Gly Leu Arg Leu Pro Arg Glu Thr Asp

35

40

45

Glu Glu Pro Glu Glu Pro Gly Arg Arg Gly Ser Phe Val Glu Met Val

50

55

60

Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val Glu Met Thr

65

70

75

80

Val Gly Ser Pro Pro Gln Thr Leu Asn Ile Leu Val Asp Thr Gly Ser

85

90

95

Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe Leu His Arg Tyr

100

105

110

Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg Lys Gly Val

115

120

125

Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu Leu Gly Thr Asp

130

135

140

Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val Arg Ala Asn Ile

145

150

155

160

Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly Ser Asn Trp

165

170

175

Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile Ala Arg Pro Asp Asp

180	185	190	
Ser Leu Glu Pro Phe Phe Asp Ser Leu Val Lys Gln Thr His Val Pro			
195	200	205	
Asn Leu Phe Ser Leu Gln Leu Cys Gly Ala Gly Phe Pro Leu Asn Gln			
210	215	220	
Ser Glu Val Leu Ala Ser Val Gly Gly Ser Met Ile Ile Gly Gly Ile			
225	230	235	240
Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr Pro Ile Arg Arg			
245	250	255	
Glu Trp Tyr Tyr Glu Val Ile Ile Val Arg Val Glu Ile Asn Gly Gln			
260	265	270	
Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp Lys Ser Ile Val			
275	280	285	
Asp Ser Gly Thr Thr Asn Leu Arg Leu Pro Lys Lys Val Phe Glu Ala			
290	295	300	
Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu Lys Phe Pro Asp			
305	310	315	320
Gly Phe Trp Leu Gly Glu Gln Leu Val Cys Trp Gln Ala Gly Thr Thr			
325	330	335	
Pro Trp Asn Ile Phe Pro Val Ile Ser Leu Tyr Leu Met Gly Glu Val			
340	345	350	

Thr Asn Gln Ser Phe Arg Ile Thr Ile Leu Pro Gln Gln Tyr Leu Arg

355

360

365

Pro Val Glu Asp Val Ala Thr Ser Gln Asp Asp Cys Tyr Lys Phe Ala

370

375

380

Ile Ser Gln Ser Ser Thr Gly Thr Val Met Gly Ala Val Ile Met Glu

385

390

395

400

Gly Phe Tyr Val Val Phe Asp Arg Ala Arg Lys Arg Ile Gly Phe Ala

405

410

415

Val Ser Ala Cys His Val His Asp Glu Phe Arg Thr Ala Ala Val Glu

420

425

430

Gly Pro Phe Val Thr Leu Asp Met Glu Asp Cys Gly Tyr Asn Ile Pro

435

440

445

Gln Thr Asp Glu Ser His His His His His His

450

455

<210> 33

<211> 25

<212> PRT

<213> Homo sapiens

<400> 33

Ser Glu Gln Gln Arg Arg Pro Arg Asp Pro Glu Val Val Asn Asp Glu

1

5

10

15

"  
Ser Ser Leu Val Arg His Arg Trp Lys  
"

" 20 25  
"

"  
<210> 34  
"

"  
<211> 19  
"

"  
<212> PRT  
"

"  
<213> Homo sapiens  
"

"  
<400> 34  
"

Ser Glu Gln Leu Arg Gln Gln His Asp Asp Phe Ala Asp Asp Ile Ser

1 5 10 15

"  
Leu Leu Lys  
"

"  
<210> 35  
"

"  
<211> 29  
"

"  
<212> DNA  
"

"  
<213> Homo sapiens  
"

"  
<400> 35  
"

gtggatccac ccagcacggc atccggctg

29

"  
<210> 36  
"

"  
<211> 36  
"

"  
<212> DNA  
"

"  
<213> Homo sapiens  
"



<400> 36

gaaagcttcc atgactcacc tgcctgtgga atgttg

36

<210> 37

<211> 39

<212> DNA

<213> Homo sapiens

<400> 37

gacgatgac tatarctgac tctcggcgtg aacaggacg

39

<210> 38

<211> 39

<212> DNA

<213> Homo sapiens

<400> 38

gacggctctt gtccacggcg agagtcagag atagtcacc

39

<210> 39

<211> 77

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Hu-Asp2

<400> 39

cggcacccgg ctgcacctgc gtagcgggtt gggcgggtgt ccaactgggtc tgcgtctgcc 60

ccggggagacc gacgaag

77

<210> 40

<211> 77

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Hu-Asp2

<400> 40

cttcgtcggc ctcctggggc agatgcagac ccagtggagt accaccaga ccgtatgca 60  
ggggcagccg gatgccg 77

<210> 41

<211> 51

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Caspase 8

Cleavage Site

<400> 41

gacgatgac tatctctgac tctcctcgg actctggat cgaaccgac g 51

<210> 42

<211> 51

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Caspase 8

## Cleavage Site

&lt;400&gt; 42

atcccgctccg ttcccatacc aagctccacc ggagagtcag agatactcat c

51

&lt;210&gt; 43

&lt;211&gt; 32

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 43

aaggatcctt tgtggagatg gtggacaacc tg

32

&lt;210&gt; 44

&lt;211&gt; 36

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 44

gaaagcttcc atgactcacc tgtctgtgga atgttg

36

&lt;210&gt; 45

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: 6-His tag

&lt;400&gt; 45

gategcatca tcaccatcac catg

24

<210> 46

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: 6-His tag

<400> 46

gatacatggt gatggtgatg atgc

24

<210> 47

<211> 354

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Introduce KK

motif

<400> 47

hottaanvtt nnnnngaactg accactcgac caggttcnrr maemhadata ragrahntsn 60  
ayrsksoana yrtawsddcg tmsnwrmans ymbarahr0g actgaccact cgaccagggt 120  
gsnayrsnay rh0dtgactg accactcgac caggttcact snayrtcsn asranrmdt 180  
gsnayrtena merstwr0t dthharmaca lngactgacc actcgaccag gttcttdgda 240  
n0bd0cda00 a0ca0rtntn ygtabwrddc mntsmmaryn rmatndcmnt smmarynrma 300  
tnsksoyemb abeerhvgrr eer0rsmers twrddcmntm swrddcwrd cmt 354

<210> 48

<211> 462

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Introduce KK

motif

<400> 48

```

bbttaanttn nnnknogaat taaattccag cacactggct acttcttgtt ctgcctctca 60
aagaachnrm acmhadata agrahntsna yrsks0snay rtawddcgt msnwrmansy 120
nbarahr0cg aattaaatte cagcacactg gctacttctt gttctgcata tcaaagaacs 180
nayrsnayrn dhogaatta aattccagca cactggctac ttcttgttct gcctctcaca 240
gaacgaasna yttcsnash anmadtsen ayrtcnamer stwrddcgks kdharmaca 300
dhogaattaa attccagcac actggctact tcttgttctg cactcacaag aactldgdan 360
db0cda00a0 ca0rtntryh kktabwrddc mntsmmaryn rnatndemnt smmarynrma 420
tntdecmbbe tekkmcrstw rddemntmsw rddcwrddem nt 462

```

<210> 49

<211> 380

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Introduce KK

motif

<400> 49

```

bbttaanttn nnnknogaat taaattccag cacactggct abnrmacmha dataragrah 60
ntsnayrks 0snayrtaws ddcgtmshwr mansymbara hr0ogaatta aattccagca 120
cactggctas nayrsnayrn dhogaatta aattccagca cactggctag aasnayrttc 180
snasnarmma dtesnayrtc namerstwrdd 0cmdharmah dhogaatt aaattccagc 240

```

acastggeta ttdgdan0b0 cda0Ca0ca0 rntnymknt abwrddcmnt smmzynrma 300  
 Indemntsmm arynrmatns ksCyrbmmc rbanbotkmk mg0g0gscr0 rsmcrstwrđ 360  
 demntmewđ dwwrddcmnt 380



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification:</b> <b>C12N 15/57, C07K 14/47,</b> <b>C07K 16/18, C07K 19/00,</b> <b>C12N 1/21, C12N 5/10,</b> <b>C12N 9/64, C12N 15/12,</b> <b>C12N 15/62, C12N 15/85,</b> <b>C12Q 1/37, G01N 33/68</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 00/17369</b> <b>(43) International Publication Date:</b> 30 March 2000 (30.03.2000)
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------	--------------------------------------------------------------------------------------------------------------------------------------------------

**(21) International Application Number:** PCT/US99/20881**Published****(22) International Filing Date:** 23 September 1999 (23.09.1999)**(30) Priority Data:**  
60/101,594 24 September 1998 (24.09.1998) US**(60) Parent Application or Grant**

PHARMACIA & UPJOHN COMPANY [ ]; () GURNEY,  
Mark, E. [ ]; () BIENKOWSKI, Michael, Jerome [ ];  
() HEINRIKSON, Robert, Leroy [ ]; () PARODI, Luis, A. [ ];  
() YAN, Riqiang [ ]; () GURNEY, Mark, E. [ ];  
() BIENKOWSKI, Michael, Jerome [ ]; () HEINRIKSON,  
Robert, Leroy [ ]; () PARODI, Luis, A. [ ]; () YAN, Riqiang  
[ ]; () WOOTTON, Thomas, A. [ ]

**(54) Title:** ALZHEIMER'S DISEASE SECRETASE**(54) Titre:** SECRETASE DE LA MALADIE D'ALZHEIMER**(57) Abstract**

The present invention provides the enzyme and enzymatic procedures for cleaving the 'beta' secretase cleavage site of the APP protein and associated nucleic acids, peptides, vectors, cells and cell isolates and assays.

**(57) Abrégé**

La présente invention porte sur l'enzyme et les procédures enzymatiques de clivage du site de clivage de la 'beta' secrétase de la protéine APP et des acides nucléiques, des peptides, des vecteurs, des cellules et des isolats cellulaires associés, et sur des dosages.

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12N 15/57, 15/62, 15/85, 5/10, 9/64,</b> <b>C07K 19/00, 14/47, C12N 15/12, C07K</b> <b>16/18, C12Q 1/37, G01N 33/68, C12N</b> <b>1/21</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 00/17369</b> <b>(43) International Publication Date:</b> 30 March 2000 (30.03.00)
<b>(21) International Application Number:</b> PCT/US99/20881 <b>(22) International Filing Date:</b> 23 September 1999 (23.09.99) <b>(30) Priority Data:</b> 60/101,594 24 September 1998 (24.09.98) US <b>(71) Applicant (for all designated States except US):</b> PHARMACIA & UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> GURNEY, Mark, E. [US/US]; 910 Rosewood Avenue, S.E., Grand Rapids, MI 49506 (US). BIENKOWSKI, Michael, Jerome [US/US]; 3431 Hollow Wood, Portage, MI 49024 (US). HEINRIK- SON, Robert, Leroy [US/US]; 81 South Lake Doster Drive, Plainwell, MI 49080 (US). PARODI, Luis, A. [US/SE]; Grevgafan 24, S-115 43 Stockholm (SE). YAN, Riqiang [US/US]; 5025 Queen Victoria Street, Kalamazoo, MI 49009 (US).		<b>(74) Agent:</b> WOOTTON, Thomas, A.; Pharmacia & Upjohn Com- pany, Intellectual Property Legal Services, 301 Henrietta Street, Kalamazoo, MI 49001 (US). <b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GI, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). <b>Published</b> <i>With international search report.</i> <b>(88) Date of publication of the international search report:</b> 23 November 2000 (23.11.00)
<b>(54) Title:</b> ALZHEIMER'S DISEASE SECRETASE		
<b>(57) Abstract</b>		
The present invention provides the enzyme and enzymatic procedures for cleaving the $\beta$ secretase cleavage site of the APP protein and associated nucleic acids, peptides, vectors, cells and cell isolates and assays.		



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

# INTERNATIONAL SEARCH REPORT

Internat. Application No.  
PCT/US 99/20881

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/57 C12N15/62 C12N15/85 C12N5/10 C12N9/64  
C07K19/00 C07K14/47 C12N15/12 C07K16/18 C12Q1/37  
G01N33/68 C12N1/21

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K C12Q G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, STRAND, WPI Data, BIOSIS, CHEM ABS Data, MEDLINE, EMBL

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 848 062 A (SMITHKLINE-BEECHAM CORPORATION) 17 June 1998 (1998-06-17) cited in the application  page 2, line 10 -page 3, line 40 page 4, line 20 - line 33 page 5, line 8 - line 20 page 8, line 1 -page 9, line 25; tables 1,2  -/--	1-3, 5-21,24, 25, 28-31, 34, 37-47, 49-64, 66-69, 72-75, 77, 80-91, 95-97, 114-129, 140,141

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*F\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*S\* document member of the same patent family

Date of the actual completion of the international search

26 July 2000

Date of mailing of the international search report

02.08.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2  
NL - 2280 HV Rijswijk  
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Montero Lopez, B

3

Form PCT/ISA/210 (second sheet) July 1992

# INTERNATIONAL SEARCH REPORT

Intern. Application No.

PCT/US 99/20881

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
X	<p>page 10, line 28 - line 44  page 11, line 10 -page 12, line 8  ---  EP 0 855 444 A (SMITHKLINE-BEECHAM P.L.C.)  29 July 1998 (1998-07-29)  cited in the application</p>	<p>1-3,  5-21,24,  25,  28-31,  34,  37-47,  49-64,  66-69,  72-75,  77,  80-91,  95-97,  114-129,  140,141</p>
X	<p>page 2, line 8 -page 3, line 44  page 5, line 3 - line 15  page 5, line 49 -page 6, line 3; tables  1,2  page 7, line 34 - line 50  page 10, line 20 -page 11, line 1  page 12, line 1 - line 19  page 12, line 45 -page 13, line 44  ---  WO 96 40885 A (ATHENA NEUROSCIENCES)  19 December 1996 (1996-12-19)</p> <p>page 3, line 1 -page 5, line 26  page 8, line 1 - line 34  page 14, line 19 -page 17, line 22  page 23, line 31 -page 25, line 20  page 28, line 7 -page 48, line 13  ---  -/--</p>	<p>1-4,6,7,  9,10,  12-21,  24,25,  28-31,  34,  37-47,  49,50,  52,53,  55-63,  67,68,  72-75,  77,  80-90,  108-129,  136-139,  141</p>

## INTERNATIONAL SEARCH REPORT

Intern. Application No.

PCT/US 99/20881

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 26059 A (ATHENA NEUROSCIENCES, INC.) 18 June 1998 (1998-06-18)	1-4,6,7, 9,10, 12-21, 24,25, 28-31, 34, 37-47, 49,50, 52,53, 55-63, 67,68, 72-75, 77, 80-90, 108-129, 136-139, 141
	page 2, line 35 -page 4, line 3 page 5, line 9 -page 11, line 5 page 11, line 10 -page 22, line 3	
P,X	WO 99 34004 A (CHIRON CORPORATION) 8 July 1999 (1999-07-08)	1-4,6,7, 9-20,24, 28-31, 34, 37-47, 49,50, 52-63, 67,68, 72-75, 77, 80-92, 95-98, 101-103, 106,107, 114-117, 120,141
	page 7, line 19 -page 8, line 9 page 11, line 22 -page 14, line 24 page 16, line 26 -page 21, line 1 page 21, line 20 -page 23, line 13; figure 2; examples 2,3 --- -/--	

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 99/20881

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 99 46281 A (GENENTECH, INC.) 16 September 1999 (1999-09-16)	1-4,6,7, 9-12, 18-20, 24, 28-31, 34,37, 38, 40-47, 49,50, 52-54, 61-63, 67,68, 72-75, 77,80, 81, 84-92, 95-98, 101-103, 106,107, 114-118, 120-128, 140,141
	page 15, line 10 - line 23 page 65, line 5 - line 25 page 130, line 30 - line 35 page 149, line 3 -page 155, line 6 page 160, line 20 - line 22 page 173, line 35 -page 175, line 23; figures 72,73; examples 32,99-107 -----	
A	US 5 795 963 A (MICHAEL JOHN MULLAN) 18 August 1998 (1998-08-18) column 3, line 58 -column 6, line 21 -----	130-135, 141
T	YAN RIQIANG ET AL.: "Membrane-anchored aspartyl protease with Alzheimer's disease beta-secretase activity" NATURE, vol. 402, 2 December 1999 (1999-12-02), pages 533-537, XP002136300 LONDON GB -----	

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 99/20881

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

The International Searching Authority found multiple inventions in this International application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box 1.2

Claims Nos.: claims 32, 33, 35, 36, 70, 71, 76, 78 and 79 and partially claims 1, 18, 28, 44, 61, 72 and 141

Present claims 1, 18, 28, 44, 61, 72 and 141 relate to an extremely large number of possible products. In fact, the claims encompass so many possible compounds that a lack of clarity (and/or conciseness) within the meaning of Article 6 PCT arises to such an extent as to render a meaningful search of the claims impossible.

Moreover, in view of the large number and also the wording of the claims presently on file, which renders it difficult, if not impossible, to determine the matter for which protection is sought, the present application fails to comply with the clarity and conciseness requirements of Article 6 PCT (see also Rule 6.1(a) PCT) to such an extent that a meaningful search is impossible.

In addition, the obscure definition of claims 32, 33, 35, 36, 70, 71, 76, 78 and 79, relating to an unidentified SEQ ID, and referring to the examples renders as well the search of these claims impracticable.

Consequently, the search has been carried out for those parts of the application which do appear to be clear, namely the particular sequences SEQ ID NOs.: 1, 2, 3, 4, 5, 6, and 8, variants, and uses thereof

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-31, 34, 37-69, 72-75, 77, 80-129, 136-140 and partially 141

Proteases capable of cleaving the beta secretase cleavage site of APP, variants thereof; polynucleotides encoding them; vectors and host cells comprising the same; antibodies for the polypeptides and uses of the foregoing in screening tests.

2. Claims: 130-135 and partially 141

APP isoform wherein the last two carboxy terminus amino acids are Lysine residues.



# INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern: al Application No

PCT/US 99/20881

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 848062	A	17-06-1998	JP 11069981 A US 6025180 A	16-03-1999 15-02-2000
EP 855444	A	29-07-1998	CA 2221686 A JP 10327875 A JP 2000060579 A	28-07-1998 15-12-1998 29-02-2000
WO 9640885	A	19-12-1996	US 5744346 A AU 6383396 A EP 0871720 A JP 11507538 T US 5942400 A	28-04-1998 30-12-1996 21-10-1998 06-07-1999 24-08-1999
WO 9826059	A	18-06-1998	AU 1684097 A	03-07-1998
WO 9934004	A	08-07-1999	AU 1726199 A AU 2014899 A WO 9933963 A	19-07-1999 19-07-1999 08-07-1999
WO 9946281	A	16-09-1999	AU 3072199 A AU 3075099 A WO 9947677 A AU 1532499 A WO 9927098 A AU 3757099 A WO 9954467 A AU 1070399 A WO 9920756 A	27-09-1999 11-10-1999 23-09-1999 15-06-1999 03-06-1999 08-11-1999 28-10-1999 10-05-1999 29-04-1999
US 5795963	A	18-08-1998	US 5455169 A	03-10-1995